

CIRCULATION OF MARGINAL AND SEMI-ENCLOSED SEAS (SEA OF JAPAN AND RELATED PROCESS STUDIES)

Christopher N.K. Mooers
Ocean Prediction Experimental Laboratory (OPEL)
Rosenstiel School of Marine and Atmospheric Science
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149-1098
phone: (305) 361-4825 fax: (305) 361-4797 email: cmooers@rsmas.miami.edu
Award #: N000149610399

LONG-TERM GOALS

My basic goal is to understand the general principles governing the mean and transient circulation in realistic marginal and semi-enclosed seas, especially the roles of mesoscale variability. My applied goal is to improve the scientific basis for the design of nowcast/forecast systems to be implemented in marginal and semi-enclosed seas. I am particularly interested in treating realistic continental margins and realistic atmospheric and throughflow forcing for such seas, using numerical simulation and data assimilation models for the circulation.

OBJECTIVES

My scientific objectives include:

- 1) developing methods for dealing with open boundary forcing of marginal and semi-enclosed seas due to the large-scale external circulation;
- 2) developing methods for model validation; i.e., for quantitative validation of models versus observations;
- 3) conducting model versus observations comparisons;
- 4) conducting model versus model comparisons; and
- 5) conducting observing system design studies through numerical simulation.

APPROACH

The Princeton Ocean Model (POM), as implemented for the Sea of Japan (SOJ) with mesoscale eddy-admitting resolution, is used for sensitivity and process studies. Model simulations are evaluated in comparison to CREAMS current meter time series. Ms. HeeSook Kang carries out these calculations for her graduate research project. Dr. Jia Wang provides technical advice and is responsible for the POM implementation in Prince William Sound and the Labrador Sea, where an ice coupling strategy is being developed. Ms. Renellys C. Perez has been making model versus observations comparisons in the Gulf of Mexico and will soon shift to data assimilation studies with SOJ-POM. Mr. Derrick Snowden is beginning to implement a model for Prince William Sound that has an open boundary well offshore in the Gulf of Alaska to explore use of the NCEP Pacific Ocean Re-Analysis for open boundary conditions. Prof. Victor I. Kuzin, RAS (Akademgorodok), has implemented a finite element model for SOJ so that model-model comparisons can soon begin.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Circulation of Marginal and Semi-enclosed Seas (Sea of Japan and Related Process Studies)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami,Rosensteil School of Marine and Atmospheric Science,4600 Rickenbacker Causeway,Miami,FL,33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED

Our SOJ-POM implementation has been used to conduct sensitivity studies relative to throughflow strength and structure, different climatological atmospheric forcing fields, and increased vertical resolution. It has also been used to make statistical comparisons to CREAMS current meter data, prior to introducing seasonally varying and synoptic atmospheric forcing. And it has been used to perform a case study for the circulation and convective response to impulsive forcing (windstress, heat flux, and moisture flux) characteristic of Siberian cold air outbreaks.

RESULTS

With ca. 10km horizontal resolution, a 26 (sigma) level simulation (with relatively high resolution in surface and bottom boundary layers) yielded results superior to 15 and 20 level simulations. The major circulation feature of the SOJ is the cyclonic gyre over the Japan Basin, and which is the locus of strong mesoscale variability and topographic coupling of barotropic and baroclinic modes, as can be seen from the 1,000-day mean and standard deviation fields for the sea surface height (SSH) and depth-integrated stream-function (ISF) (Fig. 1). Comparison of the 26-level simulation (with steady forcing) versus CREAMS intermediate level current meter data over the Japan Basin for 1,000-days indicated close agreement of the spectral levels and slopes for the oceanic mesoscale frequency band but (understandably) two orders of magnitude lower energy levels in the atmospheric synoptic frequency band (Fig. 2). However, the model mean flows were substantially greater than the observed mean flows for unknown reasons. The detailed structure of the upper layer circulation is also quite sensitive to the forcing and vertical resolution.

IMPACT/APPLICATIONS

A well-validated modeling system for marginal and semi-enclosed seas in general, and the Sea of Japan in particular, will form the basis for an ocean (circulation) prediction system. As such, it will form the basis for observing system design studies and hypothesis development. It will also form the basis for marine ecosystem, sediment transport, etc. simulation models.

TRANSITIONS

The results will aid in the final design of the CREAMS II field experiment. NRL (SSC) NAVO, and FNMOC are briefed annually on progress and plans.

RELATED PROJECTS

POM is also being used in simulations of the circulation of Prince William Sound (PWS) and the Intra-Americas Sea (IAS). PWS is a much smaller domain than SOJ, while IAS is significantly larger. Both PWS and IAS, as SOJ, are strongly influenced by throughflow and atmospheric forcing. PWS, unlike SOJ and IAS, is also strongly influenced by tidal forcing. SOJ and PWS, unlike IAS, are significantly influenced by wintertime ventilation and occasionally deep convection. Thus, these three domains have important dynamical similarities and differences, allowing exploration of general principles in different parts of parameter space. The Dynalysis of Princeton POM implementation for the Gulf of Mexico is the subject of an ongoing evaluation in comparison to the copious LATEX datasets.

REFERENCES

- (1994) (with D.S. Ko) Nowcast System Development for the Straits of Florida. In: (Eds. M.L. Spaulding, K. Bedford, A. Blumberg, R. Cheng, and C. Swanson), *Estuarine and Coastal Modeling III*, ASCE, pp. 158-171.
- (1994) (with S. Jin and D.S. Ko) Preliminary Evaluation of a Gulf of Mexico Circulation Model at 92W. VII JECCS/PAMS Workshop Proceedings, *La Mer*, 32 (4): 307-324.
- (1995) (with H.S. Kang) Initial Spin-Up of a Sea of Japan Numerical Circulation Model. In: Advanced Mathematics: Computations and Applications (Eds. A.S. Alekseev and N.S. Bakhvalov), NCC Publisher, Novosibirsk, pp 350-357.
- (1996) (with H.S. Kang) Preliminary Results from a Sea of Japan Numerical Circulation Model. Proceedings of the PICES Workshop on the Sea of Okhotsk and Adjacent Areas, PICES Scientific Report No. 6, pp. 202-214.
- (1996) (with J. Wang) Three-Dimensional Perspectives of the Florida Current: Transport, Potential Vorticity, and Related Dynamical Properties. *Dyn. Atmos. and Ocn.* (in press).
- (1997) (with G. Maul) Intra-Americas Sea Coastal Ocean Circulation. In: Global Coastal Ocean (Eds. A.R. Robinson and K.H. Brink), *The Sea*, John Wiley & Sons, NY (in press).
- (1997) (with J. Wang) On the Development of a Three-Dimensional Circulation Model for Prince William Sound, Alaska. Con. Shelf Res. (in press).
- (1997) (with E. Deleersnijder and J. Wang) A Two-compartment Model for Understanding the Simulated Three-Dimensional Circulation in Prince William Sound, Alaska. Con. Shelf Res. (in press).
- (1997) (with J.Wang and V. Patrick) A Three-Dimensional Tidal Model for Prince William Sound, Alaska. In: Computer Modelling of Seas and Coastal Regions III (eds. J.R. Acinas and C.A. Brebbia), Computational Mechanics Publications, Southampton, pp. 95-104.
- (1997) H.S. Kang Implementation, Sensitivity Testing, and Evaluation of a Numerical Model for the East/Japan Sea Circulation, MS Thesis, RSMAS/University of Miami, 174pp.




Figure. 1 Mean and standard deviation of sea surface height (SSH) and depth-integrated streamfunction (ISF).





Figure. 2

Frequency-domain comparison of model output (red) versus CREAMS (Prof. M. Takematsu) current meter data (black) at 2,000 and 3,000 m in Japan Basin.