

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Ionospheric Profiles from Ultraviolet Remote Sensing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Alabama A & M University, Department of Physics, P. O. Box 447, Normal, AL, 35762				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

IONOSPHERIC PROFILES FROM ULTRAVIOLET REMOTE SENSING

Arjun Tan

Department of Physics

Alabama A & M University

P. O. Box 447

Normal, AL 35762

phone: (205) 851-5313, fax: (205) 851-5622, e-mail: atan@asnaam.aamu.edu

Award #: N00014-97-1-0267

LONG-TERM GOAL

The long-term goal of this project is to obtain ionospheric profiles from ultraviolet remote sensing of the ionosphere from orbiting space platforms.

SCIENTIFIC OBJECTIVES

Remote sensing of the nighttime ionosphere is a more straightforward process because of the absence of the complications brought about by daytime solar ultraviolet radiation. This, therefore, constitutes the first objective of this investigation.

APPROACH

We first focus on the reconstruction of the topside ionosphere where the major ions are O^+ and H^+ . The H^+ densities are expressed in terms of O^+ densities by assuming a chemical equilibrium condition (Hanson and Ortenburger, 1961). The intensity of the O^+e^- recombination reaction at different observation angles is then a linear combination of the altitudinal distribution of $[H][O^+]^2/[O]$. The latter can be retrieved by matrix inversion from which the densities of O^+ , H^+ , and e^- can be calculated sequentially.

WORK COMPLETED

A computer code to calculate the path matrix (path lengths through stratified layers at various observation angles) given the altitude of the orbiting platform, has been written. The forward model to calculate the expected nighttime emission intensities from the O^+e^- recombination as seen from the orbiting platform has been completed.

A matrix inversion program in double precision BASIC has been incorporated. It has successfully inverted path matrices of dimensions of up to at least 29x29. Standard Chapman profiles have been reconstructed using the matrix inversion routine which are accurate up to about 1000 km.

RESULTS

The nighttime intensity of the radiative recombination of O^+e^- at 91.1 nm as observed from 833 km altitude is calculated using the forward model. The reaction rate is assumed to be $4.4 \times 10^{-13} \text{ cm}^3/\text{s}$ (Dymond, et al., 1995). The standard Chapman profile is given by 3 parameters: peak electron density $N_m = 10^6 \text{ cm}^{-3}$; altitude of the peak $z_m = 350 \text{ km}$; and atomic oxygen scale height $H = 60 \text{ km}$.

Parametric study of the intensity of radiation in z_m , H and N_m are obtained for nadir angles (complements of observation angles) between 66 deg and 73 deg. The peak intensity nadir angle

translates about 1.1 deg for every 50 km increase of z_m . There is a slight but perceptible increase in the peak intensity with z_m . The peak migrates to lower nadir angles with increasing H .

IMPACT/APPLICATION

Existing remote sensing methods for ionospheric profiles do not take into account the H^+ densities. Faithful reconstruction of the far topside ionosphere is, therefore, not feasible at present. The current study aims to rectify this situation.

TRANSITIONS

Several current and future ultraviolet remote sensing experiments to be conducted by the U. S. Naval Research Laboratory will monitor the ionosphere on a continuous basis from orbiting space platforms. The present study can find applications in these measurements.

REFERENCES

- K. F. Dymond, S. E. Thonard, and R. P. McCoy, 1996. *I. E. S. Alexandria, Virginia, May.*
W. B. Hanson and I. B. Ortenburger, 1961. *J. Geophys. Res.*, 66, 1424 .