

SIMULATIONS OF ATMOSPHERIC FLOWS IN THE BOUNDARY LAYER OVER INHOMOGENEOUS SURFACE CONDITIONS

Darko R. Koracin

Desert Research Institute, 5625 Fox Ave., Reno, NV 89506
702-677-3344 (voice); 702-677-3157 (fax); e-mail: darko@sage.dri.edu

Award # N00014-96-1-1235

LONG TERM GOALS

The goal of this project is to improve the description and prediction of resolvable and subgrid-scale atmospheric fields over inhomogeneous surfaces and to develop an integrated modeling system for weather phenomena and dispersion effects on a spectrum of spatial and temporal scales.

OBJECTIVES

Specific objectives include: 1) developing a fully-compressible and computationally efficient model that can be applied to atmospheric and dispersion fields on microscale to mesoscale domains, 2) investigating turbulence transfer within the atmospheric boundary layer, and 3) formulating a concept of nonlocal turbulent mixing. This work is supported by the Office of Naval Research, Marine Meteorology and Atmospheric Effects.

APPROACH

We have developed a fully-compressible, 2D atmospheric model based on the MacCormack numerical scheme, a subgrid-scale turbulence transfer algorithm, and an interactive shortwave radiation scheme based on the Monte Carlo method (Koraćin et al. 1997a, website 1). To simulate atmospheric transport and dispersion, we developed a Lagrangian particle model (LAP, Koraćin et al. 1997d, websites 3-5) which uses data from measurements or atmospheric model results as input. Our approach to the LAP model involved a new treatment of the range of randomness derived from the turbulence kinetic energy. In order to gain confidence in dispersion estimates, we also developed a novel approach to evaluating predicted atmospheric fields in complex terrain by using tracer measurements (Koraćin et al. 1997c).

We have also developed a new approach to nonlocal mixing applicable to different models. Our method uses Donaldson's set of so called "super-equilibrium equations" for initial fields and then determines nonlocal mixing potential from an integrated value of the turbulence kinetic energy and momentum fluxes (Koraćin and Isakov 1997; Koraćin et al. 1997b, website 2).

WORK COMPLETED

Work involved the reproduction of several published numerical experiments appropriate to fully-

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997	2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997		
4. TITLE AND SUBTITLE Simulations of Atmospheric Flows in the Boundary Layer over Inhomogeneous Surface Conditions			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Desert Research Institute, 5625 Fox Ave, Reno, NV, 89506			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

compressible models. A simulation of a buoyant bubble in a neutral environment closely correlated with results from Droegemeier's fully compressible model. The same experiment using a compressible Advanced Regional Prediction System (ARPS) led to similar results. Our model, including the Monte Carlo radiation scheme, was also evaluated using Nicholls' (1984) measurements over the North Sea. The model evaluation and analysis of simulation results were presented by Koraćin et al. (1997a).

The LAP model was applied to data from a tracer experiment conducted in complex terrain (Koraćin et al. 1997d, website 4). Atmospheric fields of various complexities were evaluated by using the Tracer Potential method (Koraćin et al. 1997c) prior to being used as input to the dispersion model. This method for determining nonlocal mixing was applied to an idealized convective experiment as well as to data from the first Lagrangian period of the Atlantic Stratocumulus Transition Experiment (ASTEX) (Koraćin and Isakov 1997; Koraćin et al. 1997b, website 2; Tjernström et al. 1997).

RESULTS

The destruction of a "hot" buoyant bubble was simulated by the recently developed fully-compressible atmospheric model (Fig. 1) (Koraćin et al. 1997a, website 1). The model retained computational stability while simulating vertical motions ranging between -11 ms^{-1} and 22 ms^{-1} . A numerical experiment with a broken-cloud field indicated several important issues in using the Monte Carlo scheme in mesoscale and cloud-resolving models: accuracy was determined by the number of photons used in the experiment (at least 10,000 photons were found to be necessary to provide 2% accuracy in flux calculations); the most critical parameter in shortwave flux calculations was found to be cloud optical thickness; and geometry of cloud elements was also important.




Figure 1 Contour plot of potential temperature (K) representing positively buoyant bubble in neutral environment.




Figure 2 Measured (dashed) and simulated by LAP (solid line) contours of SF_6 surface tracer concentrations in the southwest U.S. averaged from 6-14 August 1992.




Figure 3 Non-diagonal coefficients of the transilient matrix describing turbulent transfer.

The LAP model reproduced the main features of measured tracer concentrations in complex terrain at distances from the source ranging from 20 to 300 km, including channeling of the transported tracers (Fig. 2). The LAP model simulated the correct order of magnitude of tracer concentration and reproduced similar temporal patterns as measurements.

According to preliminary results of the method for determining nonlocal turbulent transfer, nonlocal mixing contributed to evolution of the boundary layer by approximately 10% as compared to local mixing in the stratocumulus case (Fig. 3). The contribution increased to approximately 30-40% in the convective test case (Koraćin and Isakov 1997; Koraćin et al. 1997b, website 2).

IMPACT

The models and methods we have developed will increase understanding and lead to improved prediction of the atmospheric boundary layer, particularly in the areas of turbulence transfer and atmospheric transport and dispersion. This has obvious application to naval and aircraft operations as well as to defense against chemical or biological weapons. Most of the models can be operated on a PC platform, permitting use by tactical military units.

TRANSITIONS

The cloud-resolving numerical model will be adapted and available for operational applications. A research team from Flinders University in Australia would like to obtain the code and use it for studies of broken clouds off the Australian coast. Faculty at the University of Zagreb in Croatia would like to use the model for air quality studies in complex terrain and along the Adriatic coast. A team from the University of Uppsala in Sweden plans to use our method of nonlocal mixing in a collaborative study of the ASTEX data. The Naval Postgraduate School in Monterey is interested in obtaining high resolution atmospheric data as input for ocean models.

RELATED PROJECTS

Work on this project benefitted from experience and results from another ONR-funded project (N00014-96-1-0980) focused on simulations of coastal dynamics. Darko Koraćin is also a P.I. on this project. Performance has been enhanced by collaboration with Dr. Luis Mendez of the University of California, Davis, and Dr. Michael Tjernström of the University of Uppsala, Sweden. Research conducted in this project has led to submission of a proposal to DOD-ONR focused on modeling the dispersion of vapor and aerosol particulates in complex terrain as well as a proposal (with NCEP) to NOAA to develop a high resolution, compressible ETA model. The Tracer Potential

method and Lagrangian model were also used in several EPA-related projects (website 4) as well as in planning a modeling study for a future airport near Las Vegas (website 3) and the NOAA-funded weather modification program (website 5).

REFERENCES

Koračin, D. and V. Isakov, 1997: Simulations of local and nonlocal instabilities in marine stratocumulus. IAMAS and IAPSO, 1 - 9 July 1997, Melbourne, Australia. Paper IM5pp.

Koračin, D., V. Isakov and L. Mendez-Nunez, 1997: A Cloud-resolving Model with the Radiation Scheme Based on the Monte Carlo Method. *Atmos. Res.(in print)*

Koračin, D., M. Tjernström and G. Svensson, 1997: Turbulence transfer and nonlocal mixing processes in the cloudy marine atmospheric boundary layer. 12th Symposium on Boundary Layers and Turbulence, 28 July - 1 August 1997, Vancouver, B.C. p.60-61.

Koračin, D., V. Isakov and J. Frye, 1997: A method of evaluation of atmospheric and dispersion models by using tracer measurements. To be presented at the 10th Joint Conference on the Applications of Air Pollution Meteorology, Phoenix, AZ, 11-16 January 1998. Pap. 1.3, 6 pp.

Koračin, D., V. Isakov and J. Frye, 1997: A Lagrangian particle dispersion model (LAP) applied to transport and dispersion of chemical tracers in complex terrain. To be presented at the 10th Joint Conference on the Applications of Air Pollution Meteorology, Phoenix, AZ, 11-16 January 1998. Paper 5B.5, 4 pp.

Tjernström, M., D. Koračin and G. Svensson, 1997: A numerical investigation of the First ASTEX Lagrangian. IAMAS and IAPSO, 1 - 9 July 1997, Melbourne, Australia. Paper IM6kk.

Website 1: <http://www.dri.edu/ResearchAreas/Modeling/newmod.html>

Website 2: <http://www.dri.edu/ResearchAreas/Modeling/flows.html>

Website 3: <http://www.dri.edu/ResearchAreas/Modeling/vegasair.html>

Website 4: <http://www.dri.edu/ResearchAreas/Modeling/mohave.html>

Website 5: <http://www.dri.edu/ResearchAreas/Modeling/wmod.html>