

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 03/11/2015		2. REPORT TYPE Conference Presentation		3. DATES COVERED (From - To) 2014-2015	
4. TITLE AND SUBTITLE Investigation into Blue Light Emission for Copper-containing Pyrotechnics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) David Dye, Jonathan Dilger, Christina Yamamoto				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Surface Warfare Center, Crane Division Code WXP, Building 3323 300 Highway 361 Crane, IN 47522-5001				8. PERFORMING ORGANIZATION REPORT NUMBER NSWCCR/RDTR-15/56	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Innovative Science and Engineering Program				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Pyrotechnic signals typically generate colored light via photonic emission from thermally-excited states of gas-phase metal combustion products. Blue light can be generated in pyrotechnic compositions with the use of copper-containing compounds as a fuel within the formulation. In the presence of chlorine containing compounds within the composition, these pyrotechnics exhibit emission bands in the blue region of the visible spectrum with peaks observed from 435-480 nm. Color degradation occurs at temperatures above 1200 °C, leading to the hypothesis that the blue emission must arise from a relatively unstable molecular species. Several have hypothesized that this species is likely copper(I) chloride (CuCl) or even the trimeric Cu ₃ Cl ₃ , though the identity of this emissive species as either CuCl or Cu ₃ Cl ₃ has been theoretically disputed in the literature with NASA CEA thermodynamic calculations suggesting that excellent blue colors can be achieved with temperatures in excess of 2200 °C. While many of these potential emissive copper-containing species have been empirically characterized, the identity and electronic structure of these copper(I) halide emitters have yet to be fully characterized. Here, we investigate these cuprous halide emitters with the use of flame ionization spectroscopy to measure high-resolution molecular emission spectra of various cupric salts that can be utilized in pyrotechnic signaling devices. These spectroscopically measured visible emission bands will be compared with calculated electronic transitions of candidate emitter species using time-dependent density functional theory to confirm the molecular and electronic identity of the emitters. These techniques will establish methods for the characterization and development of light emitters for next generation visible-light generating pyrotechnic formulations.					
15. SUBJECT TERMS pyrotechnics, copper halides, blue visible light, density functional theory					
16. SECURITY CLASSIFICATION OF: U			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 17	19a. NAME OF RESPONSIBLE PERSON David Dye
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code) 1-812-854-6616

Investigation into Blue Light Emission for Copper-containing Pyrotechnics

Dr. David Dye*, Dr. Jonathan Dilger*, Christina Yamamoto
Naval Surface Warfare Center, Crane Division
Code WGRP, Building 3323
300 Highway 361
Crane, IN 47522-5001
Phone: 1-812-854-1452
email: david.f.dye@navy.mil

*These authors contributed equally

Pyrotechnic signals typically generate colored light via photonic emission from thermally-excited states of gas-phase metal combustion products. Blue light can be generated in pyrotechnic compositions with the use of copper-containing compounds as a fuel within the formulation. In the presence of chlorine-containing compounds within the composition, these pyrotechnics exhibit emission bands in the blue region of the visible spectrum with peaks observed from 435-480 nm. Color degradation occurs at temperatures above 1200 °C, leading to the hypothesis that the blue emission must arise from a relatively unstable molecular species. Several have hypothesized that this species is likely copper(I) chloride (CuCl) or even the trimeric Cu_3Cl_3 , though the identity of this emissive species as either CuCl or Cu_3Cl_3 has been theoretically disputed in the literature with NASA CEA thermodynamic calculations suggesting that excellent blue colors can be achieved with temperatures in excess of 2200 °C. While many of these potential emissive copper-containing species have been empirically characterized, the identity and electronic structure of these copper(I) halide emitters have yet to be fully characterized. Here, we investigate these cuprous halide emitters with the use of flame ionization spectroscopy to measure high-resolution molecular emission spectra of various cupric salts that can be utilized in pyrotechnic signaling devices. These spectroscopically measured visible emission bands will be compared with calculated electronic transitions of candidate emitter species using time-dependent density functional theory to confirm the molecular and electronic identity of the emitters. These techniques will establish methods for the characterization and development of light emitters for next-generation visible-light generating pyrotechnic formulations.



HARNESSING TECHNOLOGY for the WARFIGHTER

*Captain J. T. Elder
Commanding Officer
NSWC Crane*

Investigation into Blue Light Emission for Copper-Containing Pyrotechnics

POC: Dr. David F. Dye, david.f.dye@navy.mil

***Prepared for: EuroPyro
May, 2015, Toulouse, France***

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)





Blue Pyrotechnic Emitters: Background

- Blue Pyrotechnic Emitters:
 - Copper Chloride
 - Traditionally identified as CuCl monomer
 - Some dispute¹: clusters detected via mass spec
 - Clusters know to be photoluminescent²
 - Copper Iodide
 - More intense than CuCl³
 - No clear explanation as to why



1) Dolata, Propellants, Explosives, Pyrotechnics, 30 (2005), No 1, p 63.

2) Ford, Chemical Reviews, 99, (1999), pp 3625-3647.

3) Klapötke, Antewandte Chemie, 53, (2014), 9665 –9668.






- Quantum mechanics can elucidate differences in molecular emission
- Computational tool: Time-Resolved Density Functional Theory (TD-DFT)
 - Gaussian 09W, B3LYP, Def2-TZPPD¹
- Calculations will focus on electronic transitions
 - Influence of molecular vibrations—Franck-Condon factors



¹Dmitrij Rappoport and F. Furche, *Property-optimized Gaussian basis sets for molecular response calculations*, *J. Chem. Phys.* 133, 134105 (2010).



Computational Results: $(\text{CuCl})_n$ Geometries



CuCl Calculated Geometry

Cu-Cl	2.091 Å
-------	---------

$(\text{CuCl})_2$ Calculated Geometry

Cu-Cl	2.281 Å
Cu-Cu	2.441 Å
Cu-Cl-Cu	64.7°
Cl-Cu-Cl	115.3°

$(\text{CuCl})_3$ Calculated Geometry




Cu-Cl	2.199 Å
Cu-Cu	2.671 Å
Cu-Cl-Cu	74.8°
Cl-Cu-Cl	165.2°
Cu-Cu-Cu	60.0°

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)





Computational Results: $(\text{CuI})_n$ Geometries



CuI Calculated Geometry

Cu-I	2.393 Å
------	---------

$(\text{CuI})_2$ Calculated* Geometry

Cu-I	2.565 Å
Cu-Cu	2.455 Å
Cu-I-Cu	57.2°
I-Cu-I	122.8°

$(\text{CuI})_3$ Calculated* Geometry

Cu-I	2.548 Å
Cu-Cu	2.252 Å
Cu-I-Cu	52.4°
I-Cu-I	63.8°
Cu-Cu-Cu	60.0°


*Preliminary optimization using LANL2DZ

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)






Computational Results: CuCl Monomer



$$\nu_{1(\text{calc})} = 391.39 \text{ cm}^{-1}$$

- Single vibration
- Two dominant visible electronic absorption bands




DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)






Computational Results: CuI Monomer



$$\nu_{1(\text{calc})} = 391.39 \text{ cm}^{-1}$$

- Single vibration
- Two dominant visible electronic absorption bands
- Very similar to CuCl




DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)




- SrCl calculated as reference
 - $\lambda_{1(calc)} = 647.28$ nm
- Cluster growth produces hypsochromic absorption shift
- “Low” energy of first electronic transition for CuX monomers of interest

Calculated (CuCl)_n Absorption Spectra




Calculated (CuI)_n Absorption Spectra





Preliminary Computational Results: Emission Energies



Electronic Excitation vs. Emission Energies		
	Emission	Absorption
SrCl	648.4 nm	647.28 nm
CuCl	505.9 nm	482.99 nm
CuI	616.5 nm	528.41 nm

SrCl State Energies	
GS1	0 eV
ES1	1.9155 eV
ES2	1.9138 eV
GS2	0.0017 eV
Emission	1.9121 eV
Emission	648.4 nm


CuCl State Energies	
GS1	0 eV
ES1	2.5670 eV
ES2	2.5278 eV
GS2	0.0762 eV
Emission	2.4506 eV
Emission	505.9 nm

CuI State Energies	
GS1	0 eV
ES1	2.3463 eV
ES2	2.2256 eV
GS2	0.2145 eV
Emission	2.0111 eV
Emission	616.5 nm

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)




- Optimization of excited state geometries is ongoing
 - ES electronic structure causes distortion
 - Confirmation of minimized structures via vibrational calculations
 - This geometry is NOT minimized!
 - Software can sometimes produce misleading results if you aren't careful



Calculated Geometry	
Cu-Cl ₁	2.168 Å
Cu-Cl ₂	2.558 Å
Cu-Cu	3.518 Å
Cu-Cl-Cu	77.3°
Cl-Cu-Cl ₁	108.5°
Cl-Cu-Cl ₂	86.9°



Computational Results: Trimeric Species



DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)



Flame Emission Spectroscopy



- Metered gases to control flame temperature
- High-resolution spectroscopic measurements



salts of arsenic, copper, sodium, lithium, and calcium burn within a flame

<http://www.somethingaboutscience.com/?p=1877>

Concentrated metal salt solutions nebulized within premixed gas fuel



<http://faculty.sdmiramar.edu/fgarces/labmatters/instruments/aa/aa.htm>



Conclusions

- Calculations of electronic emission spectra for $(\text{CuX})_n$ species are underway
 - Preliminary calculations show all species have electronic excitation energies in correct range for possible blue emission
 - Vibrational contributions to emission spectra will be valuable for determining emitting species
- Future flame ionization studies will be used to for comparison with computational results

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)





Acknowledgements

- Dr. Aurora Clark, Washington State University
- Dr. Jonathan Dilger, NSWC Crane
- Dr. Eric Miklaszewski, NSWC Crane
- Dr. Hank Webster, NSWC Crane
- Joshua Geary, NSWC Crane

- Funding: NSWC Crane NISE/219 program

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)






Backup Slides

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)







Preliminary Computational Results




Copper Iodide Monomer (CuI)




Copper Iodide Dimer (CuI)₂




(CuI)₃ – 146.2 cm⁻¹ vibration



(CuI)₃ – 278.6 cm⁻¹ vibration



CuCl Calculated Electronic Absorption Spectrum



(CuI)₃ Calculated Vibrational Spectrum

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited (All data should be considered preliminary and is subject to change)

