

Characterization of Caribbean Meso-Scale Eddies

Jose M. Lopez
Department of Marine Sciences
P.O. Box 9013
Mayaguez, Puerto Rico 00681-9013
phone: (787) 834-7620 fax: (787) 834-8025 email: jlopez@uprm.edu

Jorge E. Corredor
Department of Marine Sciences
P.O. Box 908
Lajas, Puerto Rico 00667-908
phone: (787) 899-2049 fax: (787) 899-3838 email: j_corredor@cima.uprm.edu

Julio M. Morell
Department of Marine Sciences
P.O. Box 908
Lajas, Puerto Rico 00667-908
phone: (787) 899-2049 fax: (787) 899-3838 email: j_morell@cima.uprm.edu

Jorge E. Capella
Department of Marine Sciences
P.O. Box 908
Lajas, Puerto Rico 00667-908
phone: (787) 899-2049 fax: (787) 899-3838 email: jcapella@caribe.net

Fernando Gilbes
Department of Geology
P.O. Box 9001
Mayaguez, Puerto Rico 00681-9001
phone: (787) 834-7620 fax: (787) 265-6340 email: gilbes@cacique.uprm.edu

Grant Number: N000140310904

LONG-TERM GOALS

Our long-term goal is to improve predictivity of physical, biogeochemical and optical properties of Eastern Caribbean waters under the influence of mesoscale eddies and their interaction with regional features (i.e. massive riverine discharge). An expected outcome from this research is the capacity to infer subsurface properties and processes as well as their temporal and spatial evolution utilizing remotely acquired surface observations.

This program will contribute to the development of infrastructure for marine research and education at the University of Puerto Rico. A particular expectation is the recruitment and training of graduate students who will focus their graduate research in current scientific issues attuned with the above expressed goals.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Characterization of Caribbean Meso-Scale Eddies				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Marine Sciences,,P.O. Box 9013,,Mayaguez,,PR, 00681				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Our long-term goal is to improve predictivity of physical, biogeochemical and optical properties of Eastern Caribbean waters under the influence of mesoscale eddies and their interaction with regional features (i.e. massive riverine discharge). An expected outcome from this research is the capacity to infer subsurface properties and processes as well as their temporal and spatial evolution utilizing remotely acquired surface observations.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

SPECIFIC OBJECTIVES

- To determine the distribution of physical, chemical, biological and optical variables across frontal and eddy boundaries in the Eastern Caribbean Sea
- To develop sub regional derivation of empirical relationships between horizontal and vertical physical/chemical gradients and remotely sensed ocean properties
- Verification of inherent optical properties predicted by NRL's Automated Software System
- Verification of the eddy structure and trajectory predictions by the 1/16° operational global NRL Layered Ocean Model
- Correlations phytoplankton pigment composition and size class distribution to inherent optical properties

APPROACH

We are implementing a series of research cruises to carry out observations of the optical and physical structure and upper water column biogeochemistry of eddies. Eddy distribution and displacement will be followed using the output from 1/16° operational global Naval Research Laboratory (NRL) Layered Ocean Model (NLOM), near real time altimetry data, SeaWiFS and MODIS imagery (F.Gilbes). During research cruises we will obtain continuous flow surface records of T, S and Chl-a (J. Corredor), vertical sections of these variables plus apparent optical properties using the Nv-shuttle undulating towed body (J. Lopez). Discrete vertical profiles (J. Morell, J. Corredor) of physical (T, S), chemical (DO, nutrients, CDOM), biological and inherent (a, b, c) and apparent (R, R_{RS} , T, K_z) optical properties (F. Gilbes) will be obtained. Selected samples will be subjected to 3-D spectrofluorometry and scanning spectrophotometry for CDOM characterization (J. Corredor, J. Morell). Current structure across fronts and eddies will be characterized by means of ship-lowered ADCP current profiler (J. Capella). These studies will also include assessment of rates of biomass accumulation, primary production and photosynthetic efficiency (J. Lopez).

WORK COMPLETED

Cruise CaVortEx I aboard R/V CHAPMAN, August 12 – 19 2003 was undertaken in order to characterize a cyclonic eddy detected by satellite altimetry and ocean color traversing the east-central Caribbean Sea. The eddy, approximately 230 km in diameter was, observed as a sea surface height (SSH) anomaly by satellite altimetry in July 2003. Subsequently, SeaWiFS ocean color imagery revealed that the eddy core, at the time centered ca. latitude 14.8 ° N and longitude 067.8 ° W, was surrounded by a high-chlorophyll (HC) surface water mass. Imagery showed this as a large filament continuous with the Orinoco/Amazon River plume, creating a spiral pattern of alternating bands of HC and LC (low-chlorophyll) surface water with HC waters occupying the entire periphery of the eddy enclosing a core of LC waters. The cruise track, a diametric north-south section across the eddy with observations extending to 1000 m depth, allowed characterization of the physical, chemical, optical and biogeochemical structure of the eddy.

RESULTS

Observations during the initial field campaign CaVortEx I confirmed the applicability of ocean color imagery to determine the precise location of eddies, particularly when these interact with Chlorophyll/CDOM-rich river plume waters (Figure 1). Eddy core position estimates based on ocean altimetry were displaced to the east (the oncoming direction) of the actual position probably due to data smoothing procedures.




Figure 1: Contour plot of MODIS Chl-a data product for 14 August 2003
[Oceanographic stations and dates occupied aboard R/V CHAPMAN during the Caribbean Vorticity Experiment (CaVortEx I) are superimposed as white diamonds. The eddy core in the image is located approximately 14° 50' N 067° 50' W. Higher-chlorophyll, near-surface river plume water (reds and greens) surrounds lower-chlorophyll near-surface waters of the eddy core (blue). A filament of plume water spirals within the eddy core. Real-time shipboard salinity and chlorophyll-a measurements across the track (inset) closely reflect the pattern observed in the satellite imagery, yet surface Chlorophyll-a concentrations, calculated with the MODIS algorithm, exceed those observed in-situ by a factor of ca. 2.5.]

The hydrography evidenced moderate doming of the isopycnals down to ca. 800m with no evident near surface cross-isopycnal mixing. The salinity section evidences interaction of the eddy with the Orinoco/Amazon river plume. Contrary to purely oligotrophic oceanic eddies, the eddy-plume interaction resulted in the eddy core area exhibiting relatively higher upper water transparency than in non-eddy areas thus the presence of a deeper Chl a maximum.




Figure 2: North South oceanographic section of the cyclonic eddy.

[Left: Baroclinic structure of the eddy is apparent in the seawater density (sigma-t) section as a dome extending throughout the top of the water column to ca. 700 m, below which, isopycnals exhibit a negative dip. Center: High-salinity oceanic waters interact with the low-salinity plume water in the upper portion of the dome.

Right: As the subsurface influence of the northern edge of the eddy dome extends beyond the surrounding surface river filament, high chlorophylls in the buoyant plume create a halo around the doming, deep chlorophyll maximum (DCM) of the eddy. Low light availability in waters directly below the plume limits the abundance of phytoplankton and results in attenuation of the oceanic DCM.]

IMPACT/APPLICATIONS

Cyclonic eddies have been invoked as powerful drivers of biogeochemical fluxes in the tropical ocean as well as significant vectors for inter-hemispheric transport. Such roles have aroused interest from scientists in a wide range of fields. The capability to remotely characterize these features as well as to predict their properties and dynamics should allow for more precise modelling of the role these mesoscale features play in the global ocean. In a regional context, the occurrence of these mesoscale events further contributes to the optical and biogeochemical complexity of Eastern Caribbean waters where other mesoscale phenomena, the discharge of two of the world's major rivers and coastal upwelling, sustain pronounced physical chemical and biological gradients. Knowledge of the interaction of these features with eddies will allow a more complete understanding of the optical heterogeneity of the region and its forecasting.

RELATED PROJECTS

A UPRM/NASA-sponsored research program, entitled Tropical Center for Earth and Space Studies, includes an oceanography component focused on identifying factors limiting biological carbon fixation in the Western Tropical Atlantic and Caribbean Sea. This program has funded the acquisition of instrumentation applicable to the effort here described and provided ocean color satellite imagery and ship time funding for the CaVortEx I cruise.