

Lagrangian Floats For Deep Convection

Eric A. D'Asaro

Applied Physics Laboratory, University of Washington

1013 NE 40th Street, Seattle, WA 98105

phone: 206 685 2982 fax: 206 543 6785 email: dasaro@apl.washington.edu

N000149410025

<http://poseidon.apl.washington.edu/~dasaro/Labsea/>

LONG TERM GOALS

I aim to understand the process of deep convection in the ocean.

OBJECTIVES

Near surface water is mixed to great depth at a few high latitude locations, thereby forming the deep and bottom masses of the ocean. This proposal has supported the development and deployment of neutrally buoyant floats in the Labrador Sea in the winters of 1997 and 1998 and the analysis of the resulting data. These floats provide detailed information on the processes and rates of deep convection.

APPROACH



Lagrangian Floats (see figure) accurately follow water motions through a combination of a density which matches that of seawater and a high drag. The density is matched to that of the ambient water by actively changing the float's volume and will stay matched, despite changes in pressure and temperature, through a combination of active control and a hull compressibility which is close to that of seawater. High drag is achieved through a large circular cloth drogue attached to the float. The horizontal motion of the float is determined by acoustic tracking (RAFOS) and its vertical motion is determined from pressure. Data is relayed at the end of the 2-month mission via satellite (ARGOS). These data are supplemented by meteorological and oceanographic data from other investigators involved in the Labrador Sea Deep Convection experiment.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Lagrangian Floats For Deep Convection				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 NE 40th Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			


WORK COMPLETED

Our data from both 1997 (13 floats) and 1998 (7 floats) has been completely processed and calibrated and most of the basic data analysis completed. The first cut at a detailed comparison between the data and LES modelling of deep convection by Ramsey Harcourt (NPS, Monterey) has been completed. Much of this work was done by graduate student Elizabeth Steffen who will use this data for her Ph.D. work.

RESULTS

The two figures show comparison between the float data and the results of an LES model (Ramsey Harcourt, NPS) of deep convection in the Labrador Sea. Data from simulated Lagrangian floats, some perfect and some imperfect, is compared to the real data.

The first figure (left) compares the profile of vertical heat flux, $\langle w'T \rangle$. The major difference between model and data is a near surface maximum in measured heat flux which is not reproduced well by the model. The major reason for this is the low resolution of the model (50 m) which is insufficient to resolve this feature. Physically, this maximum may result from near-surface processes, such as horizontal Ekman heat flux, which are not included in the model, or from the non-Lagrangian features of the float motion.



The second figure (right) compares the rms vertical velocity from the model and data. The comparison is good in the middle of the boundary layer, particularly for the "buoyant float" simulation which most accurately represents the float dynamics. The model fails near the surface, probably again due to its low resolution.

Overall, the model does a good first order prediction of the measured properties, but clearly shows some differences with the observations. The significance of these will be investigated as the major part of E. Steffen's Ph.D. work.

IMPACT/APPLICATIONS

We expect this study to provide the first direct evaluation of models of deep convective turbulence and thus lead to improved simulations of this important process.

RELATED PROJECTS

These floats are close relatives of those used to study mixing in the Littoral zone and upper ocean mixed layer funded by ONR 322PO. Mixing processes in these various environments are similar in many ways and we learn the most by comparing and contrasting them.

PUBLICATIONS

The Lab Sea Group, 1998, The Labrador Sea Deep Convection Experiment, *Bulletin of the American Meteorological Society*, Vol. 79, No. 10, pp. 2033–2058.