

Measurements of the Aerosol Light-Scattering Coefficient at Ambient and 85% Relative Humidity on the ONR Pelican During ACE-2

Dean A. Hegg

University of Washington
Department of Atmospheric Sciences
Box 351640
Seattle, WA 98195-1640

phone: (206) 685-1984 fax: (206) 685-7160 email: deanhegg@atmos.washington.edu

David S. Covert

University of Washington
Department of Atmospheric Sciences
Box 354235,
Seattle, WA 98195-4235

phone: (206) 685-7461 fax: (206) 685-3397 email: dcovert@u.washington.edu

Award #: N00014-97-1-0132

http://cargsun2.atmos.washington.edu/~santiago/ACE2/html/2panel_page.html

LONG-TERM GOALS

The ultimate goal of this study is the determination of aerosol hygroscopicity and its impact on physical (e.g., size) and optical properties (e.g., scattering coefficient) of marine aerosols on large spatial scales. Related to this, a methodology for retrieving the dry aerosol properties from ambient measurements on large spatial scales, for example, by satellite remote sensing, is of great interest.

OBJECTIVES

The immediate objectives of the study involve data analysis from two separate field studies already completed. The first such study is the ACE-2 study completed in the summer of 1997 and the second the DECS study completed this past summer (1999). Objectives for each study are given below:

- Determine the impact of humidity on the aerosol scattering coefficient for different aerosol types (ACE-2).
- Determine the impact of humidity on upwelling radiance from aerosol backscatter (ACE-2)
- Test the constancy of the dry aerosol number-to-volume ratio, and the impact of humidity on it, in marine air (ACE-2 and DECS).
- Explore the feasibility of remotely retrieving aerosol hygroscopicity.
- Explore the feasibility of a remote retrieval of dry aerosol volume and CCN number concentration (ACE-2 and DECS).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Measurements of the Aerosol Light-Scattering Coefficient at Ambient and 85% Relative Humidity on the ONR Pelican During ACE-2				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Department of Atmospheric Sciences, Seattle, WA, 98195				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

- Explore the formation of new particles near clouds.

APPROACH

The approach utilized to measure the RH dependence of the aerosol light-scattering coefficient was described in the last annual report and will not be re-iterated here. The approach to appraising the constancy of the dry aerosol number to volume ratio is, in part, related to this. Once the functional dependence of the scattering on RH is determined, it can be applied to the measured ambient scattering (for which the measurement RH is known) to correct the values to low RH (£30%, i.e., "dry") for comparison with similarly corrected and simultaneous ambient measurements of the aerosol volume from the PCASP measurements.

To access the impact of humidity on upwelling radiance, and the closely related issue of remote retrieval of aerosol hygroscopicity and aerosol dry volume and CCN concentration, forward radiative transfer simulations (from the in-situ measurements of aerosol size and composition) are necessary. For the data gathered during ACE-2, the well-known 6S code has been utilized (Vermote et al., 1997). For the DECS data set, a version of the DISTORT radiative transfer code will be utilized (MODTRAN, Berk et al., 1998) together, of course, with the measurements themselves. For documenting particle formation near cloud, a suite of three CN counters, operating at different supersaturations, has been employed. The measurements were continuous (1 Hz) as the aircraft passed in and out of cloud, thus permitting assessment of small particle gradients near cloud.

WORK COMPLETED

In addition to the values obtained for the aerosol hygroscopicity parameter (g) reported in last year's report for the ACE-2 data set, several other tasks have been completed. An assessment has been made of the constancy of the aerosol number-to-volume ratio for the ACE-2 data set. A preliminary assessment of the feasibility of a retrieval of the dry value of this parameter has also been made. The impact of hygroscopicity on upwelling irradiance due to aerosol backscatter for the ACE-2 data set has been examined.

RESULTS

Analysis of the ACE-2 data set has yielded a mean, dry aerosol number-to-volume ratio of $168 \pm 21 \mu\text{m}^{-3}$, in agreement with several previous studies in different marine locals. However, the mean value is quite "noisy," with a large standard error. This is in part due to uncertainties generated by the need to correct the measurement to "dry" conditions. Nevertheless, analysis suggests that this noise will render the remote retrieval of marine CCN concentration difficult when based on retrieval of the dry aerosol volume alone. This can be seen in Figure 1, which shows a comparison of CCN predicted from the calculated dry aerosol volume to actual CCN concentrations. On the other hand, if concurrent retrieval of both the dry volume and number-to-volume ratios (from the retrieved effective radius) is feasible, the recovery of actual CCN concentrations proved appreciably more accurate (Figure 2). These results have been reported in Hegg and Jonsson (1999).

Observed vs CCN predicted from v alone




Figure 1.

Observed vs CCN predicted from v and n/v




Figure 2.

The radiative transfer calculations demonstrating the separable impact of aerosol hygroscopicity on upwelling radiance, were carried out for the radiometer wavelength bands (and viewing angles) of both the MISR and MODIS radiometers. The most promising results were found for the near infrared bands of the MODIS sensor. These results are shown in Figure 3. It can be seen that, at least for the higher optical depths, the impact of hygroscopicity (γ) on the radiance does indeed suggest that retrieval of hygroscopicity is possible if estimate of the ambient RH with $\sim 10\%$ can be made. This analysis has been reported in Gasso et al. (1999).




Figure 3.

Work has just commenced on reduction of data from the DECS study conducted this last summer. The analysis will center on analysis of particle formation and light scattering near clouds, and comparisons with MISR and MODIS retrievals.

IMPACT/APPLICATIONS

These results suggest that remote retrieval of both aerosol hygroscopicity and the CCN number concentration may be possible in marine air, where measurement by more conventional means is rare and expensive.

TRANSITIONS

As was the case with the ACE-2 data, the DECS hygroscopic data will be used by all participants in the study. More indirectly, the analysis of the potential for remote retrieval of aerosol hygroscopicity and of CCN concentrations should be of wide interest to the aerosol community.

RELATED PROJECTS

The impact of aerosol hygroscopicity on aerosol radiative properties, and the resultant impact on potential retrievals of dry aerosol parameters via remote sensing, is of great interest to both the MODIS and MISR science teams at NASA. We are cooperating with scientists on both teams.

REFERENCE

Berk, A., L. S. Bernstein, G. P. Anderson, P. K. Ackorya, D. C. Robertson, J. H. Chetwynd and S. M. Adler-Golden, MODTRAN cloud and multiple scattering upgrades with application to AVIRIS, *Remote Sensing of Environments*, 65, 367-375, 1998.

Vermote, E. J., D. Tanré, J. L. Deuzé, M. Herman and J. J. Mockette, Second simulation of the satellite signal in the solar spectrum, 6S: An overview, *IEEE Trans. Geos. and R. Sens.*, 35, 675-686, 1997.

PUBLICATIONS

Collins, D. R., H. H. Jonsson, J. H. Seinfeld, R. C. Flagan, S. Gasso, D. A. Hegg, P. B. Russell, B. Schmid, J. M. Livingston, E. Ostrom, K. J. Noone, L. M. Russell and J. P. Putaud, In situ aerosol size distributions and clear column radiative closure during ACE-2, *Tellus*, In press, 1999.

Durkee, P. A., K. E. Nielsen, P. J. Smith, P. B. Russell, B. Schmid, J. M. Livingston, B. N. Holben, C. Tomasi, V. Vitale, D. Collins, R. C. Flagan, J. H. Seinfeld, K. J. Noone, E. Ostrom, S. Gasso, D. Hegg, L. M. Russell, T. S. Bates and P. K. Quinn, Regional aerosol optical depth characteristics from satellite observations: ACE-1, TARFOX and ACE-2 results, *Tellus*, In press, 1999.

Gasso, S., D. A. Hegg, D. Collins, K. Noone, E. Ostrom, B. Schmid, P. B. Russell, J. M. Livingston, P. A. Durkee and H. Jonsson, Influence of humidity on the aerosol scattering coefficient and its effect on the upwelling radiance during ACE-2, *Tellus*, In press, 1999.

Hegg, D. A. and H. Jonsson, The aerosol number-to-volume relationship and relative humidity in the Eastern Atlantic, *J. Geophys. Res.*, In press, 1999.

Livingston, J. M., V. N. Kapustin, B. Schmid, P. B. Russell, P. K. Quinn, T. S. Bates, P. A. Durkee, P. J. Smith, V. Freudenthaler, S. Gasso, D. A. Hegg, D. R. Collins, R. C. Flagan, J. H. Seinfeld, V. Vitale and C. Tomasi, Shipboard sunphotometer measurements of aerosol optical depth spectra and columnar water vapor during ACE-2 and comparison with selected land, ship, aircraft and satellite measurements, *Tellus*, In press, 1999.

Schmid, B., J. M. Livingston, P. B. Russell, P. A. Durkee, H. Jonsson, D. R. Collins, R. C. Flagan, J. H. Seinfeld, S. Gasso, D. A. Hegg, E. Ostrom, K. J. Noone, E. J. Welton, K. J. Voss, H. R. Gordon, P.

Formenti and M. O. Andreae, Clear sky closure studies of lower tropospheric aerosol and water vapor during ACE-2 using airborne sunphotometer, airborne in-situ, space-borne, and ground-based measurements, *Tellus*, In press, 1999.