

Toward a Universal Sea Spray Source Function (UNISOURCE)

Gerrit de Leeuw
TNO Physics and Electronics Laboratory
P.O. Box 96864
2509 JG The Hague
The Netherlands
phone: +31 70 3740462 fax: +31 70 3740654 email: deleeuw@fel.tno.nl

Award Number: N00014-96-1-0581

<http://www.tno.nl>

LONG TERM GOALS

The long-term goals of the research are to understand and assess the effects of the atmosphere on the detection of targets at low altitudes over sea in coastal regions using IR and radar systems. Effects considered are transmission losses due to aerosols and water vapor, effects of turbulent fluctuations of the air temperature on blurring and scintillation, and effects of vertical temperature and water vapor gradients on IR and rf refractivity.

OBJECTIVES

The objectives of the research performed in the framework of the present grant are to further analyze data from the EOPACE and RED experiments, with new efforts focussing on the sea spray source function, i.e.:

- to determine the turbulence and refractivity in the inhomogeneous coastal boundary layer and their effects on imaging of low altitude point targets;
- to determine effects of scintillation and refraction in the MW IR band as function of environmental conditions through measurements with a camera mounted ashore at a range of levels above the sea surface, looking at a source mounted on FLIP;
- to improve the description of the aerosol size distribution as function of height and meteorological parameters;
- to determine the effect of sea spray aerosol on IR propagation as function of environmental conditions;
- to determine the generation of sea spray aerosol from breaking wind waves by the bubble-mediated mechanism and by direct tearing from the wave tops, the dispersal of freshly produced aerosol in the surface layer and the influence of wave induced turbulent phenomena, and the subsequent transport of the aerosol throughout the atmospheric boundary layer.

APPROACH

Data from the EOPACE IOP's at the California coast and in Duck, North Carolina in 1996-1999 and from the RED experiments at the coast of Hawaii in August/September, 2001, are further analyzed and interpreted. Research in FY03 was focused on the analysis of aerosol and bubble data from the RED experiments, and on refraction effects on near-surface IR propagation using results from EOPACE and other experiments. Contributions to EOPACE and RED experiments were described in previous annual

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Toward a Universal Sea Spray Source Function (UNISOURCE)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) TNO Physics and Electronics Laboratory,,P.O. Box 96864,2509 JG The Hague,,The Netherlands, ,				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

reports. Preparations are made to determine a formulation for the production of primary marine aerosol that applies in a wide variety of conditions. As part of this, a sea spray aerosol workshop is organized in the UK in the spring of 2004.

WORK COMPLETED

Work completed in years prior to FY03 was summarized in earlier reports. In FY03 the following work was completed:

- The publication of the results from the analysis of IOP4 has appeared in Applied Optics [Doss-Hammel et al., 2002]. A publication with results from other IOP's, focussing on refraction effects, has been submitted for publication [De Jong et al., 2003]. Results from laboratory experiments on primary marine aerosol production by bubble bursting have been published in J. Geophys Res. [Mårtensson et al., 2003]. An analysis of data from a Mediterranean site (Porquerolles, France) has been published in Opt. Eng. [Piazzola et al., 2003]. See the publications section for articles published and submitted in FY03 on related subjects.
- TNO-FEL contributed to a RED overview paper that has been submitted for publication in BAMS [Anderson et al., 2003].
- The data from the Rough Evaporation Duct (RED) experiments off Oahu (HI) in August/September 2001 have been partly analyzed. Initial results were presented at several scientific conferences (see listing in 'Publications').
- An overview of primary marine source functions is in press [Schulz et al., 2003].

RESULTS

During the Rough Evaporation Duct (RED) experiments off Oahu (HI) in August/September 2001, aerosol particle size distributions were measured with optical particle counters and a volatility system. The volatility system was also utilised in a combination with a sonic anemometer and a CPC to measure primary marine aerosol production by application of the eddy covariance technique. Particle size distribution profiles were measured with Rotorod impaction samplers [De Leeuw, 1986]. Bubble size distributions were measured with an optical bubble measuring system [De Leeuw and Cohen, 2001]. Lidar measurements were made to characterise the boundary layer structure (cf. Kunz et al. [2002] for a description of the lidar system and the type of measurements). All these measurements were made from FLIP. IR propagation measurements were made between FLIP and a number of sites ashore [De Jong, 2002].

The aerosol size distributions and volatility spectra have been analysed and used in the Coastal Aerosol Transport model CAT [Vignati et al., 2001] to derive the sea salt source function. The volatility spectra recorded at a temperature of 600°C after subtraction of those recorded at the higher temperature, were used to ascertain that only sea spray was measured. The size spectra were binned in 1 ms^{-1} wind speed intervals. Particles larger than $1 \text{ }\mu\text{m}$ in diameter, the concentrations of which were falling off due to inefficient collection by the volatility system, were deleted. Likewise, the concentrations of particles smaller than $2 \text{ }\mu\text{m}$ measured with the optical particle counter were high due to the occurrence of other constituents than sea salt. Although it cannot be ruled out that larger particles did not contain other constituents than sea salt, these particles were retained to construct sea salt size spectra in the size range from 0.2 to $>40 \text{ }\mu\text{m}$ for wind speeds from 5 to 10 ms^{-1} . The results are shown in Figure 1. It is noted that the relative humidity (RH) during the RED experiments were always close to 80%, and hence no humidity corrections were needed.




Figure 1. Wind speed binned size spectra derived from the volatility measurements (denoted by VWs nn, see legend) and the optical particle counters (denoted by PWs nn).

These size spectra were subsequently used to construct bimodal lognormal size distributions in the radius range 0.1-10 μm :

$$y = A \exp \left[C \ln \left(\frac{R_{80}}{B} \right)^2 \right] + D \exp \left[F \ln \left(\frac{R_{80}}{E} \right)^2 \right]$$

where R_{80} is the particle radius in micron at 80 % relative humidity and A, B, C, D, E, F are given in the following table for wind speed from 5 to 10 m/s.

Table 1. Coefficients from fitting Lognormal size distributions to the combined volatility and optical particle counter data, binned in 1 ms^{-1} wind speed intervals between 5 and 10 ms^{-1} .

WS	5	6	7	8	9	10
A	8	8.5	12.496	11.497	14.494	14.494
B	0.0643	0.0643	0.0643	0.0643	0.0643	0.0643
C	1.126	1.126	1.027	0.829	0.829	0.631
D	1	1	1.8	2.2	2.7	3.5
E	0.7996	0.7996	0.7996	0.7996	0.7996	0.7996
F	1.099	1.099	1.099	1.099	1.099	1.099

These values were used to derive the wind speed dependence of the parameters A, C and D :

$$\begin{aligned}
A(ws) &= 1.4129ws + 0.9831 & R^2(A) &= 0.87 \\
C(ws) &= -0.1018ws + 1.6917 & R^2(C) &= 0.93 \\
D(ws) &= 0.5143ws - 1.8238 & R^2(D) &= 0.96
\end{aligned}$$

where R^2 are the correlation coefficients:

The CAT model has been run using wind speed from 5 to 10 m/s. The column has been advected for 300 km over the open sea in neutral conditions. The Boundary Layer height has been set to 1000 m and the concentrations at 16 m height are reported. The initial size distribution was set to zero for both sea salt and continental aerosols. Sea salt is supposed to be generated by the sea surface. At the end of the trajectory the aerosol concentrations predicted by CAT have been compared to the concentrations measured by PMS and the volatility system, see Figure 2 for results for 5 and 10 ms^{-1} . The submicron particles were given most weight because they are only sea salt. For the larger particles they are high by roughly a factor of 2. This supports the earlier suggestion that these larger particles do not exclusively consist of sea spray, likely due to the occurrence of organics, or due to residues from other origin, long range transported and transformed due to a variety of processes.




Figure 2. Comparison of CAT-calculated sea spray size distributions with sea spray size distributions derived from the volatility measurements (blue) and the optical particle counters (red).

Thus exercise resulted in the following expression for the sea salt source function:

$$\frac{dF_N}{dr} = c \left(\sum_{i=1}^2 A_i e^{-C_i \ln\left(\frac{r}{r_i}\right)^2} \right), \quad \text{in } \mu\text{m}^{-1} \text{ m}^{-2} \text{ s}^{-1}$$

where:

$$A_1 = 1.41u + 0.98$$

$$A_2 = 0.51u - 1.82$$

$$C_1 = -0.1u + 1.69$$

$$C_2 = 1.09$$

$$R_1 = 0.0643$$

$$R_2 = 0.7996$$

$$c = (0.24u + 0.4) \cdot 10^4$$

In Figure 3, this source function is compared with the ones derived by Monahan et al. [1986] and Smith et al. [1993], that are commonly referred to in the literature, and the recently published source function by Vignati et al. [2001]. For the smaller particles the source functions are quite close, although at the lower end the Vignati et al. source function decreases toward smaller sizes, in good agreement with recent measurements. For particles with radius around 1 μm , the current source function is higher than all other ones, while the curvature suggests that for larger particles it may approach Vignati et al. The higher concentrations in the 1 μm range may be a water temperature effect: during RED it was about 28°C, whereas the other source functions are based on data collected in colder water of the North Atlantic. However, the preliminary analysis of direct covariance measurements of the primary marine aerosol flux, including data from RED and spanning a water temperature range from 0°C to 28°C, does not show a clear dependence. This analysis is continued, and new laboratory experiments are planned.




Figure 3. Comparison of source functions derived by Monahan et al. [1986] (blue), Smith et al. [1993] (green), Vignati et al. [2001] (red) and current work (black).

IMPACT/APPLICATIONS

The results can be used to assess the effects of the atmosphere on the performance of thermal imagers over sea, and in particular the performance of LR-IRST systems. Another important application is in the field of numerical weather forecasting because of the influence on the solar irradiation at the Earth surface, as well as the assessment of the impact of aerosols on climate. Sea spray aerosol has been estimated to contribute 44% to the total aerosol optical depth, but with an uncertainty of a factor of three depending on the source function estimate used (IPCC, 2001). The surf-produced aerosol affects atmospheric processes involving sea spray particles, such as heterogeneous reactions, at fetches of at least 10 km in off-shore winds. Reaction between sea spray and HNO_3 has consequences for atmospheric inputs of nitrogen compounds in coastal waters, and thus eutrophication [De Leeuw et al., 2001a;b, 2003a;b]. Over land, sea spray influences fragile coastal eco-systems, and the corrosive properties cause damage to buildings, structures and cultural heritage.

TRANSITIONS

The EOPACE and RED results of TNO-FEL are exchanged with other EOPACE and RED participants, for common analysis combining all required expertise to achieve the EOPACE and RED goals. Common EOPACE publications have been published, others are submitted or in preparation. Accurate sea spray source functions are important in regional and global scale transport models and work is in progress with regional and global scale modelers to promote the use of our results.

RELATED PROJECTS

The efforts described above are in conjunction with other projects addressing electro-optical propagation over sea, in part basic research, in part applied research. They take place in conjunction with studies funded by the Netherlands Ministry of Defense, including work on long-range transmission, IRST and backgrounds. Data from other areas, e.g. the North Sea, the North Atlantic, the Mediterranean and the Baltic, are from other projects supported by the Netherlands Ministry of Defense, the EU or other funding agencies.

REFERENCES

- Anderson, K., P. Caffrey, A. Clarke, K. Crahan, K. Davidson, A. De Jong, G. de Leeuw, D. Dion, S. Doss-Hammel, P. Frederickson, C. Friehe, D. Hegg, T. Hristov, D. Khelif, J. Reid, S. Reising, E. Terrill, and D. Tsinikidis (2003). The Rough Evaporation Duct (RED) Experiment; An Assessment of Boundary Layer Effects in a Trade Winds Regime on Microwave and Infrared Propagation over the Sea. [Submitted].
- De Jong, A.N., TNO Physics and Electronics Laboratory, Report, 2002.
- De Jong, A.N., G. de Leeuw, H. Winkel, P.J. Fritz and M.M. Moerman (2003). Atmospheric refraction effects on optical/IR sensor performance in a littoral maritime environment. [Submitted].

- De Leeuw, G. Vertical profiles of giant particles close above the sea surface. *Tellus* 38B, 51-61, 1986.
- De Leeuw, G., and L.H. Cohen. Bubble size distributions on the North Atlantic and the North Sea. in *Gas Transfer and water Surfaces*, edited by M.A. Donelan, W.M. Drennan, E.S. Salzman, and R. Wanninkhof, pp. 271-277, AGU, 2001.
- De Leeuw, G., G.J. Kunz, G. Caulliez, D.K. Woolf, P. Bowyer, I. Leifer, P. Nightingale, M. Liddicoat, T.S. Rhee, M.O. Andreae, S.E. Larsen, F.A. Hansen and S. Lund, LUMINY - An overview. in *Gas Transfer and water Surfaces*, edited by M.A. Donelan, W.M. Drennan, E.S. Salzman, and R. Wanninkhof, pp. 291-294, AGU, 2001.
- De Leeuw, G., L.H. Cohen, L.M. Frohn, G. Geernaert, O. Hertel, B. Jensen, T. Jickells, L. Klein, G. J. Kunz, S. Lund, M.M. Moerman, F. Müller, B. Pedersen, K. von Salzen, K. H. Schlünzen, M. Schulz, C. A. Skjøth, L.L. Sorensen, L. Spokes, S. Tamm and E. Vignati. Atmospheric input of nitrogen into the North Sea: ANICE project overview. *Continental Shelf Research*, (21) 18-19 (2001),pp. 2073-2094, 2001.
- De Leeuw, G., C.Ambelas Skjøth, O. Hertel, T. Jickells, L. Spokes, E. Vignati, L. Frohn, J. Frydendall, M. Schulz, S. Tamm, L.L. Sørensen and G.J. Kunz, Deposition of Nitrogen into the North Sea, *Atm. Env.*, 37, 145-165 (2003).
- De Leeuw, G., L. Spokes, T. Jickells, C. Ambelas Skjøth, O. Hertel, E. Vignati, S. Tamm, M. Schulz, L.-L. Sorensen, B. Pedersen, L. Klein, and K. H. Schlünzen (2003). Atmospheric Nitrogen Inputs into the North Sea: Effect on Productivity. *Continental Shelf Research*, in press.
- Doss-Hammel, S.M., C.R. Zeisse, A.E. Barrios, G. de Leeuw, M.Moerman, A.N. de Jong, P. A. Frederickson, and K. L. Davidson (2002), Low altitude infrared propagation in a coastal zone: refraction and scattering. *Appl. Opt.* 41(18), 3706-3724, 2002.
- Kunz, G.J., G. de Leeuw, C. O'Dowd and E. Becker, LIDAR studies of atmospheric boundary layer structure and sea spray aerosol plumes generation and transport at Mace Head, Ireland (PARFORCE experiment). *J. Geophys. Res., special issue on PARFORCE*, doi:10.1029/2001JD001240, 2002.
- Mårtensson, M., E. D. Nilsson, G. de Leeuw, L.H. Cohen, and H-C Hansson, Laboratory simulations of the primary marine aerosol generated by bubble bursting, *JGR-Atmospheres* 108 (D9), 10.1029/2002JD002263 (2003).
- Monahan, E.C., C.W. Fairall, K.L. Davidson and P.J. Boyle, Observed inter-relations between 10 m winds, ocean whitecaps and marine aerosols, *Quart. J. R. Met. Soc.* 109, 379-392, 1983.
- Piazzola, J., F. Bouchara, A.M.J. van Eijk and G. de Leeuw (2003) Development of the Mediterranean extinction code (MEDEX), *Opt. Eng.*, 42, 912-924 (2003).
- Schulz, M., G. de Leeuw, and Y. Balkanski, Sea-salt aerosol source functions and emissions, in "Emissions of Atmospheric Trace Compounds", eds. C. Granier, P. Artaxo and C. Reeves, Kluwer, 2003. (in press).

- Smith, M.H., P.M. Park, and I.E. Consterdine, Marine aerosol concentrations and estimated fluxes over the sea, *Q. J. R. Meteorol. Soc.*, 119, 809-824, 1993.
- Vignati, E., G. de Leeuw and R. Berkowicz. Modeling coastal aerosol transport and effects of surf-produced aerosols on processes in the marine atmospheric boundary layer, *JGR-Atmospheres*, 106, 20225-20238, 2001.

PUBLICATIONS

Refereed Journals:

- R.1 Doss-Hammel, S.M., C.R. Zeisse, A.E. Barrios, G. de Leeuw, M. Moerman, A.N. de Jong, P. A. Frederickson, and K. L. Davidson (2002), Low altitude infrared propagation in a coastal zone: refraction and scattering. *Appl. Opt.* 41(18), 3706-3724, 2002 [published].
- R.2 O'Dowd, C.D., K. Hämeri, J. Mäkelä, M. Väkeva, P. Aalto, G. de Leeuw, G. Kunz, E. Becker, H.-C. Hansson, E. Becker, A.G. Allen, R.M. Harrison, C. Kleefeld, M. Geever, S.G. Jennings and M. Kulmala, Coastal new particle formation: Environmental conditions and aerosol physico-chemical characteristics during nucleation bursts, *J. Geophys. Res.*, 10.1029/2000JD000206, 2002 [published].
- R.3 O'Dowd, C.D., K. Hämeri, J. Mäkelä, L. Pirjola, M. Kulmala, S.G. Jennings, H. Berresheim, H.-C. Hansson, G. de Leeuw, G.J. Kunz, A.G. Allen, C.N. Hewitt, A. Stroh, Y. Viisanen and T. Hoffmann, A dedicated study of new particle formation and fate in the coastal environment (PARFORCE): Overview of objectives and initial achievements, *J. Geophys. Res.*, 10.1029/2001JD000555, 2002 [published].
- R.4 Kleefeld, C., C.D. O'Dowd, S. O'Reilly, S. G. Jennings, P. Aalto, E. Becker, G. Kunz and G. de Leeuw, The relative contribution of sub and super micron particles to aerosol light scattering in the marine boundary layer (MBL). *J. Geophys. Res.*, 10.1029/2000JD000262, 2002 [published].
- R.5 Kunz, G.J., G. de Leeuw, C. O'Dowd and E. Becker, LIDAR studies of atmospheric boundary layer structure and sea spray aerosol plumes generation and transport at Mace Head, Ireland (PARFORCE experiment). *J. Geophys. Res.*, 10.1029/2001JD001240, 2002 [published].
- R.6 De Leeuw, G., G.J. Kunz, G. Buzorius and C. O'Dowd (2001). Meteorological influences on coastal new particle formation. *J. Geophys. Res.*, 10.1029/2001JD001478, 2002 [published].
- R.7 Hertel, O., C. Ambelas Skjøth, L.M. Frohn, E. Vignati, J. Frydendall, G. de Leeuw, Uwe Schwarz and Stefan Reis. Assessment of the Atmospheric Nitrogen Inputs into the North Sea using a Lagrangian model. *Phys. Chem. Earth* 27, 1507-1515, 2002 [published].
- R.8 Schmid B., J. Redemann, P. B. Russell, P. V. Hobbs, D. L. Hlavka, M. J. McGill, B. N. Holben, E. J. Welton, J. R. Campbell, O. Torres, R. A. Kahn, D. J. Diner, M. C. Helmlinger, D. A. Chu, C. Robles Gonzalez, and G. de Leeuw, Coordinated airborne, spaceborne, and ground-based measurements of massive, thick aerosol layers during the dry season in southern Africa, *J. Geophys. Res.*, 108(D13), 8496, doi: 10.1029/2002JD002297, 2003 [published].

- R.9 Robles-Gonzalez, C., M. Schaap, G. de Leeuw, P.J.H. Builtjes and M. van Loon, Spatial variation of aerosol properties derived from satellite observations and comparison with model calculations, *Atmospheric Chemistry and Physics*, 3, 521-533, 2003 [published].
- R.10 Mårtensson, M., E. D. Nilsson, G. de Leeuw, L.H. Cohen, and H-C Hansson, Laboratory simulations of the primary marine aerosol generated by bubble bursting, *JGR-Atmospheres* 108 (D9), 10.1029/2002JD002263 (2003) [published].
- R.11 Piazzola, J., F. Bouchara, A.M.J. van Eijk and G. de Leeuw (2003) Development of the Mediterranean extinction code (MEDEX), *Opt. Eng.*, 42, 912-924 (2003) [published].
- R.12 De Leeuw, G., C.Ambelas Skjøth, O. Hertel, T. Jickells, L. Spokes, E. Vignati, L. Frohn, J. Frydendall, M. Schulz, S. Tamm, L.L. Sørensen and G.J. Kunz, Deposition of Nitrogen into the North Sea, *Atm. Env.*, 37, 145-165 (2003) [published].
- R.13 Leifer, I., G. De Leeuw, and L.H. Cohen. Optical measurement of bubbles: System' design and application. *J. Atm. Ocean. Tech.* 20(9), 1317-1332, 2003 [published].
- R.14 Leifer, I., G. de Leeuw and L.H. Cohen. Calibrating optical bubble size by the displaced mass method. *Chem. Eng. Sci.*, in press (2003) [published].
- R.15 De Leeuw, G., L. Spokes, T. Jickells, C. Ambelas Skjøth, O. Hertel, E. Vignati, S. Tamm, M. Schulz, L.-L. Sorensen, B. Pedersen, L. Klein, and K. H. Schlünzen (2003). Atmospheric Nitrogen Inputs into the North Sea: Effect on Productivity. *Continental Shelf Research* [in press].
- R.16 Schulz, M., G. de Leeuw, and Y. Balkanski, Sea-salt aerosol source functions and emissions, in "Emissions of Atmospheric Trace Compounds", eds. C. Granier, P. Artaxo and C. Reeves, Kluwer, 2003. [in press].
- R.17 Anderson, K., P. Caffrey, A. Clarke, K. Crahan, K. Davidson, A. De Jong, G. de Leeuw, D. Dion, S. Doss-Hammel, P. Frederickson, C. Friehe, D. Hegg, T. Hristov, D. Khelif, J. Reid, S.Reising, E. Terrill, and D. Tsinikidis (2003). The Rough Evaporation Duct (RED) Experiment; An Assessment of Boundary Layer Effects in a Trade Winds Regime on Microwave and Infrared Propagation over the Sea. [Submitted].
- R.18 De Jong, A.N., G. de Leeuw, H. Winkel, P.J. Fritz and M.M. Moerman (2003). Atmospheric refraction effects on optical/IR sensor performance in a littoral maritime environment [Submitted].

Proceedings:

- P.1 Van Eijk, A.M.J., L.H. Cohen, L.J. Navarro and G. de Leeuw (2002). Near-surface aerosol transmission. SPIE conference 'Optics in Atmospheric Propagation and Adaptive Systems V, Crete, 23-27 September 2002 [published].
- P.2 De Leeuw, G., C. Robles Gonzalez, J. Kusmierczyk-Michulec, R. Decae and P.J. Veeffkind (2002). Retrieval of aerosol optical depth from satellite measurements using single and dual view algorithms. In Proceedings of SPIE Vol. 4882 Remote Sensing of Clouds and the Atmosphere VII, edited by K.P.

Schäfer, O. Lado-Bordowsky, A. Camerón and R.H. Picard, (SPIE< Bellingham, WA, 2003, pp. 275-283 [published].

P.3 M. van Loon, P. Builtjes and G. de Leeuw (2002). PM_{2.5} concentrations over Europe: Combining satellite data and modelling. In: Borrego and Schayes (Eds.) Air pollution modeling and its application XV, Kluwer, New York, pp. 233-224 [published].

P.4 Nilsson, E.D., M. Mårtensson, S. van Ekeren, G. de Leeuw, M. Moerman, C. O'Dowd, R. Flanagan, and M. Geever (2003). Eddy correlation measurements of the primary marine aerosol source, AMS conference, Long Beach, CA [published].

P.5 De Leeuw, G., M. Moerman, L. Cohen, B. Brooks, M. Smith and E. Vignati (2003) Aerosols, bubbles and sea spray production studies during the RED experiments, Proceedings AMS conference, Long Beach, CA, 9-13 Feb., 2003 [published].

P.6 G. de Leeuw, E. Vignati, M. Moerman, B. Brooks and M.H. Smith (2003). Determination of the sea salt source function over the Pacific Ocean, J. Aerosol Sci., Abstracts of the European Aerosol Conference 2003, Madrid (31 Aug-5 Sept. 2003), pp. 537-538, 2003 [published].

P.7 E. D. Nilsson, M. Mårtensson, S. van Ekeren, G. de Leeuw, M. Moerman, C. O'Dowd, (2003). The primary marine aerosol source measured by eddy correlation for particles of different size and volatility. Aerosol Sci., Abstracts of the European Aerosol Conference 2003, Madrid (31 Aug-5 Sept. 2003), pp. 539-540, 2003 [published].

P.8 J. Kusmierczyk-Michulec, G. de Leeuw, C. Robles Gonzalez and R. Decae (2003). Retrieval of aerosol properties over land and over water with SCIAMACHY, J. Aerosol Sci., Abstracts of the European Aerosol Conference 2003, Madrid (31 Aug-5 Sept. 2003), pp. 887-888, 2003 [published].

P.9 C. Robles Gonzalez, G. De Leeuw, R. Decae and J. Kusmierczyk-Michulec (2003). Aerosol optical depth retrieved over land and water from the INDOEX campaign area from ATSR-2 data, J. Aerosol Sci., Abstracts of the European Aerosol Conference 2003, Madrid (31 Aug-5 Sept. 2003), pp. 893-894, 2003 [published].

P.10 R. Decae, G. Bazalgette Courrèges-Lacoste and G. de Leeuw (2003). The Dedicated Aerosol Retrieval Experiment (DARE). Scientific requirements for a dedicated satellite instrument to measure atmospheric aerosols. SPIE, Barcelona, Sept. 2003 [in press].

P.11 K. Smorenburg, G. Bazalgette Courreges-Lacoste, R. Decae, A. Court, G. de Leeuw and H. Visser (2003). DARE: Dedicated aerosol retrieval instrument. SPIE, Barcelona, Sept. 2003 [in press].