

# Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>30 SEP 1999</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1999 to 00-00-1999</b>	
4. TITLE AND SUBTITLE <b>The Chemistry and Bioaccumulation of Copper and Other Heavy Metals By Phytoplankton in the Water Column of San Diego Harbor and Its Relationship to Ecological Assessment and Water Quality</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Woods Hole Oceanographic Institution, Department of Marine Chemistry and Geochemistry, Woods Hole, MA, 02543</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>2</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# **The Chemistry and Bioaccumulation of Copper and Other Heavy Metals By Phytoplankton in the Water Column of San Diego Harbor and Its Relationship to Ecological Assessment and Water Quality**

James W. Moffett  
Department of Marine Chemistry and Geochemistry  
Woods Hole Oceanographic Institution  
Woods Hole MA 02543  
phone: (508) 289-3218 fax: (508) 457-2164 email: [jmoffett@whoi.edu](mailto:jmoffett@whoi.edu)  
Award #: N000149710718  
<http://www.whoi.edu>

## **LONG-TERM GOALS**

My long term objective for AASERT Awards has been to attract excellent graduate students into areas of environmental quality that are relevant to the Navy's interest. This AASERT was used to support a student interested in contaminant fate and transport in systems highly relevant to the Harbor Processes Program.

## **OBJECTIVES**

The objectives of this project were ultimately determined by the student's dissertation project. The student, Elizabeth Kujawinski, has been studying the role of protozoan grazing on the fate and transport of organic contaminants, specifically PCBs. Protozoans are important components of microbial assemblages in many carbon-rich contaminated waters. Ingestion of bacterial sized particles by these organisms can have a major impact on particle size distribution and chemistry. She is interested in the role of the grazing process in bioaccumulation and as a source of colloidal organic that is important in contaminant mobility.

## **APPROACH**

The approach is to study the partitioning of a suite of PCB congeners between bacterial, protozoan and colloidal particles in cultures of actively grazing marine protozoans and bacteria. The processes are characterized by filtration, headspace partitioning, and radiotracer measurements, and ancillary measurements of organic carbon concentration and composition, and surfactant concentrations are also being made.

## **WORK COMPLETED**

Cultures of 3 representative protozoans have been studied using the bacterium *Halomonas halodurans* as prey. A suite of PCB congeners were added to these systems and the kinetics of partitioning between bacterial and protist size classes was measured to distinguish between grazing and equilibrium partitioning as mechanisms for PCB accumulation in the protozoans.

Protozoan grazers can “clear” a dense bacterial culture in several days. The resulting media has high surfactant concentrations. In order to evaluate grazers as a source of surfactants, measurements of surfactant concentration, normalized to a Triton X-100 standard, were made using a standard polarographic technique.

Filtrates from “cleared” cultures were equilibrated with C-14 labelled PCBs in specially designed vessels to measure changes in headspace partitioning resulting from colloidal binding of PCBs. The concentration of colloidal organic C in these experiments was measured, and the lipid composition (which might contribute to PCB partitioning) was measured as well.

## **RESULTS**

Partitioning of PCBs amongst these size classes was shown to be dominated by physical-chemical processes rather than grazing, even for the most hydrophobic PCB congeners. This is a consequence of the high surface area to volume ratios of these unicellular organisms, and large cross membrane fluxes arising from preconcentration within phospholipid bilayers.

Ingestion of bacteria was associated with large production of surfactant material. Less than 2% of this material was lipid in composition. There were significant interspecies differences. Interestingly, the protists that produced the highest surfactant levels were the least effective at dissolving refractory iron oxide phases (from the thesis work of my former student K. Barbeau – also an AASERT recipient). Results suggests that protozoan grazing may be a major source of surfactant material in seawater.

The surfactant material was correlated with PCB partitioning in culture filtrates, suggesting that common materials are involved. Partition coefficients ( $K_{oc}$ ) for these samples were 7-10 fold higher than previous measurements in seawater. This suggests that materials derived directly from grazing may be more important in PCB mobility than more refractory materials in the water column. Thus areas of localized, high grazing activity might show enhanced PCB mobility.

## **IMPACT/APPLICATIONS**

Protozoan grazing is especially important in C-rich, nutrient poor regimes, including systems with high levels of hydrocarbon contamination. Many systems established for bioremediation have these characteristics. The contribution of protozoans to the overall chemistry of such regimes and their influence on contaminant transport will need to be considered.

## **PUBLICATIONS**

Kujawinski, E.B. , J.W. Moffett and J.W. Farrington (1999). The importance of passive diffusion in the uptake of PCBs by phagotrophic protozoa. Submitted to Applied and Environmental Microbiology.