

Tropical Cyclone Structure And Motion

Elizabeth A. Ritchie
Department of Meteorology, Code MR/Ri
Naval Postgraduate School
589 Dyer Rd., Room 254
Monterey, CA 93943-5114
Telephone: (831) 656-3430, FAX: (831) 656-3061, E-mail: ritchie@nps.navy.mil

Additional Principle Investigators: R. L. Elsberry, and P. A. Harr

N0001499WR30003

LONG-TERM GOALS

To improve tropical cyclone track and intensity prediction through a research program combining high resolution modeling and detailed observations to investigate physical processes by which the motion and structure of a tropical cyclone is modified.

OBJECTIVES

The objective of this project is to investigate the physical processes that occur as a tropical cyclone interacts with the environment such that motion and structure changes occur. Specific interactions being studied are with mesoscale convective systems (MCS) that have been hypothesized to impact the development, motion, and structure of tropical cyclones, with baroclinic environments in the tropics and in the midlatitudes during extratropical transition, and with topography as a tropical storm makes landfall. As a storm makes landfall, significant asymmetries in the low-level wind structure are expected to develop with marked impact on precipitation and wind damage patterns. During extratropical transition, radical changes to the storm structure occur as the warm core is eroded by intruding cold, dry air from the midlatitudes. Re-intensification to a strong midlatitude system is possible, and to further complicate matters, passage over islands with significant topographic features can occur during this transitioning period. In cases where forecast models did poorly in predicting the motion and re-intensification of the storm during these transitional periods, better understanding of these processes should improve motion and intensity forecasts.

APPROACH

Due to the scarcity of detailed observations in regions where tropical cyclones develop and move, high-resolution, idealized modeling is combined with observations in all studies described here. The degree of physical complexity included in current mesoscale models allows detailed examination of environmental and mesoscale convective system (MCS) impacts on the motion, structure, and intensity of tropical cyclones. However, caution must be taken when applying cause and effect arguments to describe the complex physical interactions that develop in these high-resolution models that may be a product of the model parameterizations rather than realistic physical processes. Thus, a tiered approach is employed, in which understanding of basic processes comes first and is built upon by gradually adding to the complexity of the modeling system, isolating each physical process in turn. The U.S. Navy's coupled ocean-atmosphere mesoscale prediction system is the primary model used in ongoing

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Tropical Cyclone Structure And Motion				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Meteorology, Code MR/Ri, 589 Dyer Rd., Room 254, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

studies into landfall and extratropical transition effects on tropical cyclone motion and structure. Where available, detailed observations such as those available from the ONR-sponsored TCM-92 and TCM-93 field experiments are used to verify processes examined in the model experiments.

WORK COMPLETED

Simulations using the PSU/NCAR mesoscale model investigated the possible role of mesoscale convective systems in producing tropical cyclone track deflections. New diagnostics were developed to aid in this effort. A journal article (Ritchie and Elsberry 1999a) from this study is currently in press in the *Monthly Weather Review*.

Simulations using the U.S. Navy's coupled ocean-atmosphere mesoscale prediction system (COAMPS) investigated the influence of topography, and midlatitude baroclinic environment on the structure of TY Peter. First-guess fields and boundary conditions were provided from the Navy Operational Global Prediction System at 1-degree latitude/longitude resolution and interpolated to the high-resolution domains used in COAMPS. Additional observations including high-resolution cloud-drift and water vapor winds were assimilated into the analysis to test the impact of better resolving mesoscale features in the initial fields. Sensitivity studies were conducted by varying the degree of idealization of topography in the model.

Simulations using COAMPS investigated the structural changes that occur as a simulated tropical cyclone interacts with an idealized midlatitude baroclinic zone. Idealized initial conditions for three environments were created based on composite wind fields at each of the three steps of transformation of a tropical cyclone identified in Klein et al. (1999). The environmental mass fields were then calculated using geostrophic and hydrostatic balance. The three-dimensional tropical cyclone wind field was spun-up from analytic wind fields with the initial mass fields calculated using gradient wind and hydrostatic balance. Boundary conditions were calculated directly from the idealized fields and held constant at the initial values through the simulation. The structural changes of the simulated tropical cyclone as it interacted with each of these three environments were validated with the observed cloud structure changes. Sensitivity studies were run in which the strength of the environmental baroclinic zone was changed. Finally, a single baroclinic environment was developed that included the aspects of all three steps of transformation. As the tropical cyclone advected through the three environments, the physical changes that occur because of the interaction between the tropical cyclone circulation and the environment were studied.

RESULTS

New interpretations have been obtained with respect to the role that a large MCS may have played in the track deflection of TY Robyn (1993) in the western North Pacific during the ONR-sponsored TCM-93 field experiment (Ritchie and Elsberry 1999a). Interaction between a mesoscale convective vortex associated with the MCS and a simulated tropical cyclone with a structure similar to that observed in TY Robyn was modeled. The simulations show that as much as a 2 m s^{-1} deceleration in Robyn's westward motion might have resulted due to an interaction with the mesoscale convective vortex, and then a more northward track deflection, which is similar to that observed. This study demonstrates that better analysis of mesoscale convective vortices may be important in numerical weather prediction models to obtain a correct steering current over the tropical cyclone as the majority of the steering in these simulations was located between 300 and 750 mb, a region of the atmosphere that is not well sampled by observing systems over the oceans.

The physical processes of transitioning and landfalling tropical cyclones have been investigated using COAMPS. As a tropical cyclone makes landfall, interaction with steep topography is hypothesized to produce localized extremes in low-level winds, which in turn may produce extreme precipitation events. A case study of TY Peter (1997) was used to simulate the processes that occur as a tropical cyclone makes landfall (Ritchie and Elsberry 1998; 1999d). The case was further complicated as the tropical cyclone was also beginning to transition to an extratropical storm while interacting with an upper-level trough. The tropical cyclone track guidance from both the Navy's global prediction system and COAMPS was slow. After assimilating additional satellite wind, and other observations into the analyses, the high-resolution COAMPS track speed guidance improved. In addition, the simulated tropical cyclone path was very close to that observed, and rainfall patterns closely matched SSM/I data (Ritchie and Elsberry 1999d). Interestingly, the majority of the rainfall (spatial distribution and intensity) occurred over the open ocean north of Japan rather than over the mountain regions of the islands. Idealized studies that simulated the same case over open ocean, and then over flat terrain, demonstrated that the majority of the rainfall occurs as a result of the baroclinic processes associated with interaction with the upper-level trough as the tropical cyclone transitions to an extratropical system rather than because of interaction with topography. However, very heavy rainfall occurred locally in association with steep orography.

Extratropical transition of tropical cyclones has been hypothesized to occur in two stages (Klein et al. 1999). The first stage, called transformation, occurs as a tropical cyclone encounters cooler air and waters and a baroclinic zone associated with midlatitude westerlies. Re-intensification subsequently occurs in some systems when, as part of an interaction with an upper-level trough, re-deepening of the system to a significant midlatitude storm takes place. An idealized study using COAMPS investigated processes associated with a mature tropical cyclone that transitions into an extratropical storm by simulating the effects of strong shear such as that found in the midlatitudes on the tropical cyclone structure. As the environmental vertical shear increases, the ability of the simulated tropical cyclone to remain coherent against the shear is reduced and the upper levels of the tropical cyclone begin to tilt. This tilting disrupts the convective processes that are important for maintaining vertical coherency of the vortex and an asymmetric pattern of convection develops to the northeast of the vortex core. At a critical vertical shear, the convection weakens in the upper-level portion of the tropical cyclone, which is then advected downstream. However, the lower portion of the tropical cyclone remains vertically coherent and upright. Whereas the central pressure of the tropical cyclone begins to rise similar to that observed, a sea-level trough in the pressure develops below the upper-level warm core that has been advected downstream. Thus, elongation of the sea-level pressure contours in the direction of the vertical shear also resembles observations. Sensitivity tests varying the environmental vertical wind shear are in progress to determine what tropical characteristics (intensity, size, etc.) lead to (or prevent) this vertical decoupling during extratropical transition.

The Klein et al. (1999) study identified three steps in the transformation stage of extratropical transition during which the tropical cyclone interacts with different aspects of the midlatitude, baroclinic environment. The composite wind fields of the Steps 1, 2, and 3 of tropical cyclone transformation that they calculated were used as a guide to create an idealized baroclinic environment. The tropical cyclone was then located within that environment at the location where the wind conditions matched those of the composite fields for each of the three steps of transformation. As the tropical cyclone interacted with the environment, the important structural changes were studied and verified against the observations in Klein et al. (1999) for each step. The main results were that the cloud fields developed in a manner very similar to that observed in the Klein et al. (1999) study. Physical processes that were identified in Klein et al. (1999) to be occurring during transformation

based on 1° lat./long. NOGAPS analyses were generally verified in the higher-resolution, but idealized COAMPS simulations. In addition, the higher-resolution COAMPS fields allowed investigation of a hypothesis proposed in Klein et al. (1999) that it was interaction with the low-level temperature gradient associated with the baroclinic zone that produced the dry slots in the clouds, and finally the erosion of the southern eyewall convection of the tropical cyclone. The COAMPS simulations suggest that it is the interaction between the tropical cyclone circulation and upper-level environmental winds that produce deep mechanically forced subsidence into the western and southern portions of the tropical cyclone core. This subsidence not only suppresses convection in that region of the tropical cyclone, but also produces subsidence warming in the middle to low levels of the tropical cyclone core. By Step 3, when the tropical cyclone is being advected more rapidly by the mean flow, this low-level warm anomaly tends to lag the remnants of the tropical cyclone warm core.

RELATED PROJECTS

1 - The simplest baroclinic environment within which a tropical cyclone can interact is that of vertical wind shear with no horizontal variation, and thus a constant, weak temperature gradient. The motion and structural development of tropical cyclones in such an environment has been numerically simulated using the PSU/NCAR mesoscale model in a related project with Prof. W. M. Frank (PSU) (Frank and Ritchie 1999a). Important results include the identification of persistent patterns of asymmetric convection and rainfall in the left quadrant of the storm. The rotational motion of the storm due to storm tilt that was identified in dry simulations is reduced in simulations that include parameterizations of convective processes, since the storm tilt due to environmental shear is almost completely eliminated in the presence of strong vertical momentum transport. In a weak environmental vertical wind shear (3 m s^{-1} over the entire troposphere), the storm is advected at 1 m s^{-1} with only a slight motion to the right of the zero-shear storm track. Currently, the model resolution is being increased to 5 km with a third inner mesh and all convective processes that were previously parameterized on the 15- and 45-km meshes are calculated using an explicit physical representation of convection scheme. Because the higher resolution model with explicitly resolved convection results in a more intense simulated tropical cyclone, it is possible to explore the parameter range of more realistic vertical wind shear values. Observational studies (e.g., DeMaria and Kaplan 1994) have shown that tropical cyclones tend to exhibit weakening when the environmental vertical wind shear exceeds about 10 m s^{-1} . Thus, three simulations with vertical wind shear values of 5, 10, and 15 m s^{-1} have been designed to explore the physical interactions between the tropical cyclone and the environment that result in eventual weakening of the tropical cyclone (Frank and Ritchie 1999b).

REFERENCES

- DeMaria, M., and J. Kaplan, 1994: A statistical hurricane intensity prediction scheme (SHIPS) for the Atlantic basin. *Wea. and Forec.*, **9**, 209-220.
- Frank, W.M., and E.A. Ritchie, 1999a: Effects of environmental flow upon tropical cyclone structure. *Mon. Wea. Rev.*, **127**, 2044-2061.
- Frank, W.M., and E.A. Ritchie, 1999d: Effects of vertical shear upon tropical cyclone intensity and structure. *Mon. Wea. Rev.* (*In preparation*)
- Klein, P. M., P. A. Harr, and R. L. Elsberry 1999: Extratropical transition of western North Pacific tropical cyclones: An overview and conceptual model of the transformation stage. *Mon. Wea. Rev.*,

(submitted)

Ritchie, E. A., and R. L. Elsberry, 1998: A study of the development of mesoscale asymmetries in tropical cyclone structure during landfall. *Preprints, 12th Conf. on Numerical Weather Prediction, Phoenix, Arizona, Amer. Meteor. Soc., Boston, MA 02108, 345-348.*

Ritchie, E.A., and R. L. Elsberry, 1999a: Simulated impacts of a mesoscale convective system on the track of Typhoon Robyn during TCM-93. *Mon. Wea. Rev. (In Press)*

Ritchie, E. A., and R. L. Elsberry, 1999d: A modeling study of the development of asymmetries in tropical cyclone structure during landfall. *Proceedings of the U.S. – Japan NR Task Committee “E” on Design for Wind Hazard Mitigation, 24-28 May, Tsukuba, Japan.*

PUBLICATIONS

Frank, W.M., and E.A. Ritchie, 1999a: Effects of environmental flow upon tropical cyclone structure. *Mon. Wea. Rev.*, **127**, 2044-2061.

Frank, W. M., and E. A. Ritchie, 1999b: Environmental effects on hurricane core structure. *Proceedings of the 23rd Conference on Hurricanes and Tropical Meteorology, American Meteorological Society, Boston, MA 02108, 680-681.*

Frank, W. M., and E. A. Ritchie, 1999c: Mechanisms for rapid intensity changes in hurricanes. *Proceedings of the 1st USWRP Science Symposium, March 29-31, Boulder, Colorado.*

Ritchie, E.A., and R. L. Elsberry, 1999a: Simulated impacts of a mesoscale convective system on the track of Typhoon Robyn during TCM-93. *Mon. Wea. Rev. (In Press)*

Ritchie, E. A., and R. L. Elsberry, 1999b: Mechanisms contributing to extratropical transition of tropical cyclones in the western North Pacific: an idealized study. *Proceedings of the 23rd Conference on Hurricanes and Tropical Meteorology, American Meteorological Society, Boston, MA 02108, 811-814.*

Ritchie, E. A., and R. L. Elsberry, 1999c: Mechanisms contributing to extratropical transition of tropical cyclones in the western North Pacific: A modeling study. *Proceedings of the International Workshop on the Extratropical Transition of Tropical Cyclones, 10-14 May 1999, Kaufbeuren, Germany. WMO – Technical document ###, World Meteorological Organization, Geneva (in press).*

Ritchie, E. A., and R. L. Elsberry, 1999d: A modeling study of the development of asymmetries in tropical cyclone structure during landfall. *Proceedings of the U.S. – Japan NR Task Committee “E” on Design for Wind Hazard Mitigation, 24-28 May, Tsukuba, Japan.*

Ritchie, E. A., and G. J. Holland 1999: Large-scale patterns associated with tropical cyclogenesis in the western Pacific. *Mon. Wea. Rev.*, **127**, 2027-2043.