

AWARD NUMBER:

W81XWH-13-2-0011

TITLE:

The Phase of Illness Paradigm: A
Checklist Centric Model to Improve
Patient Care in the Burn Intensive
Care Unit

PRINCIPAL INVESTIGATOR:

LTC Jeremy Pamplin, MD

RECIPIENT:

The Geneva Foundation
Tacoma, WA 98402

REPORT DATE:

April 2016

TYPE OF REPORT:

Annual

PREPARED FOR:

U.S. Army Medical Research and Materiel
Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE April 2016		2. REPORT TYPE Annual		3. DATES COVERED 30 Mar 2015 - 29 Mar 2016	
4. TITLE AND SUBTITLE "The Phase of Illness Paradigm: A Checklist Centric Model to Improve Patient Care in the Burn Intensive Care Unit."				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-13-2-0011	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Dr. Jeremy Pamplin - Jeremy.c.pamplin.mil@mail.mil				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Geneva Foundation 917 Pacific Avenue Suite 600 Tacoma, WA 98402				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We will develop the checklists for the POIP using cognitive systems engineering methodologies and iteration. To validate the model, we will measure clinician perception of patient condition and care priorities; we will observe the frequencies that healthcare teams complete key elements of patient care; and we will measure communication, teamwork, cognitive work load, and provider quality of life before and after units implement the POIP. To assess the model's effect, we will observe patient outcomes and complication rates before and after units implement the POIP. To further assess the model's capacity to rapidly incorporate new knowledge into burn critical care, we will update the phase specific checklists six months after initial implementation and continue data collection.					
15. SUBJECT TERMS Team Communication, Burn Intensive Care, Illness severity, Care Goals, Clinical Decision Support Tools, Phases of Illness, Cognitive Workload, Quality of Life, Card Sorting					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT U	18. NUMBER OF PAGES 163	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

TABLE OF CONTENTS

	<u>Page</u>
1. Introduction	4
2. Keywords	4
3. Overall Project Summary	4
4. Key Research Accomplishments	17
5. Conclusion	17
6. Publications, Abstracts, and Presentations	18
7. Inventions, Patents and Licenses	19
8. Reportable Outcomes	19
9. Other Achievements	20
10. References	22
Appendices	

A. Introduction

This project seeks to validate the phases of Illness paradigm (POIP) (Pamplin 2011) and its effect on a variety of measures in three Burn ICUs. This paradigm describes patients with similar illness severity for which clinicians define standard goals of care, treatment objectives, and specific care tasks. Checklists may help to identify a patient's illness severity and priorities of care as they progress or regress through the continuum of illness during their time in the ICU. Within each "phase of illness" – or range of illness severity – phase-specific checklists may help ensure adherence to local protocols, best practices, clinical guidelines, and specific care bundles. These checklists may help to standardize supportive care elements such as types of monitoring, frequency and type of laboratory assessment, sedation strategies, modes of mechanical ventilation, and physical therapy interventions. Through this standardization, the POIP may create a shared mental model of patient care amongst clinicians in the BICU and thus enhance distributed cognition (Hutchins 2000) and assist the work of the multidisciplinary ICU care team. The objectives of this program are as follows:

- a. Understand the work domain in the Burn ICU in terms of patient condition, patient progress, and dependent clinician behaviors in order to create ecologically valid checklists that support clinician work including decision making according to the Phases of Illness Paradigm.
- b. Validate the Phases of Illness Paradigm and its effect on a variety of measures in three Burn ICUs
- c. Implement the POIP to improve the multidisciplinary burn ICU team's understanding of patient illness severity, daily care priorities, and anticipated care goals.

B. Keywords

Team, Communication, Burn Intensive Care, Illness severity, Care Goals, Clinical Decision Support Tools, Phases of Illness, Cognitive Workload, Quality of Life, Card Sorting

C. Overall Project Summary

This project seeks to understand the work domain in the Burn ICU in terms of patient condition, patient progress, and dependent clinician behaviors in order to create ecologically valid checklists that support clinician work including decision making according to the Phases of Illness Paradigm. We will implement the POIP to improve the multidisciplinary burn ICU team's understanding of patient illness severity, daily care priorities, and anticipated care goals. This project aims to validate the Phases of Illness Paradigm and its impact on a variety of measures in three Burn ICUs. In addition, we aim to further develop the Phases of Illness Paradigm by investigating the ecology of clinical behaviors in the team environment it is meant to support. Using surveys, we will evaluate the perception of the clinicians implementing the checklists on teamwork and communication effectiveness.

The objectives of this program are as follows:

- a. Implement the POIP to improve the multidisciplinary burn ICU team's understanding of patient illness severity, daily care priorities, and anticipated care goals.
- b. Understand the work domain in the Burn ICU in terms of patient condition, patient progress, and dependent clinician behaviors in order to create ecologically valid checklists that support clinician work including decision making according to the Phases of Illness Paradigm.

- c. Validate the Phases of Illness Paradigm and its effect on a variety of measures in three Burn ICUs.

The project tasks are as follows:


TASK ONE: Describe the patient progress through intensive care from patient-centric and provider-centric perspectives. This will include identification of general patient characteristics, provider perspectives, care priorities, therapeutics, activities, and care team goals at various times during a notional patient's progress through intensive care.

TASK TWO: Using the information discovered in task 1, create a representation that maps patient progress through the ICU in the form of checklists that identify patients' and care team goals, objectives, and tasks that are commonly associated with a patient's current condition (i.e. "phase of illness").

TASK THREE: Implement the phases of illness paradigm in three Burn Centers and assess its impact on provider understanding of patient status, care priorities, patient outcomes, and effect on communication, teamwork, quality of life, and cognitive workload. Comparative data for providers and patients will be obtained/initiated throughout the project beginning in month 3.

TASK FOUR: Review and update the Phases of Illness Paradigm (POIP) checklists and assess the time it takes for new checklist items to be reliably completed without new/additional education for the healthcare team.

1. Gantt Chart



Key: red line = Core Site progress; purple line = Houston Site progress; dark blue line = Dallas Site progress

2. Project Summary by Task

TASK ONE. Describe the patient progress through intensive care from patient-centric and provider-centric perspectives. This will include identification of general patient characteristics, provider perspectives, care priorities, therapeutics, activities, and care team goals at various times during a notional patient's progress through intensive care.

Data collection for provider-centric perspectives was completed last year at all three sites. Data regarding the patient-centric progress through intensive care will be more thoroughly evaluated using retrospective data obtained during the study period at each participating institution. While there is no additional data at this time, further analyses of previously collected data remains ongoing and continues to reveal confirmation of findings previously reported as well as new insights.

Indeed, mental models that clinicians use differ widely with respect to patient condition and care priorities. We investigated this using two different “surveys,” card sorting (i.e. the “clinician card sort test” or CCST) and a “clinician understanding survey” or CUS. In both tests, we asked clinicians to identify the illness severity of a patient they were caring for by placing a mark on a horizontal spectrum that represented the continuum from “most sick, could die today” to “least sick, could transfer today.” This scale is shown in figure 1. Figure 2 shows representative examples of two patients for whom multiple clinicians were surveyed (in these examples, the survey was the card sort test).



Figure 1. Scale used by clinicians to rate a patient’s current condition or “illness severity.” The blue line would have been made by a clinician before compiling a card sort or a condition understanding survey. This scale may be divided into 10 equal parts and used to compare a clinician’s perception of patient condition to other clinicians caring for the same patient (see below).




Figure 2. Examples of differences clinician perception of patient condition or illness severity. Clinicians individually marked blank scales as in Figure 1. These two examples are synthesized from multiple individual scales to show the differences in clinical perspective about two different patients. In the top example, five clinicians rated the patient. The average score was 6.4 (range 4-8). In the bottom example, seven clinicians rated the patient. The average score was 2.5 (range 1-5). Both examples represent significant differences in clinical perspective about patient condition that could impact clinical decision making about care goals, care priorities, clinician activity synchronization and planning, and, ultimately, patient outcomes.

Interestingly, but not surprisingly, clinician perception about patient condition is more consistent (aligned) when patients are most sick and when they are least sick. Figure 3. Shows the variability (in terms of standard deviation) of individual illness severity ratings compared to the average illness severity rating for patients whom at least five clinicians rated. The main conclusion from this analysis is that the greatest potential for clinicians to disagree about a patient's condition exists when patients are "middle sick."




Figure 3. Standard deviation of illness severity ratings for comparisons of at least five individual clinician ratings. Variance of clinician perception is greatest in "middle sick" patients (average illness severity scores between 4 and 7). Ranges of individual ratings were as high as 4 points different as in the examples in figure 2.

Perceptions about clinician information use and prioritize to make judgements about patient condition was also collected during the CCST. Although this data continues to be analyzed, the initial reviewed also demonstrated notable differences in how clinicians of different clinical backgrounds (i.e. nurses vs. physicians) and of difference experience levels (i.e. attending vs. resident) use information to make these judgements. A simple method for displaying these differences are word clouds. Although these are not a traditional format in medical literature, social sciences use them to demonstrate thematic differences between groups or to reveal patterns. In the word clouds shown in figures 4 and 5, words size represents the frequency (i.e. picks/subjects) that a card was selected to assess patient condition. Information frequency also changes by illness severity; in other words, some information is prioritized more frequently to assess patient condition if a patient is more vs. less sick (data not shown).

SpO2
Vasopressors

Initial Burn Size
Ventilator Goal/Plan
Wound Care Complexity
CXR
Pulse Ox
Mechanical Ventilation
Rehabilitation Goal/Plan
Diagnosis/Problem
Feeding
Organ Failures
Vital Signs
Mental Status
Open Wounds
Mean Arterial Pressure
BUN/Creatinine
Mode
Electrolytes
Burn Depth
Renal Replacement Therapy
Organ Support
Wounds

Nurses
61 Card Sorts, 56 Patients

[illegible]

Figure 4. Comparison of word clouds generated from attending physician and nurse card sorts. Relative word size represents the frequency that information element was chosen to form an assessment of a patient's illness severity. Larger words were chosen more frequently than smaller words. Attendings appear to focus more on problems, diagnoses, categorical ideas (i.e. organ failures, mental status, ventilator mode), plans and goals to assess patient condition whereas nurses appear to be less consistent of focus, evidenced by smaller, more numerous words, and seem to focus on more data driven information elements (i.e. vital signs, urine output, blood pressure), acuity/workload (i.e. renal replacement therapy, mechanical ventilation and its plan), and gestalt (i.e. "general condition") to make these assessments.



Figure 5. Comparison of word clouds generated from attending physician and resident card sorts. Relative word size represents the frequency that information element was chosen to form an assessment of a patient's illness severity. Larger words were chosen more frequently than smaller words. Interestingly, residents (less experienced physicians) chose more information sources, and more cards overall, than did attending physicians as evidenced by the larger word cloud. Resident physicians seem to have difficulty focusing on any particular information source or concept when making assessments of patient illness severity.

As part of the CCST, clinicians were also asked to describe their patient using key terms. Some key terms were identified during CCST tool development through in depth interviews with clinician experts. These were offered to subjects completing CCSTs for simplicity (i.e. they could circle terms they would use), but clinicians were also permitted to offer their own terms. In general, clinicians chose multiple terms to describe their patients (5+/-2 terms on average). Very few terms had any discriminatory power for identifying patients of any particular illness severity. An apparent pattern does exist (see Figure 6), but it is clear that clinicians in three burn ICUs lack a common language that could help clinical teams perceive patient condition more consistently.

#surveys/group	11	13	16	20	23	15	25	25	13	8	169				
Illness Severity	1	2	3	4	5	6	7	8	9	10	Term Count	Mean Severity	A Priori Strength within Group	Final Strength within Group	
Ready for transfer	91%	38%		5%				4%			17	1.9	0.34	0.34	
Not Sick	9%	8%	19%								5	2.4	0.13	0.13	
Just Monitoring	9%	15%	6%	10%		7%					7	3.1	0.06	0.06	
Getting Better	45%	85%	31%	30%	17%	13%	4%	8%			36	3.4	0.32	0.32	
Stable	73%	38%	75%	70%	30%	20%	8%	8%			53	3.6	0.27	0.27	
Improving	55%	77%	69%	30%	39%	7%	12%	16%	8%		51	3.8	0.34	0.34	
Going in the right direction	55%	69%	56%	50%	48%	20%	16%	12%	15%		57	4.1	-0.37	0.15	
Normotensive	36%	15%	44%	30%	22%	27%	28%	8%		13%	38	4.7	-0.13	-0.13	
Stabilized	27%	31%	25%	40%	30%	53%	28%	8%	8%		44	4.8	0.02	0.02	
Medium Sick	18%	8%		35%	30%	20%	12%	8%			25	4.9	0.12	0.12	
Ventilated	9%	15%	44%	55%	61%	53%	72%	60%	69%	38%	88	6.2	-0.22	-0.22	
Infected	9%			20%	17%	27%	36%	24%	38%	13%	34	6.7	-0.03	-0.03	
Critically Stable			6%	5%	43%	40%	64%	48%	15%	13%	49	6.7	0.05	0.05	
Getting Worse				5%		7%	12%	20%	23%	13%	14	7.7	-0.14	0.14	
Profound lung injury		8%		5%		7%	4%	20%	38%	38%	17	7.9	-0.27	0.22	
3+ sick					4%	7%	4%	12%	23%	13%	10	7.9	-0.12	0.12	
Profoundly ill				10%		7%	28%	40%	85%	25%	33	7.9	-0.38	0.38	
Most Interventions				5%		7%	16%	24%	23%	50%	19	8.1	0.21	0.21	
Multiple Organ Failure				5%	4%	7%	4%	24%	23%	50%	17	8.1	0.23	0.23	
In Shock							4%	4%	31%		6	8.5	0.10	0.10	
The sickest patient						7%	4%	28%	38%	88%	21	8.8	0.39	0.39	
Shock Like								4%	23%	13%	5	9.0	0.11	0.11	
Actively trying to die								4%	8%	25%	4	9.3	0.09	0.09	

Figure 6. Contour plot showing the frequency of terms chosen according to patient illness severity as indicated by clinicians. Percentage = the count of a term divided by the number of surveys/illness severity score. Also shown are the number of times a term was chosen overall (**term count**), the mean illness severity for which a term was chosen (**mean severity**), and the difference between the frequency a term was chosen within the A Priori Group (least sick = 1-3, middle sick = 4-7, most sick = 8-10) and Final Group (where most sick = 7-10) vs. outside the group (**strength within group**). This statistic helps differentiate terms that are used often (i.e. by many clinicians) and are consistent with a particular illness severity group. Of note, illness severity in reality is not discrete: it represents a continuum of patient condition. Here, we try to demonstrate that with a gradient of color whereas in the analysis we uses three discrete groups.

Patterns of care priorities were discussed with and described by clinicians during the second group interviews that were designed to validate the POIP model. As shown below in figure 7, clinicians alter care priorities according to their perception of patient condition. Laboratory evaluation, monitoring, mechanical ventilation, venous access, and medication management (i.e. ensuring the patient is on the correct medications) are all more prioritized in sicker patients whereas optimizing nutrition, sleep, wound care, and rehabilitation are all more prioritized in less sick patients. After evaluating patient cases, clinicians identified patient condition and then rated care elements according to highest, middle, and least priority. Highest priority was given a numerical rating of 8, middle priority a rating of 5, and lowest priority a rating of 3. Priorities were then averaged for all clinicians evaluating a patient as the same level of illness (not necessarily grouped by case). Thus, the representation in figure 7 depicts the clinician perception of care priority according to patient illness severity, not according to the patient case presented. Some care elements may have a bimodal importance (i.e. mechanical ventilation, venous access, and wound care). When discussed, this pattern generally reflected the following positions: when patients are “sicker” mechanical ventilation is added or increased in support, when they improve mechanical ventilation is weaned or discontinued. Similarly, venous access is added and removed when patients worsen or improve respectively. Finally, wounds must be thoroughly evaluated when patients worsen and wound care optimized when patients are improved/improving to prevent complications – also, patients tend to be more perioperative in middle illness categories. Renal replacement therapy seems to be generally important whenever it is utilized for any degree of patient illness. Similarly, analgesia and sedation, while not the highest priority at any time, remains a consistently priority during all phases. Summing all of the priorities, even with relative scores, produced a rank-ordered list of overall care priorities of clinicians in the BICU:

1. Wound Care
2. Renal Replacement Therapy
3. Vascular Access & Monitoring
4. Medication Management
5. Laboratory Evaluation, Analgesia, & Sedation
6. Mechanical Ventilation
7. Nutrition

8. Sleep
9. Rehabilitation

SOI	LABS	MON	MV	ACC	MEDS	CRRT	A&S	NUT	SLEEP	WC	REHAB
Red	8	8	7	8	8	7	6	3	4	4	3
	8	7	8	7	7	8	6	5	5	6	3
	5	8	3	5	3	5	3	5	5	5	3
	8	8	8	8	5	8	6	4	3	4	3
	7	6	8	7	7	5	7	6	6	7	4
Orange	8	8	8	8	8	3	5	3	5	8	3
	8	8	8	5	8	8	5	5	3	5	3
Yellow	6	6	6	7	5	7	4	7	6	6	5
	5	8	4	6	6	6	6	7	6	6	6
	6	7	8	7	8	6	6	7	6	6	6
	7	5	5	6	6	8	5	5	7	8	6
Green	5	5	3	5	5	5	7	7	7	8	8
	3	3	8	5	5	8	5	8	5	3	8
	4	4	3	8	7	8	7	7	5	8	7
	3	3	3	4	5	4	7	4	7	7	6
	3	6	3	3	5	3	5	4	4	6	5
	3	3	3	3	3	5	5	3	3	8	5
	3	3	3	4	4	3	5	4	5	7	4

Figure 7. Red = Most sick, orange = “high middle sick,” yellow = “low middle sick,” and green = least sick. Dark blue = highest priority (score of 8), pale blue = middle priority (score of 5), light grey = least priority (score of 3). Numerical values represent average score of all clinicians who participated in group interviewed. An 8 indicates that all clinicians felt a care item was highest priority for that level of illness, whereas a 3 indicates it is of least priority. SOI: Severity of Illness; LABS: lab draws; MON: Monitors; MV: Mechanical Ventilation; ACC: Venous Access; MEDS: Medications; CRRT: Continuous Renal Replacement Therapy; A&S: Analgesia and Sedation; NUT: Nutrition; SLEEP: Sleep; WC: Wound Care; REHAB: Rehabilitation.

Summary of Findings

1. Multiple measurements demonstrate that mental models of patient condition and treatment priorities are highly variable, and change according to:
 - a. patient condition,
 - b. institution,
 - c. profession or clinician type,
 - d. and years of experience.
2. Consistent patterns of information use, descriptive terminology, and treatment preferences exist that could be used to help create explicit models of patient condition and therapeutic appropriateness.
3. These patterns, however, are spread along a condition versus in discrete phases. Therefore, clinician perceptions are often overlapping and, at times, at odds with each if individual clinician perspective is at distant locations on this continuum compared to other clinicians.
4. Tools can be made to support explicit models of these patterns that should facilitate clinician communication, better decision making, and consistency of patient care.
5. These findings were presented at regional and national conferences (appendix A) and manuscripts (appendix B) of these findings are currently in draft.

TASK TWO. Using the information discovered in task 1, create a representation that maps patient progress through the ICU in the form of checklists that identify patients' and care team goals, objectives, and tasks that are commonly associated with a patient's current condition (i.e. “phase of illness”).

Each site has created tools that are similar in function, but differ in design and content (appendix C). These tools are:

1. A patient illness severity assessment tool.

2. A checklist tool that matches treatment options to patient condition.
3. A “TeamView” tool that helps clinicians to visually identify patient condition and care goals.

While each of these tools have similar function, they differ slightly at each of the participating locations. These pragmatic differences stem from the cultures, personalities, other systematic processes, and leadership at each organization. For example, the TeamView at the Dallas site incorporated a diagram of the wounds a patient has, but this does not appear on either the Houston or USAISR TeamViews because these locations have other mechanisms for sharing that information. Both illness severity assessment and checklist tools also vary somewhat in that each participating location. These differences help improve the ecological validity of the model these tools support. Whether or not these differences are clinically important remains to be determined. Ultimately, identifying these differences, categorizing them, and, if possible, identifying “best” options will be a significant contribution of this project to the science of checklist creation and process improvement.

One major difference in the development and implementation of these tools is worth highlighting. Tool development at Dallas site has been challenging for a variety of reasons: change in research team, physical move of the hospital to a new location, etc. Their version of tools are notably different than those at Houston or the USAISR. While the illness severity assessment tool is similar, they have chosen to create discrete checklist for different phases of patient illness (more similar to the original concept of POIP described by Pamplin, et. Al) instead of showing these checklist items along a continuum as was done at the other sites. It will be interesting to compare their tool use and its impact on patient care to that from the other locations.

Also different at the Dallas site was a major challenge from their hospital leadership regarding the acceptability of posting the TeamView outside of patient rooms. While this practice was not an issue at the USAISR or Houston locations. Indeed, The Joint Commission inspection highlighted these tools as a “best practice” for communication while visiting the USAISR location. Nevertheless, the Dallas site is not showing the TeamView in site for the entire team to see. Instead, these are contained in a binder at the patient bedside which must be opened and interacted with. Again, these differences will be noted and described to help identify best practices to facilitate tool use.

Finally, it is worth noting that using these tools changes how clinicians interact and perceive patient care. Observations of this are below (task three) and will be further described as we continue to analyze data. One observation, however, is clear: clinicians find completing and maintaining these tools to be time consuming and, at times, tedious. This is a similar finding in other checklist projects. To reduce this burden, we intend to create software that facilitates tool completion if the project is approved for a one year, no cost extension. This software will be digital representation of the paper tools currently used, but will help clinicians through prompts and interface to more rapidly assess necessary updates. Future work could integrate such a tool with electronic medical records to further reduce task burden for the clinician.

Summary of Findings

1. Each site created tools to support the POIP model. Tools have similar functions, however they vary somewhat in form and content. These differences account for the differences in clinical cultures and expectations at each organization and improve the ecological validity of the tools for the work they support.
2. Consistent tool completion, even if valuable to the team, remains challenging at all locations because they are an additional burden to individually overburdened, time pressured, task saturated clinicians.

TASK THREE. *Implement the phases of illness paradigm in three Burn Centers and assess its impact on provider understanding of patient status, care priorities, patient outcomes, and effect on communication, teamwork, quality of life, and cognitive workload. Comparative data for providers and patients will be obtained/initiated throughout the project beginning in month 3.*

Numerous delays in implementation at the Houston and Dallas site occurred. Both sites have changed research coordinators during the course of the study. The Dallas site moved into a new hospital in 2015.

Also, the Dallas site has had some difficulty implementing the tools because, despite clinician preferences and empiric data recorded as a part of this project, the Dallas hospital leadership was reluctant to post the TeamView tool outside of patient rooms due to concerns about violating the Health Insurance Portability and Accountability Act (HIPAA). Leadership at the USAISR and Houston sites did not perceive this to be a problem because no the TeamView does not contain any personally identifiable information (PII). Furthermore, the Joint Comission surveyed the USAISR site in 2015 and identified the TeamView tool as a “best practice” for team communications. Despite these arguments, the Dallas site was unable to convince leadership to post the TeamView so that it’s information would be available at the patient bedside for all team members, including patient family and friends, to see.

After education of the staff, tools were incorporated into clinical workflow at each of the three participating centers:

- USAISR on May 2014
- Houston on April 2015
- Dallas on November 2015

Clinician perceptions about patient condition were measured using the clinician card sort test and the condition understanding survey (CCST and CUS respectively, see task 1) at the beginning of the study (pre-baseline time point). These perceptions were subsequently measured before implementation of the process improvement (i.e. before tools were introduced in the the clinical environment) and then at six months and one year (at the core site only, data at spoke sites is still being collected). Data for the core site about clinician perception of patient illness severity from the CUS is shown in figure 8 below. Comparisions at one year and from spoke locations at mid point evaluation will be analyzed soon. The core site data demonstrate that discussions about patient condition that occurred through tool development improved clinian agreement about patient condition and that introduction of POIP tools further improved clinician agreement. This suggests that the POIP improves the clinical team’s shared mental model of a patient.




Figure 8. The variance in clinician perception at three time points in from the core study location. Each data point represents the standard deviation of reported severity of illness scores for patients whom at least five surveys were returned. The average variance in these standard deviations was: pre-baseline, ± 1.73 and ± 1.37 ; baseline, ± 1.38 and ± 1.33 ; six months, ± 0.7 and ± 0.76 . The decreased variance for both current and anticipated condition was statistically significant at each time point (ANOVA, $P = 0.01$)

To assess workload perception we used the National aeronautics and space administration task load index (NASA-TLX). We used the TeamSTEPPS perception questionnaire (TPQ) to assess teamwork perceptions among staff. And the POIP User Satisfaction Survey to assess clinician opinions about the POIP tools and model.

NASA-TLX

The pre-baseline NASA-TLX is a tool developed to assess cognitive workload across six scales: mental, temporal, performance, effort, frustration and physical. Baseline (pre-implementation) NASA-TLX data from all three sites assessed workload perception differences among clinician types, years of experience, institution and time spent with a patient. We sought to characterize clinician subjective sense of workload when performing two tasks.

1. Identify if the patient is better, same, or worse than yesterday (severity of illness (SOI)).
2. Identify the most important objectives of care for the patient today (priorities of care (POC)).

After multidisciplinary rounds (MDR) on one patient the entire clinical team present was asked to assess their workload perception associated with those two tasks. For statistical analysis, we grouped clinicians into the following groups:

- Student: medical students
- Nurse: all types including registered nurse (RN), licensed practical/vocational nurse (LPN/LVN), clinical nurse specialist (CNS), etc.
- Physician: all attending physicians (burn surgeons, intensivists), fellows and residents.
- Other: all other credentialed providers not represented in any other category.

Summary of Findings

This data and analysis was accepted in manuscript form and is currently in press for the Journal of Burn Care and Research. The Manuscript describes the cognitive work performed by clinicians during MDR in the Burn ICU. The NASA-TLX effectively revealed workload perception differences and similarities in cognitive work associated with completing two critical tasks performed during MDR: identifying a patient's condition (severity of illness, SOI) and prioritizing associated treatments (priority of care, POC). Significant findings include (see PDF of the submitted manuscript for full details in appendix B):

1. Mental demand, temporal demand, performance, and effort were the primary determinants for the cognitive work performed for the identified tasks on MDR with mental demand being greatest;
2. students, nurses, and physicians all had higher perceived total workload for both SOI and POC than "others";
3. students perceived the most effort on rounds and had the most mental demand when identifying POC and significantly more than "others" when identifying SOI;
4. students, nurses, and physicians experienced significantly more temporal demand when identifying POC than did "other" healthcare providers while only nurses and physicians perceived this significantly more than others for identifying SOI;
5. clinicians with the least experience had higher perceived workload when identifying SOI and POC as compared to those with more experience;
6. and some individuals perceived more frustration and physical demand than most others during MDR.

We will analyze the change in clinician cognitive work for these two tasks over time using this same tool at six months and study endpoint at each participating location.

TeamSTEPPS

The TeamSTEPPS-Teamwork Perceptions Questionnaire (T-TPQ) was completed at all three sites to establish baseline teamwork perceptions among clinicians. This tool was specifically designed for health care and has been validated across many different types of clinical settings.

Summary of Findings

The pre-baseline T-TPQ data has been collected at all three sites. We collected 129 surveys (physician: 19, nurse: 87, other: 23). There were no significant differences among clinician groups (nurse, physician, other). There was a high degree of acceptable perception of teamwork with a majority of staff (89-100%) across all sites having mean scores greater than 3 for all subscales. We also established good to excellent internal consistency and reliability for all scales (Cronbach's alpha: .85-92).

These results suggest that teamwork in the BICU is perceived as being better than many other ICU environments and that determining the underlying reason for better perceptions of teamwork in the BICU could be used to improve teamwork in other ICU settings.

This data was presented in abstract form (see appendix A) and a manuscript is in draft. We will analyze the change in clinician perception of teamwork over time using this same tool at six months and study endpoint at each participating location.

USER SATISFACTION

The User Satisfaction Survey is a ten-question survey. Each question assesses clinician perspectives about the POIP as a whole (the tools and overall model taken together) using a five point Likert scale (1 – strongly disagree, 2 – disagree, 3 – neutral, 4 – agree, 5 – strongly agree). For purposes of analysis, we grouped responses into disagree (1 or 2), agree (4 or 5), and neutral (3).

Six-month data was preliminarily analyzed for the core site only and will be presented at the American Burn Association Annual Conference this May. We surveyed 48 end-users: 3 Physicians, 30 Nurses, and 15 other healthcare providers (respiratory therapists, rehabilitation specialist, dietitians, pharmacist, etc.). Mean scores are shown below in figure 9. Overall satisfaction was rated above neutral (3.23 ± 0.98). Physicians were more satisfied than nurses: 100% Satisfaction (n=3) vs. 36% dissatisfaction (n=30).



Figure 9. Mean scores for clinician perception of the Phases of Illness Paradigm and the tools used to introduce this model of patient care to one Burn ICU.

Summary of Findings

1. The POIP checklist tools supported teamwork and communication without interfering with workflow for most participants
2. After 6 months of use, most clinicians were comfortable with the tools yet daily use was lower than expected
3. We used this and other data to update tools (see below) to improve use and, hopefully, user satisfaction.
4. We will analyze the change in clinician perception of the POIP and its tools over time using this same survey at the other participating sites and at all sites at study endpoint.

We are currently developing a retrospective protocol that will assess the impact of these interventions on clinician performance (i.e. adherence to the suggested practice patterns outlined by the POIP tools) and on patient outcomes. This retrospective protocol will be submitted for IRB and HRPO review in the next quarter.

TASK FOUR. *Review and update the Phases of Illness Paradigm (POIP) checklists and assess the time it takes for new checklist items to be reliably completed without new/additional education for the healthcare team.*

The core site updated their tools from March to July 2015. The new tools and workflow were introduced in July 2015 (appendix C). Key finding and changes included:

1. The tools when used, were valuable for team situational awareness, but when not completed consistently led to dissatisfaction with their outputs and little impact on patient care. Users requested to incorporate the tools key features (illness severity assessment and corresponding checklists of care priorities) into their daily workflow. This was accomplished by developing both nurse and physician report tools. Nurses could use these tools during change of shift report and residents could use these tools when reporting for multidisciplinary rounds.
2. The TeamView was perceived to have the most potential benefit for team communication, but it was underutilized. The most important feedback was that the tool was not referenced or reviewed during

multidisciplinary rounds. When it was, the tool was perceived as providing value. This led to a decision that resident physicians and physician assistants would be responsible for updating the TeamView (vice nurses) and emphasis on using the tool during rounds.

3. Team View was modified so that it did not duplicate information found in other areas of the team's workflow. Specifically, the daily checklist items were removed from the TeamView, as these were already available in multiple other locations (the EMR note, the "charge nurse checklist" and the resident handoff tool). Additional areas were added to leave messages, identify patient allergies and code status, and to prioritize/synchronize commonly conflicting care tasks (i.e. procedures vs. wound care vs. rehabilitation vs. imaging).
4. A separate TeamView tool was created for patients on extracorporeal membrane oxygenation (ECMO)

Assessment of the impact of these changes is currently ongoing at the core site for final data collection. Anecdotal observations following the tool updates are: 1. The TeamView tool is updated consistently and is valued by team members for maintain the "whole picture" of a patient. Family members interact with the tool to get a daily update on patient condition. 2. Scales and checklist tool use remain inconsistent. Nursing staff do not use them. Physician staff, primarily residents and medical students, find the tools valuable to get a "gist" of patient progress through the BICU, important care elements, and how to apply these care elements.

The Houston site implemented their tools in March 2015, but due to low patient volume and a change in their site's research nurse, their mid-point assessment and tool update is currently on going (April 2016). The Dallas site implemented their tools in November 2015. They changed research nurses associated with this project between March and April 2016 and will start mid-point assessment and tool updates as soon as possible.

Assessment of tool impact on patient outcome and clinician reliability will be evaluated through retrospective review of electronic medical record data. This protocol is currently being written.

D. Key Research Accomplishments

- Models of clinician perceptions of patient condition and progress through the Burn ICU are described at all three sites.
- Tools that aid clinical assessment of patient illness severity and that help identify important treatment priorities are implemented at 3 sites.
- Differences in hospital and unit level leadership, and unit culture *directly* impact the success of process improvement. Differences in these important system features at each location have led to variability in how tools are implemented, valued, and utilized.
- There are differences in how clinicians think demonstrated by the mental models elicited from the card sort data, the condition understanding data, and through individual and group interviews. There are differences among professions or clinician types, by years of experience, and among institutions. These differences are important to acknowledge as they impact communication, care coordination, and ultimately patient safety and outcomes.
- The process of studying clinician perceptions about patient condition and implementing a model of care that highlights patient condition and care priorities makes clinician perception about patient condition more consistent.
- There are differences among professions or clinician types in workload perception, with statistical significance between:
 - o Students and others for mental and overall effort.
 - o Physicians and others for overall workload.
 - o Nurses and others for overall workload.
 - o Physicians and others for temporal demand.
 - o Nurses and others for temporal demand.

- Mental, temporal, performance and effort all contribute significantly more to the overall cognitive workload than physical or frustration subscales.
- Teamwork perception at all three sites was perceived as highly favorable.
- The T-TPQ is a valid and reliable tool to measure teamwork perception in the burn ICU.

E. Conclusions

The Phases of Illness Paradigm is a valid model for patient progress and care prioritization in the Burn ICU. Differences in clinical perception of patient condition and care priorities impacts patient treatment. When clinicians have different mental models of a patient, care may be fractured, priorities may be different, and friction occurs between clinicians. When clinicians share the same mental model of a patient, care is consistent, synchronized, and clinicians will likely perform better as a team.

Process, workflow, and leadership all play an important role in implementing systematic changes in how clinicians perform work. Even when processes may improve patient care, if they do not match clinician workflow or leadership does not prioritize them, there is marginal adoption of the process, little change in culture, and limited impact on patient care. When process matches workflow and leadership supports change, then tools like the POIP model can impact and change unit culture.

Final data collection and analysis is ongoing for this project at the core site.

F. Publications, Abstracts, And Presentations

Manuscripts

1. McInnis I, Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. "Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs." *Journal of Burn Care & Research* [In Press]

Presented Abstracts

1. Pamplin J, Murray S, Serio-Melvin M, Aden J, Huzar T, Wolf S, Chung K, Nemeth C. "Variations on a theme: How clinician descriptions of patient condition diverge." Presented at the Society of Critical Care Medicine Annual Congress, 21-24 February 2016, Orlando, FL.
2. Murray S, Pamplin J, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S. "Using Focus Group Interviews to Validate Team Communication Tools in the Intensive Care Unit." Presented at the Society of Critical Care Medicine Annual Congress, 21-24 February 2016, Orlando, FL.
3. Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. "Developing Cognitive Tools for the Intensive Care Unit: A Mixed Methods Approach." Presented at the Tri-Service Nursing Research Program Annual Conference, 2015, Bethesda, MD.
4. Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. "Revealing the different perceptions burn ICU clinicians have regarding patient illness severity and clinical treatments using card sort methodology." Presented at the Tri-Service Nursing Research Program Annual Conference, 2015, Bethesda, MD.
5. Murray S, Chung K, Mann-Salinas E, Serio-Melvin M, Huzar T, Wolf S, Nemeth C, Pamplin J. "Developing Cognitive Aids According to the Phases of Illness Paradigm for use in the Burn ICU." Presented at the Military Health Research Symposium, 17-2701 August 2015, Ft. Lauderdale, FL.
6. Brown T, Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. "Perceptions of Teamwork in the Burn ICU." Presented at the Military Health Research Symposium, 17-2701 August 2015, Ft. Lauderdale, FL.
7. McInnis I, Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. "Comparing the Workload Perceptions of Determining Patient Condition and Priorities of

Care Among Burn Providers in Three Burn ICUs.” Presented at the Military Health Research Symposium, 17-2701 August 2015, Ft. Lauderdale, FL.

8. Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. “Comparing the workload perceptions of determining patient condition and priorities of care between burn providers in three burn ICUs.” Presented at the American Burn Association Annual Meeting, Chicago, IL April 2015
9. Pamplin J, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C. “Discovering mental models that burn ICU clinicians' use for decision making using card sorts.” Presented at the American Burn Association Annual Meeting, Chicago, IL April 2015

Accepted Abstracts for Presentation

4. Leazer S, Murray S, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Huzar T, Wolf S, Nemeth C, Pamplin J. “Understanding Clinician Perspectives of Patient Condition and Care Goals in the Burn Intensive Care Unit.” Accepted for presentation at the American Burn Association, Las Vegas, NE, 4-7 May 2016
5. Pamplin J, Murray S, Serio-Melvin M, Aden J, Huzar T, Wolf S, Chung K, Mann-Salinas E. “Getting the Burn Team to play from the same playbook: Understanding clinician perception of patient condition.” Accepted for presentation at the American Burn Association, Las Vegas, NE, 4-7 May 2016
6. Murray S, Phillips S, Rodriguez J, Serio-Melvin M, Aden J, Mann-Salinas E, Chung K, Nemeth C, Pamplin J. “Achieving Ecological Validity: Creating Decision Support Tools for the Burn Intensive Care Unit.” Accepted for presentation at the American Burn Association, Las Vegas, NE, 4-7 May 2016

G. Inventions, Patents And Licenses

Not applicable.

H. Reportable Outcomes

1. The core site has described how clinicians perceive patient condition and progress through the ICU. These perspectives have been organized into a “scales tool” and a “bedside checklist tool.” The final implemented tools and the Core Site in-service are included in appendices B & D and appendix J, respectively. These tools and their development were presented in abstract form at the Military Health System Research Symposium, Ft. Lauderdale, FL, 18-21 August 2014 (previously submitted) and were presented nationally at the Society of Critical Care Medicine Annual Congress, 17-21 January 2014, Phoenix, AZ.
2. Using data collected from the Core and Houston Sites, the project has identified significant differences in the mental models that clinicians use to prioritize information related to patient condition and treatments. Physicians, nurses, physician trainees, respiratory therapists, nutritionists, and clinicians of different experience levels prioritize information and treatment options differently. These results have been presented at the American Burn Association Annual Conference, 21-24 April 2015, Chicago, IL (previously submitted).
3. We have also analyzed the NASA-TLX data from all three participating sites and have described the cognitive workload that clinicians perceive while performing the tasks of identifying patient condition and treatment priorities during multidisciplinary rounds. Clinicians of different experience levels and of different professional backgrounds perceive their workload differently. Decreasing this workload may free cognitive processes to focus on more important decisions. These results were presented to the American Burn Association Annual Conference, 21-24 April 2015, Chicago, IL (previously submitted). In addition, a manuscript on this topic is in press for publication in the Journal of Burn Care and Research (previously submitted).

4. These mental models have been used at the Core and Houston sites have validated the “scales” and “checklist” (now called “TeamView”) tools through focus group interviews. The tools have been produced for use at the core site and have undergone unit level review at all three sites, and pilot testing at 2 sites (previously submitted).
5. We have analyzed the first round of TeamStepps data from all three sites and found that there were no significant differences among the clinician groups, sites, or for experience levels. There was a high degree of acceptable perception of teamwork. The majority of staff (89-100%) across all sites had mean scores greater than 3 for all subscales. We found that the TeamStepps perception questionnaire had good to excellent internal consistency and reliability for all scales (Cronbach’s alpha= .85-, .92). A manuscript explaining this data is in progress.
6. We have analyzed the CUS data, pre-baseline round 1 from the core site, quantifying the text data (thematic analysis). The CUS tool asked all clinicians caring for the same patient to identify how sick their patient was (0-10 scale) and note the goals, objectives, and tasks for the current day and the next day. We identified 169 responses from 60 participants that met the definition of “goal.” We were able to identify that clinical teams have difficulty perceiving common goals for patients, often misidentifying objectives as goals. There was very little goal agreement between clinicians caring for the same patient. (previously submitted).
7. We also analyzed the variance in severity of illness scores (SOIs) on the first 3 rounds (pre-baseline, baseline, and at 6 months of tool use) at the Core site. We found a significant decrease in variance over the study period (ANOVA, $p=0.01$). (previously submitted).
8. We analyzed the user satisfaction surveys at the core site and found that users ($n=46$; Physicians: 3; Nurses: 30; Other: 13) were in agreement that the tools supported teamwork and communication without interfering with workflow. After 6 months of use, most clinicians were comfortable with the tools. The POIP checklist tools are ecologically valid and support the work domain of the burn ICU clinicians.
9. We analyzed the focus group interviews at the core site ($n=28$ participants, 6 groups) finding that clinicians identified SOI according to their perception of patient overall condition and current trajectory. Temporal and historical factors play important parts in not only determining how clinicians think about their patients but also how they prioritize care. Understanding team perception may improve communication and patient safety.
10. We analyzed the card sort surveys and found that the language that ICU clinicians use to describe patients poorly differentiates them according to SOI. Of the 169 clinicians and 77 unique patients (Staff Physician: 25; Nurse: 61; Resident: 40; Others: 43), on average staff physicians chose 5 ± 2 descriptors, nurses and residents chose 4 ± 2 descriptors, and clinicians in other roles chose 6 ± 2 descriptors. Eight percent added novel terms, and only one term was consistent with a specific category (“ready for transfer” with least sick).
11. Using thematic coding we analyzed the baseline Condition Understanding Surveys (CUS) responses identified as “goals” ($n= 60$ clinicians, 169 responses) finding 95 (57%) met the definition of a goal. Among the coded goals, the top 3 were: Adequate perfusion/Monitor Perfusion (19%), Ventilator Liberation (12%) and Infection Treated/Cured (12%). Top goals varied according to patient condition: for Least Sick patients, the top coded goal was to “Transfer to Ward” (29%); for Middle Sick patients, they were “Ventilator Liberation” and to “Improve Function/Rehabilitation Tolerance” (10% each); and for Most Sick patients, it was “Adequate Perfusion/Monitor Perfusion” (28%). Overall goal agreement between team members was 28% +/- 17%. Overall, we found that clinicians had difficulty identifying common goals for patient care and often misidentify objectives and tasks as goals. There was very little agreement in goals between team members.

I. Other Achievements

Nothing to report.

J. Future plans and funding options.

Because of delays in implementation and data collection at all research sites, but primarily at the Houston and Dallas sites, the PI has requested a one year no cost extension (NCE) to complete work. The Houston Site is in the process of updating tools and workflow. The Dallas site will do so shortly with a new research coordinator. We will complete final data collection and analysis after sites complete at least three months of tool use.

A retrospective protocol that assess the POIP's impact on patient outcome and clinician reliability is being written. This protocol will review electronic medical record data and compare patient condition identified electronically to POIP conditional assessments, evaluate variance in patient therapies that are recommended by the POIP checklists, consistency of POIP tool completion, and overall patient outcomes.

The core site has nearly completed final data collection. As this occurs, this site will start to implement new clinical practice guidelines that incorporate patient illness severity assessment and targets management strategies accordingly for mechanical ventilation and for pain, agitation, delirium, and sedation.

Because of savings in research coordinator costs at the core site, we are able to fund a software engineer to create a digital version of the POIP tools (illness severity assessment tool, corresponding checklists, and the TeamView) if the NCE is approved. We anticipate that this type of technology can help facilitate POIP implementation and enhance workflow, primarily by reducing clinician time.

K. References

1. Ahmed et al. The effect of two different electronic health record user interfaces on intensive care provider task load, errors of cognition, and performance. *Crit Care Med* (2011) vol. 39 pp. 1626-1634
2. Alvarez and Coiera. Interdisciplinary communication: an uncharted source of medical error?. *J Crit Care* (2006) vol. 21 (3) pp. 236-42; discussion 242
3. Bosk et al. Reality check for checklists. *Lancet*. 2009;374:444
4. Boyle and Kochinda. Enhancing collaborative communication of nurse and physician leadership in two intensive care units. *J NursAdm* (2004) vol. 34 (2) pp. 60-70
5. Catchpole KR, De Leval MR, McEwan A, et al. Patient handover from surgery to intensive care: using Formula 1 pit-stop and aviation models to improve safety and quality. *PaediatrAnaesth*. 2007;17:470-478.
6. Collins et al. Model development for EHR interdisciplinary information exchange of ICU common goals. *International Journal of Medical Informatics* (2010) pp. 1-9
7. Cook RI, Brandwijk M, Kahana M, O'Connor MF, Brunetti VL, Nemeth CP [2003] Being Bumpable: Consequences of Resource Saturation and Near-saturation for Cognitive Demand on ICU Practitioners International Anesthesia Research Society National Conference. New Orleans.
8. Dodek and Raboud. Explicit approach to rounds in an ICU improves communication and satisfaction of providers. *Intensive Care Med* (2003) vol. 29 (9) pp. 1584-8
9. DuBose et al. Measurable outcomes of quality improvement in the trauma intensive care unit: the impact of a daily quality rounding checklist. *J Trauma* (2008) vol. 64 (1) pp. 22-7; discussion 27-9
10. Frankel et al. Using the Communication and Teamwork Skills (CATS) Assessment to measure health care team performance. *Joint Commission journal on quality and patient safety / Joint Commission Resources* (2007) vol. 33 (9) pp. 549-58
11. Gawande A. *The Checklist Manifesto: How to Get Things Right*. New York: Metropolitan Books; 2009.
12. Girard et al. Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial. *Lancet* (2008) vol. 371 (9607) pp. 126-34
13. Halcomb and Davidson. Using the illness trajectory framework to describe recovery from traumatic injury. *Contemp Nurse* (2005) vol. 19 (1-2) pp. 232-41
14. Hales and Pronovost. The checklist--a tool for error management and performance improvement. *J Crit Care* (2006) vol. 21 (3) pp. 231-5
15. Hales B, Terblanche M, Fowler R, et al. Development of medical checklists for improved quality of patient care. *Int. J. Qual. Health Care*. 2008;20:22-30.
16. Hales et al. Development of medical checklists to improve quality of patient care. *Inter J Qual Health Care*. 2008;20:22
17. Hall. Interprofessional teamwork: professional cultures as barriers. *J Interprof Care* (2005) vol. 19 Suppl 1 pp. 188-96
18. Hart, S. G. (2006). NASA-Task Load Index (NASA-TLX); 20 Years Later. *Proceedings of the Human Factors and Ergonomics Society 50th Annual Meeting*, 904-908. Santa Monica: HFES.
19. Hannah S. Sorting out card sorting: Comparing methods for information architects, usability specialists, and other practitioners. 2005. Retrieved on 4 Jul 2013. <http://cardsort.steviehannah.net/Hannah2005.pdf>
20. Haynes et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* (2009) vol. 360 (5) pp. 491-9
21. Henly et al. Health and illness over time: the trajectory perspective in nursing science. *Nurs Res* (2011) vol. 60 (3 Suppl) pp. S5-14
22. Herndon DN and Blakeney PE. Teamwork for total burn care: achievements, directions, and hopes. In: Herndon DN, eds. *Total Burn Care*. 3rd ed. Philadelphia, PA: Saunders Elsevier; 2007:9-13
23. Jones and Hunter. Consensus methods for medical and health services research. *BMJ* (1995) vol. 311 (7001) pp. 376-80
24. Kohn L, Corrigan JM, Donaldson MS, eds. *To Err Is Human: Building a Safer Health System*. Washington, DC: National Academies Press; 1999.
25. Kuwamoto, et al. Process improvement data collection is as simple as asking daily questions using a checklist. *Crit Care Med*. 2010;38(12): 116
26. Levy et al. The Surviving Sepsis Campaign: results of an international guideline-based performance improvement program targeting severe sepsis. *Crit Care Med* (2010) vol. 38 (2) pp. 367-74
27. Lurie et al. Assessing teamwork: a reliable five-question survey. *Fam Med* (2011) vol. 43 (10) pp. 731-4

26. Malhotra et al. Workflow modeling in critical care: piecing together your own puzzle. *J Biomed Inform* (2007) vol. 40 (2) pp. 81-92
27. Miller et al. Interrater variability of a severity-of-illness score in critically ill adults. *Am J Crit Care* (2009) vol. 18 (2) pp. 118-22; quiz 123
28. Narasimhan M, Eisen LA, Mahoney CD, et al. Improving nurse-physician communication and satisfaction in the intensive care unit with daily goals worksheet. *Am J Crit Care*(2006) vol. 15 pp.217-222.
29. Nielsen, J. (2004). Card Sorting: How Many Users to Test. Retrieved on 1 July 2013 from Jakob Nielsen's Alertbox: July 19, 2004. <http://www.nngroup.com/articles/card-sorting-how-many-users-to-test/>.
30. Nemeth et al. Using Cognitive Artifact to Understand Distributed Cognition. *IEEE Transactions on Systems, Man, and Cybernetics, Part A.* (2003) vol. 34 pp. 726-735
31. Nemeth et al. Getting to the point: developing IT for the sharp end of healthcare. *J Biomed Inform* (2004) vol. 38 pp. 18-25
32. Nemeth, C., Brandwijk, M., and Cook, R. (2003). A Study of Pediatric Intensive Care Unit Technical Work. Departments of Pediatrics and Anesthesia and Critical Care. The University of Chicago. Available from the Cognitive Technologies Laboratory, 5841 S. Maryland Avenue, MC4028, Chicago, IL 60637.
33. Newkirk M, Pamplin JC, Kuwamoto R, Allen DA, Chung KK. Checklists change communication about key elements of patient care. *J Trauma.* [In Press]
34. Pamplin J, et al. A daily checklist can change intensive care unit hand hygiene culture. *Crit Care Med.* 2010;38(12): 525
35. Pamplin JC, et al. Phases-of-illness paradigm: better communication, better outcomes. *Critical Care* 2011, 15:6;509
36. Pronovost et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med* (2006) vol. 355 (26) pp. 2725-32
37. Pronovost et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med.* 2006;355:2725
38. Pronovost et al. Improving communication in the ICU using daily goals. *J Crit Care* (2003) vol. 18 (2) pp. 71-5
39. Reader et al. Interdisciplinary communication in the intensive care unit. *Br J Anaesth* (2007) vol. 98 (3) pp. 347-52
40. Schweickert et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet* (2009) vol. 373 (9678) pp. 1874-82
41. Sexton et al. Error, stress, and teamwork in medicine and aviation: cross sectional surveys. *BMJ* (2000) vol. 320 (7237) pp. 745-9
42. Shortell et al. Organizational assessment in intensive care units (ICUs): construct development, reliability, and validity of the ICU nurse-physician questionnaire. *Med Care* (1991) vol. 29 (8) pp. 709-26
43. Stone, WC. Design of fully redundant autonomous life support systems. In: Mitchell, CT (eds.) *Diving for Science 86. Proceedings of the American Academy of Underwater Sciences Sixth Annual Scientific Diving Symposium.* Held October 31 - November 3, 1986 in Tallahassee, Florida, USA. American Academy of Underwater Sciences.
44. Verdaasdonk et al. Requirements for the design and implementation of checklists for surgical processes. *SurgEndosc* (2009) vol. 23 pp. 715-726
45. Vincent. We should abandon randomized controlled trials in the intensive care unit. *Crit Care Med* (2010) vol. 38 (10 Suppl) pp. S534-8
46. Weiss et al. Prompting Physicians to Address a Daily Checklist and Process of Care and Clinical Outcomes: A Single-Site Study. *American Journal of Respiratory and Critical Care Medicine* (2011)
47. Winters et al. Clinical review: Checklists – translating evidence into practice. *Crit Care.* 2009;13:210
48. Zevola et al.Using clinical pathways in patients undergoing cardiac valve surgery. *Crit Care Nurse* (2002)vol. 22 pp.31-9, 44-50.



Discovering mental models that burn ICU clinicians use for decision making using card sorts.

Jeremy C. Pamplin, MD, FCCM, FACP¹; Sarah Murray, PhDc, MSN^{1*}; Maria Serio-Melvin, MSN¹; James K. Aden, PhD¹; Elizabeth Mann-Salinas, PhD¹; Kevin K. Chung, MD, FCCM, FACP¹; Todd Huzar, MD²; Steven Wolf, MD, FACS³; Christopher Nemeth, PhD⁴

*** presenter**

1. United States Army Institute of Surgical Research, Fort Sam Houston, TX;
2. Memorial Hermann Hospital Texas Medical Center, Houston, TX;
3. University of Texas – Southwestern Medical Center, Dallas, TX;
4. Applied Research Associates, Inc., Fairborn, OH



Disclosures



The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.


This study was conducted in accordance with protocols reviewed and approved by Institutional Review Boards at the Brooke Army Medical Center, Memorial Herman Hospital Texas Medical Center, and the University of Texas, Southwestern Medical Center.

I have no financial disclosures.

This work is paid for by a DoD Grant (MRMC W81XWH-13-2-0011)



Introduction





Consider this...



- Have you ever experienced the following:
 - Although a plan of care was established, the entire team was not on the same sheet of music?
 - You had a “gut” feeling that your patient was “sicker” (or “less sick”) compared to how the rest of the team was treating him/her but you couldn’t quite explain it?
 - And you certainly could not explain your “feeling” to other team members?
 - That there was a disconnect between how aggressive (or too cautious) the team was being with respect to a plan of care compared to your own “instinct” of what should be done?



Hypothesis



- **Hypothesis:**
 - A novel research method may uncover different mental models that team members have regarding patient illness severity and clinical treatments.



Q Methodology: Card Sorting



- Q Methodology: Developed by psychologist: Dr. William Stephenson (1935)
 - Framework to study subjectivity
 - Quantitative method to evaluate subjective data
- Used in
 - Information Architecture and web design:
used to understand an end-user's point of view
 - Nursing, Veterinary, Medicine, Public Health, Transportation, Education, etc.

Using QMethodology to Identify Reasons for Distress in Burn Survivors Postdischarge

Shelley Wiechman Askay, PhD,* Michael Stricklin, PhD,†
Gretchen J. Carrougner, RN, MSN,‡ David R. Patterson, PhD,*
Matthew B. Klein, MD, FACS,‡ Peter C. Esselman, MD,*
Loren H. Engrav, MD, FACS‡



Card Sorting



- We developed a card sort based on interviews with burn critical care experts (a burn surgeon, intensivist, and nursing staff).
- The final card set included 97 cards:
 - 67 cards in 10 categories of “features” used to judge patient illness severity
 - 30 cards in 9 categories “treatments,” for a total of : 67 features and 30 treatments.



- Card Sort Technique:
- Participants are asked to think about their patient and determine on a scale of 1-10, how sick their patient is
 - 1 = Least Sick, “could transfer today”
 - 10 = Most Sick, “could die today”
- Step 1:
 - Select all the cards that represent features that you consider important to how sick your patient is
 - Cards are then organized in grid
- Step 2:
 - Select all the cards that represent the treatments you think this patient should have
 - Cards are then organized in grid




Methods



- Part of a prospective, mixed methods study of clinical decision making and how to create tools that support it
- 3 academic, regional referral burn ICUs.
- Approved by the institutional review board at each site.



Card Sorting






Example Feature Card Sorts



Sickest Patient/Could Die today (SOI 10)



Least Sick Patient/Could Transfer (SOI 1)





Example Treatment Card Sorts



Sickest Patient/Could Die today (SOI 10)



Least Sick Patient/Could Transfer (SOI 1)





Results





- **169** card sorts, **77** patients
- Average time: **35.5** $\pm$ 15.8 min [10-100 min]
- Average # Feature Cards: **9** $\pm$ 2 cards [3-16]
- Average # Treatment Cards: **8** $\pm$ 2 cards [3-14]



Clinician Perception



- In the diagram below, please use a vertical line to indicate “where” your patient is today:







Perception of Illness Severity



Average Percieved Illness Severity of Patients on whom at least 5 sorts were performed



Standard Deviation of Clinician Perception of Illness Severity by average Perception of Illness Severiy



Word Clouds




Attending Physicians
25 Sorts, 23 Patients

Nurses
61 Card Sorts, 56 Patients



Physician Trainees
40 Sorts, 41 Patients





Limitations

“All models are wrong, but some are useful.” – G. Box



- Subjects review cards created by the research staff during interviews with a small number of clinicians.
 - The mental models may be anchored by the card set available
 - Example: a clinician considered pulse pressure variability as an indicator of volume status but since that card was not specifically available, the subject chooses
 - to pick a card that represented the same concept (e.g. Monitoring: CVP)
 - to opt for a different card/concept altogether or
- Did not investigate the intra-rater reliability



Conclusions



- Burn ICU clinicians have different perceptions about
 - patient condition
 - treatment priorities
- The card sort method successfully elicits mental models from clinicians during routine daily activity
 - These models vary according to patient illness severity, clinician type, clinician experience, and institution



Conclusions



- These differences likely impact effective communication:

What happens when the nurse communicates a different message than the attending to the family?

- Better understanding and awareness of these differences in mental models may improve teamwork.
- Will aid the development of decision support tools.



**Thanks to all of the clinicians who
took part in the card sorts.**

Questions?

- Burn critical care is complex and error prone due to high cognitive workload associated with information overload and miscommunication.
- Multidisciplinary Rounds (MDR) in the Burn Intensive Care Unit (BICU) are the mechanism for reviewing patient status and planning care in the burn intensive care unit (BICU).
- The cognitive work associated with MDR is often unrecognized.
- The National Aeronautics and Space Administration Task Load Index (NASA-TLX) assesses workload on 6 scales: Mental, Temporal, Performance, Effort, Frustration, & Physical.

Objectives

- We sought to characterize clinician subjective sense of cognitive workload while completing 2 tasks:

Task 1: Identify[ing] if the patient is better, same or worse than yesterday (Severity of Illness); and

Task 2: Identify[ing] the most important objectives of care for the patient today.

Methods

- After consents were obtained, research staff at 3 regional referral centers administered the NASA-TLX to the multidisciplinary team:
- Immediately after MDR on a single patient
- Using paper or electronic formats (tablet)
- At different points in time; early, middle or late in the MDR activity
- A total of 5 patient MDR's were assessed at each site
- Directions were read aloud to the entire team
- The MDR groups were defined as:
 - Student: medical students
 - Nurse: all types including RN, LVN, and master's prepared (Clinical Nurse Specialist, MSN)
 - Physician: all attending physicians (burn surgeons, intensivists), fellows and residents
 - Other: All credentialed clinicians not represented in any other category
- Statistics: One-way ANOVA

- The data represent 220 clinicians, 5 patient MDR's, across 3 sites including 27 staff physicians, 25 nurses, 25 residents, 37 in other roles, and 13 students.
- The sample reported moderate difficulty (weighted cognitive load= 42) for both tasks.
- Mental, temporal, performance, and effort all contributed significantly more to the overall cognitive load than physical or frustration subscales.

Task 1: Identify if the patient is better, same or worse than yesterday

Table 1: Cognitive Workload by Profession

★Significant Comparisons:
All comparisons: p < 0.05





Table 2: Cognitive Workload by Years of Experience

Significant Comparisons:
• A, B, G, H = p < 0.0001
• C, D = p< 0.05
• E, F = p< 0.005



Task 2: Identify the most important objectives of care for the patient today

Table 3: Cognitive Workload by Profession

Significant Comparisons:
• A = p< 0.05
• B, C= p<0.005
• D p=0.0006
• E, F, G = p<0.0001





Table 4: Cognitive Workload by Profession

Significant Comparisons:
• A, B, C, G, H, I = p<0.005
• D, E = p< 0.05



- Mental demand, temporal demand, and performance were the primary contributors to overall cognitive workload.
- Providers with less experience reported less workload when identifying patient status than those with more experience.
- Students exert more cognitive effort to identify SOI and P.
- Non-physician, non-nurse reported less workload identifying patient status than physicians and nurses.
- Physicians and nurses reported more temporal demand than physician/nurses.

- The work of identifying patient status and priorities varies according to experience level, but not by institution or patient.
- Identifying patient status and priorities can affect workflow, decrease teamwork.
- Understanding how workload differs among professions can help improve workflow and performance.
- The NASA-TLX is a valid measure of workload in the BICU.

Nicole W. Caldwell, RN, is an unparalleled regulatory specialist. Agnes Burris, RN, Senior Clinical Nurse Specialist at UT Southwestern Medical Center, and Burn Research Coordinator, thank you for your hard work and dedication.

This project is supported by the Department of the Army and Materiel Command Institutional Review Board and in accordance with approved protocol.

for patients in Burn Intensive Care Units (BICUs)

- Team care is challenged by communication lapses borne from differences.
- Professional silos may produce discordant care.
- Well designed, ecologically valid cognitive aids help clinicians make decisions more efficiently, reliably, and accurately and may improve patient care.
 - Checklists, clinical pathways, order-sets, protocols, and guidelines are examples of cognitive aids that improve outcomes in healthcare.
- The Phases of Illness Paradigm (POIP) is a theoretical framework that intends to describe patients with similar severities of illness for which clinicians may define expected goals, objectives, and tasks of care.
- This research was designed to
 - Validate the POIP framework as shared mental model
 - Develop ecologically valid cognitive aids to support the POIP

- To understand the BICU work domain in terms of patient condition, patient progress, and clinician behaviors in order to create ecologically valid cognitive aids.
- To improve the multidisciplinary Burn ICU team's understanding of patient condition, daily care priorities, and anticipated care goals.
- To validate the POIP as a shared mental model.

Secondary

- To further develop the POIP by investigating clinical behaviors in the environment it is meant to support.

Exploratory

- To determine if a shared mental model will improve clinician perception of communication, teamwork, work satisfaction, and cognitive workload in the burn ICU.

- Shared mental models may improve team priorities.
- Although patient condition is a continuum, more discretely along that continuum and
- Tools may improve recognition of discordant perspectives which may foster improved c

Key I

- The described research will develop ecologically valid clinical decision making in the BICU.
- We anticipate that the POIP will:
 - decrease cognitive load
 - improve communication
 - make care more reliable

Methodology

This Institutional Review Board approved protocol includes mixed methods, participatory research project broken down into four main tasks:

Task 1: Describe a Patient's progress through the BICU

- Condition Understanding Survey
- NASA-TLX Survey
- TeamSTEPPS Survey
- Clinician Card Sort: Q Methodology
- Group Interview

Condition Understanding Survey:

- How sick is your patient today?
- What are the top 3 Goals, Objectives, and Tasks today?
- How sick will your patient be tomorrow?
- What are the top 3 Goals, Objectives, and Tasks for tomorrow?

Task 2: Develop Representations of Patient Progress

- Delphi Consensus Building
 - The elements from the card sorts and group interviews were correlated with severity of illness scores on the scales.
 - 80% consensus was achieved on each of the elements for the final version.
- Group Interview

Task 3: Implement the POIP in the BICU

- Clinicians perceive patient condition along a continuum
- Two tools emerged from the data:
 - A "scales tool" that is granular and identifies clinician perception of patient status along the continuum of care.
 - A "checklist" tool for discrete data elements of care and to provide a summative report.

Acknowledgements

- The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense
- This study was conducted under a protocol reviewed and approved by the US Army Medical Research and Materiel Command Institutional Review board and in accordance with approved protocol
- We would like to thank Nicole Caldwell, RN for her instrumental assistance in facilitating this project.
- Funding: This project is supported by a grant from the US Army Medical Research and Materiel Command Center (TATRC) (W81XWH-13-2-0011);

- The burn intensive care unit (BICU) is a data dense environment where clinicians from many disciplines work together to provide care
- Multidisciplinary rounds (MDR) is a collaborative activity designed to discuss patient data and formulate care plans
- MDR is associated with better outcomes in the ICU^{1,2,3}
- Cognitive work associated with participation in MDR in the BICU has not been described
- The National Aeronautics and Space Administration Task Load Index (NASA-TLX) is a well-validated tool that quantifies workload associated with task completion on 6 subscales: Mental, Temporal, Performance, Effort, Frustration, & Physical⁴
- Higher workload, as measured by the NASA-TLX has been associated with poorer outcomes in the healthcare setting^{5,6}

Objectives

- We sought to describe perceived clinician workload during MDR as measured by the NASA-TLX when completing 2 key tasks:

Task 1: *Identify[ing] if the patient is better, same or worse than yesterday (Severity of Illness, SOI)*

and

Task 2: *Identify[ing] the most important objectives of care for the patient today (Priority of Care, POC)*

Methods

- This study was approved by an institutional review board
- It was conducted at 3 regional referral BICUs
- NASA-TLX was administered to a convenience sample of clinicians who participated in MDR after informed consent was obtained
- Participants included nurses, physicians, students and other providers (Dietitians, Respiratory Therapists, Pharmacists, Rehabilitation Therapists, etc.)
- Scores were analyzed using the nonparametric Wilcoxon's Test
- Significance was established when $p < 0.05$

- A total of 116 completed surveys were collected from a group that included 41 physicians, 25 nurses, 13 students and 37 other providers
- Median total load for SOI was 40 (IQR:13-67), while median total load for POC was 43 (IQR:18-68)
- Mental demand, temporal demand, performance, and effort were principle contributors to the overall cognitive load
- Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15




Figure 1. Task 1, Severity of Illness, NASA-TLX Subscale and Total Load Compared Among Profession Subtypes




Figure 2. Task 2, Priority of Care, NASA-TLX Subscale and Total Load Compared Among Profession Subtypes




Figure 3. Task 1, Severity of Illness, NASA-TLX Subscale and Total Load Compared Among Experience Groups




Figure 4. Task 2, Priority of Care, NASA-TLX Subscale and Total Load Compared Among Experience Groups

Significant between-group differences for figures 1-4 above indicated by brackets ($p < 0.05$)

This study was conducted under a protocol reviewed and approved by the US Army Medical Research and Materiel Command Institutional Review board and in accordance with approved protocol

- There are differences in perceived workload among members of the healthcare team
- Key Findings:
 1. Mental demand, temporal demand, performance, and effort were principle contributors to the overall cognitive load
 2. students, nurses, and physicians perceived total workload to be higher than "others"
 3. students perceived workload to be higher than POC and significantly higher than identifying SOI
 4. students, nurses, and physicians perceived workload to be significantly higher than identifying POC and significantly higher than identifying SOI
 5. and clinicians perceived workload to be significantly higher than POC as compared to identifying SOI
- Identifying those with the highest workload to highlight those who may benefit from intervention
- The NASA-TLX tool was used to measure workload in the BICU

Acknowledgments

- Nicole W. Caldwell, RN, MSN, MBA
- Agnes Burris, RN, Senior Lecturer, MBA at UT Southwestern
- Adrian Botello, CSTR, TRMC, Memorial Hermann Hospital, project
- This project is supported by the Department of Research and Materiel Command, Technology Research Center

1. Checkley W, Martin GS, Brown J, et al. JAMA. 2000;283:1061-1067.
2. Zimmerman JE, Shortell SM, et al. JAMA. 2000;283:1068-1074.
3. Lane D, Ferri M, Lemaire J, et al. JAMA. 2000;283:1075-1081.
4. Hart SG, Staveland LE. In: P. A. Hancock, J. T. Warr, eds. Workload. Amsterdam: North-Holland; 1991:139-156.
5. Yurko YY, Scerbo MW, Prabhu AB, et al. JAMA. 2000;283:1082-1088.
6. Mazur LM, Mosaly PR, Hoyle J, et al. JAMA. 2000;283:1089-1095.

Introduction

- The Burn Intensive Care Unit (BICU) is a complex care environment requiring a large multidisciplinary team to achieve optimal patient outcomes.¹
- Good teamwork may improve patient outcomes.²
- The Team Strategies and Tools to Enhance Performance and Patient Safety (TeamSTEPPS) is a validated survey to assess perceptions of teamwork amongst clinicians.³
- In this multicenter study we sought to measure teamwork perception at three burn center ICUs.

Objectives

- Measure perceptions of teamwork among different clinicians in three different BICUs

Methods

- We administered the TeamSTEPPS-Teamwork Perceptions Questionnaire (T-TPQ) to clinicians in three American Burn Association verified regional referral BICUs as part of an Institutional Review Board approved study.
- Admission rates from each site ranged from 270 to 300 patients per year, with an average daily census between 2 and 5.
- Clinicians surveyed included: physicians, nurses, and “others” (therapists, pharmacists, and staff not part of any other group). (Figure 2).
- Experience data collected and grouped as <5 years, 5-10 years, and >10 years.
- Scoring: A total score was calculated for each teamwork construct. Teamwork scores were dichotomized as acceptable for scores greater than 3 and needs improvement for scores less than 3.
- Wilcoxon’s nonparametric method for comparisons was conducted to compare the clinician types, sites and years of experience.

Results

- Surveys were returned from 129 clinicians across all sites (19 physicians, 87 nurses, and 23 others).(Figure 1)
- We found no significant difference between clinician groups, sites, or years of experience in perceived teamwork.
- Clinicians had a high degree of perceived teamwork (Table 1)
- The T-TPQ had good to excellent internal consistency and reliability for all scales (Cronbach’s alpha: .85-.92, Table 2).

Table 1. Means Scores and Standard Deviation of Each Perceived Teamwork Construct with Standard Deviation Based on Clinician Type

Clinician Type	Team Structure	Leadership	Situation Monitoring	Mutual Support	Communication
Physicians	28.6 ± 3.68	28.45 ± 3.28	28.95 ± 4.17	30 ± 3.83	29.1 ± 3.88
Nurses	27.34 ± 4.21	26.77 ± 5.23	27.25 ± 3.46	28.41 ± 3.86	28.65 +/- 3.38
Others	27.71 ± 4.6	28.33 ± 5.66	28.83 ± 3.9	27.83 ± 4.45	29 ± 3.16

Table 2. T-TPQ Cronbach's Alpha Reliability Coefficients

Teamwork Construct	Cronbach’s Alpha
Team Structure	0.83
Leadership	0.90
Situation Monitoring	0.85
Mutual Support	0.82
Communication	0.87




Figure 1. Breakdown of Surveyed Staff Across All Sites



Figure 3. Breakdown of TeamSTEPPS Team Competency Outcomes



Figure 2. Example of multidisciplinary rounds by the burn care team, which includes physicians, nurses, dieticians, physical and occupational therapists, respiratory therapy, wound care coordinators, and social workers.

- This is the first rep perceptions of tea
- Clinicians at all thr perception ratings evaluated.
- The T-TPQ is a va perception in the E
- These results see BICU is perceived ICU environments
- Determining the ur perceptions of tea used to improve te outcomes in other

Acknowledgments

- Nicole W. Caldwell, USAISR for her support and guidance
- Agnes Burris, RN, Christopher Tran, Center;
- Adrian Botello, CS Coordinator at Me hard work and de

1. Stout RJ, Salas E. Enh through team training. G Practice. 1997;1:169-18
2. Wheelan SA, Burchill C patients' outcomes in int 527-534.
3. Agency for Healthcare F teamsteps.ahrq.gov.
4. Reader TW, Flin R, Mea communication in the in
5. Deering S, Johnston L a and communication train

- Medical care in the burn intensive care unit is complex and challenging.
- Fifty-five clinicians participated in card sorts that were validated by all clinician types (surgeon, nurse, respiratory therapist, etc.) in 3 group interviews.

- Checklists, support clinical decision making and communication
- Checklist tools must be ecologically valid, they must represent the work domain that they support in order to be effective
- We combined quantitative and qualitative data to create a valid evidence-based tool to enhance communication and provide clinical decision support.

Objectives

- To understand the work domain in the Burn ICU
- To create ecologically valid checklists
- Implement the Phases of Illness Paradigm
- To validate the Phases of Illness Paradigm
- To further develop the Phases of Illness Paradigm

Methods




Figure 1: Methodology Model;Key: G1– Group Interview 1; G2—Group Interview 2; G3—Group Interview 3

- Demographic data also collected
- Delphi: Process to obtain group consensus




Figure 2: Feature Card Sorts

Card Sorts						
77 Patients, 169 Card Sorts						
	Site 1		Site 2		Site 3	
	Patients	Sorts	Patients	Sorts	Patients	Sorts
Unique	19	60	51	73	7	36
Least	6	16	16	16	2	8
Middle	12	25	33	40	5	18
Most	9	19	12	17	5	10

Table 1: Card Sort by Site and Severity of Illness

Sorts by Clinician Type								
	Site 1			Site 2			Site 3	
	Least	Middle	Most	Least	Middle	Most	Least	Most
Total	16	25	19	16	40	17	8	10
Attending	2	8	3	2	2	2	2	1
Nurse	3	7	4	9	23	8	2	3
Other	7	4	6	2	7	5	1	3
Phys Train	4	6	6	3	8	2	3	3

Table 2: Card Sorts by Site and Clinician Type

- Senior staff (>10 years of experience) and key leaders participated in the Delphi process.

Identify the patient's current Treatments				STD Dev on Question for Same Case	Include Scale	Do Not Include Scale	Include Scale with Modification	Question order
Make a noticeable "X" ANYWHERE the scales below that indicates the patient's current treatments. **Text in each section is organized by objective/goal in bold type, recommendations in regular type, and considerations in italic type.				0.76			X	15
1. Labs Goal: Information availability & minimize blood loss Most frequent: Q4-Q6 ABG, VS/G/S/O ₂ , Lactate, CBC, Chem Q12-24 LFT *Pul Tuber, TSS, Coags, Fibrinogen Less frequent: Q12-Q24 ABG, CBC, Chem Q24-Queue: LFT, Coag As Needed/Intermittent: Q00-QMWF: CBC, Chem Queue: LFT, Coag PPS Labs only				0.76				
2. Nutrition Goal: Minimize loss of lean mass Holding Enteral Feeds or Enteral Feeds as able. Consider TPN Enteral Feeds much as possible Troche (20mL/hr) Full Support 1-Enteral 2-TPN Full Support Enteral/TPN Bolus + Supplements				1.01				
3. Monitoring Standard monitoring (T _{core} , S _{uO₂} , RR, NBP) Plus: Maximize knowledge A-Line, CVP, EIT/IO, BOLD, Continuous S _{uO₂} , Abdominal Pressure, TTE/TIC measurement Assure effective Ventilation & Sedation BOLD, A-Line, ZCVP 4E Compatible Decrease NBP measurements overnight Remove Foley				1.88				
4. Rehabilitation None or ROM/Positioning as tolerated Other care is important General Progression: ROM -> Su/TIC -> Dangle/Tit/Stand Rehab (daily) more important Progress to Mark/Walk -> Gym & Discharge Rehab most important, DO NOT DELAY				0.82				

Figure 3: Delphi Tool

- We created two artifacts (tools) based on the data analysis: a summary and a contextual conditional scales tool (Nursing Illness Severity Assessment Tool) posted outside each patient's door. Each site created unique artifacts.
- The Team View is a communal display of coordinating a view of patient severity of illness, plan of care, daily treatment goals, and metrics such as bundle compliance.

Work Flow

- Bedside nurse updates Team View SOI
- Team updates daily treatment goals and plan of care
- Any team member can use the board to improve understanding of patient condition




Figure 5: Team View Workflow

Work Flow

- Bedside nurse assesses patient: determines SOI
- Off-going nurse uses assessment for shift report communication
- On-coming nurse uses assessment to make treatment suggestions




Figure 6: Nursing Illness Severity Assessment Tools

Conclusions

- Mixed methods clinical research in the ICU provides a d a work domain and can produce ecologically valid tools work
- Effective checklists create shared mental models, improve patient outcomes by improving priorities of care

Acknowledgements

Nicole W. Caldwell, RN Research Coordinator at the USAISR; Agnes Burris, RN, Senior Research Nurse, and Christopher Tran, MBA at UT Southwestern Medical Center; & Adrian Botz, RN Research Coordinator at Memorial Hermann Hospital; for their hard work and dedication to this project

- The condition of each patient in the burn intensive care (BICU) is typically complex and requires comprehensive care from a diverse and robust multidisciplinary team
- The care team must communicate the daily care plan effectively so that all team members understand the patient's daily goals
- Care plans built upon a shared mental model provide care that is consistent, efficient, and able to meet the needs of the patient, resulting in good patient outcomes
- Often, individual BICU clinicians caring for the same patient have different perceptions, or mental models, about the patient condition and treatment priorities
- Differences in patient perception by members of the team reflect a lack of common ground and may impact effective teamwork and patient safety
- Card sorts are research tools that reveal human perceptions about information^{1,2}

Objectives

- As part of a larger prospective observational study to create decision support tools and test the Phases of Illness Paradigm³ in three burn ICUs, we sought to understand the different mental models that individual BICU clinicians have about their patients
- We sought to understand what differences in perception exist within the care team to reveal and understand how they might affect care decisions
- The result is an understanding of clinician perception differences that will impact the creation of communication tools to enhance team perception and improve patient care

- 169 card sorts, 77 patients
- Average time: 35.5 ± 15.8 min [10-100 min]
- Average # Feature Cards: 9 ± 2 cards [3-16]
- Average # Treatment Cards: 8 ± 2 cards [3-14]
- Figures 3 and 4 demonstrate differences in clinician perception of SOI on same patient




Figure 3: Differences in clinician perceptions of illness on patient A






Figure 4: Differences in clinician perceptions of illness on patient B

- Perception of Illness Severity see graphs 1 & 2
- Greater distribution amongst clinicians in the middle sick category



Graph 1: Standard Deviation of clinician perception of illness severity



Graph 2: Average scores of patients with at least 5 sorts

- Word Clouds graphically display the mental models see figures 5, 6, and 7
- There are differences amongst different clinician types
- There are differences between clinicians in the same profession see figures 5 and 7




Figure 5: Attending Physicians 25
Card Sorts, 23 Patients



Figure 7: Residents 40 Card Sorts, 41 Patients



Figure 6: Bedside Nurses 61 Card Sorts, 56 Patients

- We can think about methodological differences as differences in the way we think about the world.

- Available cards that are not available
- Small sample size
- May not be representative of all ICUs

- Burn ICU about patient priorities
- The card mental m daily activ
- These mo illness se experienc

- Ac**
- would like to thank
nce in facilitatin
g; This project i
Research and
ed Technology

Methods

- We used Q-Methodology, or card sorting, to elicit how clinicians think about their patients
- We used 97 cards: 67 features and 30 treatments
- Clinicians were asked to identify a patient's condition on a scale from "could die today" to "could transfer today"




Figure 1: Severity of Illness continuous scale

- Cards were selected and arranged by priority




Figure: 2 Example of Card Sort with cards arranged from most to least important

- Terms were analyzed by frequency in category, clinician type, and compared between institutions




Figure 3: Terms were assigned a font size based upon frequency; larger fonts indicated more frequent

1. Hannah, S. (2005). So information architects, From <http://aim.uorego> 6 August 2014
2. Nielsen, J. (1995). App IEEE, 12(1), 98–100
3. Pamplin JC, Murray S, communication, better

- Focus groups are a source of social knowledge, and represent a way to collect data interactively
- We used focus group interviews to validate checklist tools using the Phases of Illness Paradigm (POIP) model
- The POIP is a model that defines patients by Severity of Illness (SOI), aligning patient condition with treatments, with the intent of improving patient care by improving team understanding of patient condition, and team communication
- This study aimed to validate the POIP model and improve team communication through the creation and use of checklist tools

Methods

- This was a prospective mixed methods study in three academic regional referral Burn ICUs
- Convenience sample focus group sessions were conducted using case studies and semi-structured interviews
- The case studies were of patients in 3 SOI categories: most sick (could die today), least sick (could transfer today), and middle sick (all others) see figure 1
- Clinicians identified how sick the patient was along condition scales and rank ordered the treatments they felt were indicated for the patient that day
- The process was repeated using the POIP checklist tools
- We identified the differences between no tool and tool
- We used thematic analysis to identify codes and themes, we then used these data to update the POIP tools

- We found no significant differences in determining SOI with or without tool use (n= 28 participants; 6 groups)
- Although statistical significance was not found, clinically important differences SOI ranking were observed and discussed
 - 34% of clinicians changed their initial SOI score after using the tool
- Clinicians chose SOI rank based on several themes
 - **Overall Picture:** the patient’s entire status, not just one element
 - Includes premorbid condition and patient history
 - **Timing and Tolerance to Treatments:** whether or not the patient should be better at a moment in time
 - Includes the assumption of what the patient ought to be doing at a moment in their hospital stay
- Interviews suggested that the POIP tools might improve care prioritization by improving communication about patient SOI

Figure 1: The Phases of Illness Severity of Illness Scale



Table 1: PRIORITIES OF CARE BY SEVERITY OF ILLNESS

SOI	LABS	MON	MV	ACC	MEDS	CRRT	A&S	NUT	SLEEP	WC	REHAB
Most Sick	8	8	7	8	8	7	6	3	4	4	3
	8	7	8	7	7	8	6	5	5	6	3
	5	8	3	5	3	5	3	5	5	5	3
	8	8	8	8	5	8	6	4	3	4	3
	7	6	8	7	7	5	7	6	6	7	4
Middle Sick	8	8	8	8	8	3	5	3	5	8	3
	8	8	8	5	8	8	5	5	3	5	3
	6	6	6	7	5	7	4	7	6	6	5
Least Sick	5	8	4	6	6	6	6	7	6	6	6
	6	7	8	7	8	6	6	7	6	6	6
	7	5	5	6	6	8	5	5	7	8	6
Highest Priority	5	5	3	5	5	5	7	7	7	8	8
	3	3	8	5	5	8	5	8	5	3	8
	4	4	3	8	7	8	7	7	5	8	7
	3	3	3	4	5	4	7	4	7	7	6
	3	6	3	3	5	3	5	4	4	6	5
	3	3	3	3	3	5	5	3	3	8	5
Middle Priority	3	3	3	4	4	3	5	4	5	7	4
Least Priority											

LEGEND: SOI: Severity of Illness; LABS: lab draws; MON: Monitors; MV: Mechanical Ventilation; ACC: Venous Access; MEDS: Medications; CRRT: Continuous Renal Replacement Therapy; SLEEP: Sleep; WC: Wound Care; REHAB: Rehabilitation.

MOST SICK

MIDDLE SICK

LEAST SICK

HIGHEST PRIORITY

MIDDLE PRIORITY

LEAST PRIORITY

- Clinicians alter c perception of pa
- Using a checklis condition helps c consistently.
- A shared unders care priorities ma planning, and re
- Group interviews establishing end
- Clinicians priorit SOI, further valid use in the ICU s
- Limitations: we c transcribe the se

- Clinicians ident perception of o current trajecto
- Temporal and H parts in determ their patients a
- Understanding communication

Ack

- We would like to thank N assistance in facilitating t
- Funding: This project is s Research and Materiel C Technology Research Ce

- Communication is a fundamental aspect of teamwork.
- Ineffective communication remains one of the leading causes of medical errors.
- People often assume their words convey an intended meaning.
- Descriptive terms that clinicians of different specialties and backgrounds choose are sometimes misunderstood or interpreted incorrectly.
- Effective teamwork requires a *shared mental model* of patient status, team resources, and goals that all team members understand, and, more importantly, *support*.

Objectives

- We sought to learn if there is any correlation between different clinician’s perspective of patient illness severity and the terms they use to describe a patient’s condition.

Methods

- This was a prospective, mixed methods study (survey plus interview) of clinicians in three academic, regional referral Burn ICUs.
- We asked clinicians during normal daily activities to identify terms they might use to describe “how sick” their patient was, or to offer their own terms.
- We also asked clinicians to identify “where” their patient was on a 1-10 scale indicating illness severity where scores of 1-3 indicated “least sick,” 4-7 indicated “middle sick,” and 8-10 indicated “most sick.”
- We describe our findings using a contour plot, average illness severity per term, confidence intervals, and a “strength within group” statistic (the frequency a term is chosen within an group less its frequency chosen in other groups).
- This study was conducted under a protocol reviewed and approved by the US Army Medical Research and Materiel Command Institutional Review board and in accordance with approved protocol

- We performed surveys and interviews with 169 clinicians caring for 77 unique patients: 25 staff physicians (“attendings”), 61 nurses, 40 residents, and 43 clinicians in other roles.
- On average, attendings chose 5±2 descriptors, nurses and residents chose 4±2 descriptors, and clinicians in other roles chose 6±2 descriptors.
- 8% of respondents added the following novel terms (with corresponding illness severity scores) not otherwise identified in pilot interviews: looks well (1), resource limited (2), just extubated (2), slightly combative when not sedated (3), geriatric patient (4, 4, 6), long-term rehab (5), wound infection (6), significant lung issues (6), poorly oxygenating (6), not ready for transfer (7), chronic critical illness (8), tenuous (9), critically ill (7 ,7, 9)
- There is a trend of terms that describe a patient’s illness severity (figure 1).
- A priori analysis demonstrated that:
 - “Not sick” was chosen exclusively, but infrequency, for least sick patients.
 - “Shock like” and “actively trying to die” were chosen exclusively, but infrequently, in the most sick patients.
 - No term was chosen exclusively within the middle sick illness severity group.
 - “Ready for Transfer” had the strongest strength within group statistic and lowest average illness severity.
 - Three terms – “most interventions,” “multiple organ failure,” and “the sickest patient” – were consistent with an illness severity above 8 and had strength within group statistics of > 20%.
- If the most sick group included illness severity scores of 7-10, two additional terms, both including the work “profound” would be notable for describing the most sick patients.

- Confidence intervals for terms widely overlap illness severity scores (data not shown).

#surveys/group	11	13	16	20	23	15	25	25	13	8	169				
Illness Severity	1	2	3	4	5	6	7	8	9	10	Term Count	Mean Severity	A Priori Strength within Group	Final Strength within Group	
Ready for transfer	91%	38%		5%				4%			17	1.9	0.34	0.34	
Not Sick	9%	8%	19%								5	2.4	0.13	0.13	
Just Monitoring	9%	15%	6%	10%		7%					7	3.1	0.06	0.06	
Getting Better	45%	85%	31%	30%	17%	13%	4%	8%			36	3.4	0.32	0.32	
Stable	73%	38%	75%	70%	30%	20%	8%	8%			53	3.6	0.27	0.27	
Improving	55%	77%	69%	30%	39%	7%	12%	16%	8%		51	3.8	0.34	0.34	
Going in the right direction	55%	69%	56%	50%	48%	20%	16%	12%	15%		57	4.1	-0.37	0.15	
Normotensive	36%	15%	44%	30%	22%	27%	28%	8%		13%	38	4.7	-0.13	-0.13	
Stabilized	27%	31%	25%	40%	30%	53%	28%	8%	8%		44	4.8	0.02	0.02	
Medium Sick	18%	8%		35%	30%	20%	12%	8%			25	4.9	0.12	0.12	
Ventilated	9%	15%	44%	55%	61%	53%	72%	60%	69%	38%	88	6.2	-0.22	-0.22	
Infected	9%			20%	17%	27%	36%	24%	38%	13%	34	6.7	-0.03	-0.03	
Critically Stable			6%	5%	43%	40%	64%	48%	15%	13%	49	6.7	0.05	0.05	
Getting Worse				5%		7%	12%	20%	23%	13%	14	7.7	-0.14	0.14	
Profound lung injury		8%		5%		7%	4%	20%	38%	38%	17	7.9	-0.27	0.22	
3+ sick					4%	7%	4%	12%	23%	13%	10	7.9	-0.12	0.12	
Profoundly ill				10%		7%	28%	40%	85%	25%	33	7.9	-0.38	0.38	
Most Interventions				5%		7%	16%	24%	23%	50%	19	8.1	0.21	0.21	
Multiple Organ Failure				5%	4%	7%	4%	24%	23%	50%	17	8.1	0.23	0.23	
In Shock							4%	4%	31%		6	8.5	0.10	0.10	
The sickest patient						7%	4%	28%	38%	88%	21	8.8	0.39	0.39	
Shock Like								4%	23%	13%	5	9.0	0.11	0.11	
Actively trying to die								4%	8%	25%	4	9.3	0.09	0.09	

Figure 1. Contour plot showing the frequency of terms chosen according to patient illness severity as indicated by clinicians. Percentage = the count of a term divided by the number of surveys/illness severity score. Also shown are the number of times a term was chosen overall (**term count**), the mean illness severity for which a term was chosen (**mean severity**), and the difference between the frequency a term was chosen within the A Priori Group (least sick = 1-3, middle sick – 4-7, most sick = 8-10) and Final Group (where most sick = 7-10) vs. outside the group (**strength within group**). This statistic helps differentiate terms that are used often (i.e. by many clinicians) and are consistent with a particular illness severity group. Of note, illness severity in reality is not *discrete*: it represents a continuum of patient condition. Here, we try to demonstrate that with a gradient of color whereas in the analysis we uses three discrete groups.

- Clinician terminology for patient illness trend in terminology
- Terms that indicate differentiate between Burn ICU patients differentiation in middle sick patients
- Changing illness severity the strength within group not change consistently
- Limitations:
 - This was not a controlled study may have and instead of using
 - Patient illness severity yet we have a

- Lack of a shared terminology to patient condition increases risks of dyssynchrony.
- Improving teamwork condition by standardizing terminology middle and patient care

Acknowledgements

- We would like to thank instrumental assistance
- Funding: This project was funded by the US Army Medical Research and Materiel Command Telemedicine and Ambulatory Care (TATRC) (W81XWH-04-2-0001)

1. Burke CS, et al. Qual a
2. Hall P. Journal of Interp
3. Lingard L. Adv Health S
4. The Joint Commissions
5. Varpio L, et al. Acad Me

- Multidisciplinary teams in the burn intensive care unit (BICU) are large, comprised of clinicians from multiple different clinical backgrounds¹
- Teamwork and communication are essential in the care of the critically ill burn [patient](#)¹
- Substantial evidence supports using checklist tools to support clinical decision making for the critically ill²⁻⁸
- In order for tools to be useful, they must be *Ecologically Valid*- or represent the work domain they intend to support^{9,10}

Theoretical Underpinnings

- The Phases of Illness Paradigm (POIP) is a theory that defines patients by severity of illness (SOI) and [aligns treatment to patient condition](#)
- [Decision support tools based on the POIP enable shared mental models which enhance communication, efficiency, and care delivery.](#)

Figure 1: The Phases of Illness, Severity of Illness Scale



Objectives

- We sought to create ecologically valid clinical decision support tools for the BICU based upon the POIP model

Methods

- This was part of a prospective mixed methods study in three academic regional referral Burn ICUs
- [After the first 6 months of tool use, we assessed user satisfaction](#) using a 10 question 5-point Likert-like scale
- [Although intended to enhance team communication, the burden of tool completion was placed on nurses.](#)
- For statistical analysis we reduced the scale from 5 points to 3 points: agree, neutral, and disagree

- We surveyed 48 end-users ([response rate?](#)) (Physicians: 3; Nurses: 30; Others: 15)
- Two tools were created, see figure 2
- Mean Overall satisfaction scores were 3.23 (SD 0.98)
 - Fewer participants rated very low/low (20%) compared to those rating very high/high satisfaction (44%) and those that were neutral (35%)
 - [Physicians were more satisfied than nurses: 100% Satisfaction \(n =3\) vs. 36% dissatisfaction \(n=30\)](#)

• [See graph to right for combined scores \(n=48\)](#)



Figure 2: Two tools

Burn Illness Severity Assessment tool

- Clinicians assess patient along 6 continuous scales
- Clinicians determine SOI and make a mark along the rainbow patterned scale
- Treatments are checked off along each of the 12 scales
 - Clinicians assess the appropriateness of treatments for the SOI category

The TeamView board is located outside each patient's room

- Clinicians update the daily SOI along the top rainbow patterned area
- Daily events: tube and line changes are tracked below
- There is space for messages, and antibiotic start and stop dates are tracked
- On rounds, the care team updates the overall daily goals & fluid goals
- Priorities of care are determined and noted on the board

- Creating tool requires frequent communication in order to produce a valid and useful tool
- [Overall](#), clinicians' time was not interfered with by the tools greatly, but tool completion during multiple rounds was
- [Using this data to improve completion of tool during multiple rounds](#)
- Daily use of tool was not a routine since it was not a required part of the workflow


- The POIP checkerboard tool and communication workflow for rounds
- After 6 months of use, the tool was comfortable with a lower than expected burden
- We used this tool for both patient and team communication
- We would like to have further help with the tool for guidance
- Thanks to the clinicians for their study

- Herndon, ed., 2007, Total burn care
- Kuwamoto 2010, F, Care Med
- Pronovost 2006, A
- Haynes 2009, A su
- DuBose 2008, Mea
- Weiss 2011, Prom Med
- Narasimhan 2007, Crit Care
- Dodek 2003, Expli Med
- Hales 2008, Dev o
- Winters 2009, Cli

- Critical care of burn patients requires a multi-professional team of clinicians
- They must effectively collaborate in order to ensure optimal outcomes
- Effective teamwork necessitates team members share common goals
- Establishing goals requires a shared mental model of a patient's current condition, anticipated future state, and treatments to bridge the two
- Better understanding of how clinicians understand patient condition, and ultimately treatment priorities, could improve teamwork, communication, and patient outcomes

Objectives

- We sought understand how a team of clinicians perceive patient condition, and measure the impact of a burn specific illness severity assessment tool on team understanding



Figures 1 &2: Pre-Baseline Card Sort Survey Examples


- Two examples from a concurrent study demonstrating the differences amongst clinicians in perceiving patient SOI
- Pre-baseline and baseline CUS data were similar to these examples

- This was part of an IRB approved, prospective, mixed methods study to understand clinician perspectives about patient condition and treatment priorities
- This understanding was used to create tools that could improve clinician decision making, teamwork, and communication according to the Phases of Illness Paradigm
- Clinician perspectives were elicited through survey and card sort methodology
- Tools were created through using data obtained from card sort (see figures 1 & 2) and refined by focus groups, the Delphi process for consensus building, and pilot testing

- To identify the impact of these tools on the mental models clinicians form about patient condition, we asked clinicians to identify their patient's *current* and *anticipated* condition on the following day by marking a continuous scale from “Most sick/Could die today” to “Least sick/Could transfer from the ICU today”
- Continuous scales divided into ten equal parts and the location of a clinician’s mark upon them was scored from 10 (most sick) to 1 (least sick)
- We then evaluated the variance of the differences between clinician perspective before beginning the project (pre-baseline), before implementing tools (baseline), and six months after implementing tools testing

Results

- The average variance reflecting clinician perception of a patient’s current and anticipated condition was as follows:
 - pre-baseline, ± 1.73 and ± 1.37 see graph below
 - baseline, ± 1.38 and ± 1.33
 - six months, ± 0.7 and ± 0.76
- The decreased variance for both current and anticipated condition was statistically significant at each time point (ANOVA, $P = 0.01$)



- The process of clinician perception of patient condition and treatment priorities is a key component of the BICU (pre)
- Introducing a developed through further improvement about patient
- Improving the current and future teamwork by of the patient
- Patient care of teamwork and regarding patient treatment priorities

Pamplin JC, M illness paradigm outcomes. Cri

Thank you to: Nico at the USAISR for guidance; Agnes B Christopher Tran, M Adrian Botello, CS at Memorial Herma dedication to this p

This project is supp Research and Mat Technology Resea

This study was cor approved by Brook board and in accor

- Patient care in the burn intensive care unit (BICU) is complex and involves large multidisciplinary teams
- If team members are not working toward common goals, communication, efficiency, synchronization, and ultimately patient safety may be compromised
- The Phases of Illness Paradigm (POIP) is a theoretical model that defines patients by severity of illness, aligning patient condition to treatments
- Understanding how clinicians perceive patient condition and associated care goals is necessary to develop ecologically valid tools to support their daily work

Objective

- To understand how clinicians perceive patient condition and associated care goals and the degree to which individual team members share the same mental model of the patient

Methods

- A “Condition Understanding Survey” was created to assess clinician’s perception of the Patient’s Severity of Illness (SOI) and associated care goals
- Surveys were administered during normal patient care duties and clinicians were asked to:
 - rate patient SOI on a 10 point scale from “most sick” (10) to “least sick” (1)
 - prioritize goals of care
- A goal was defined as “a near-term desirable outcome for the patient. There may be more than one goal . These should be patient focused.”
- We collected clinician perspectives on at least two patients in each of the following groups : least sick: 1-3; middle sick: 4-7; most sick: 8-10
- We evaluated goal agreement between clinicians for patients with whom we obtained responses from at least 3 clinicians on the basis of the same goal appearing in any of their top 4 responses
- The authors coded responses to ensure they matched the definition of goal and for thematic analysis
- Adjudication occurred by majority vote

- Overall, there were 169 responses from 60 respondents goals. Only 95 responses (57%) met the definition of goal (example below, table 1)
 - The most common goals were are shown in figure 1
 - Only six patients had at least three clinicians that responded. Comparisons between clinician responses for this six patients showed:
 - 28% +/- 17% agreement about top care priorities
 - Clinicians perceived SOI more consistently in patients who were “most sick” than in patients who were “middle sick” (figure 2)

Table 1. Example of raw and coded data.

Original Category	Utterance	Coded Category	Coded Utterance
Goal	Maintain oxygenation and ventilation	Goal	Adequate oxygenation and/or ventilation
Goal	Skin grafting	Objective	Surgical operation (implicit goal = achieve wound healing)




Fig 1. Coded goals in middle and most sick patients

- This data suggests different beliefs and priorities
- Of the top care 4 most were not shared
- Lack of common patient and decision coordination, and
- Interestingly, the each SOI group) number of themes
- Identifying common protocols, or path patient types, wh in this project, or disease process, improving team p

- Clinicians in the E mental model of p
- Creating common should decrease performance, pat outcomes
- This might be acc to address frequ complexity an inc differences

Acknowledgements

- Thank you to Nicole C with this project
- This project is support Research and Materie Technology Research

- Pamplin J, et al. A daily hand hygiene culture.
- Pamplin JC, et al. Pha communication, better
- Schroeder, R. E. et al. professionals through e Health Administration E

This study was conducted by the Brook Army Medic accordance with the appr

Journal of Burn Care & Research

Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs --Manuscript Draft--

Manuscript Number:	JBCR-D-15-00153R2
Full Title:	Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs
Article Type:	Original Article
Keywords:	Teaching Rounds; Intensive Care; Cognitive Work; Task Performance and Analysis
Corresponding Author:	Jeremy C. Pamplin, MD US Army Institute of Surgical Research JBSA Ft. Sam Houston, TX UNITED STATES
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	US Army Institute of Surgical Research
Corresponding Author's Secondary Institution:	
First Author:	Ian Christopher McInnis, DO
First Author Secondary Information:	
Order of Authors:	Ian Christopher McInnis, DO
	Sarah Murray, MSN
	Maria Serio-Melvin, MSN
	Jay K. Aden, PhD
	Elizabeth Mann-Salinas, PhD
	Kevin K. Chung, MD
	Todd Huzar, MD
	Steven Wolf, MD
	Christopher Nemeth, PhD
	Jeremy C. Pamplin, MD
Order of Authors Secondary Information:	
Abstract:	Introduction: Multidisciplinary rounds (MDR) in the Burn Intensive Care Unit serve as an efficient means for clinicians to assess patient status and establish patient care priorities. Both tasks require significant cognitive work, the magnitude of which is relevant because increased cognitive work of task completion has been associated with increased error rates. We sought to quantify this workload during MDR using the National Aeronautics and Space Administration Task Load Index (NASA-TLX). Methods: Research staff at 3 academic regional referral burn centers administered the NASA-TLX to clinicians during MDR. Clinicians assessed their workload associated with 1) "Identify[ing] if the patient is better, same, or worse than yesterday" and 2) "Identify[ing] the most important objectives of care for the patient today." Data were collected on clinician type, years of experience, and hours of direct patient care. Results: Surveys were administered to 116 total clinicians, 41 physicians, 25 nurses, 13 medical students and 37 clinicians in other roles. Clinicians with less experience

reported more cognitive work when completing both tasks ($p<.005$). Clinicians in the "others" group (respiratory therapists, dieticians, pharmacists, etc.) reported less cognitive work than all other groups for both tasks ($p<0.05$).

Conclusions:

The NASA-TLX was an effective tool for collecting perceptions of cognitive workload associated with MDR. Perceived cognitive work varied by clinician type and experience level when completing 2 key tasks. Less experience was associated with increased perceived work, potentially increasing mental error rates and increasing risk to patients. Creating tools or work processes to reduce cognitive work may improve clinician performance.



REPLY TO
ATTENTION OF

DEPARTMENT OF THE ARMY
U.S. ARMY INSTITUTE OF SURGICAL RESEARCH
3698 CHAMBERS PASS
JBSA FORT SAM HOUSTON, TEXAS 78234-7767

10 July 15

Dear Dr. Gamelli,

Thank you for accepting and allowing us to revise again our manuscript JBCR-D-15-00153, entitled "Comparing the workload perceptions of determining patient condition and priorities of care among burn providers in three burn ICUs." We are greatly appreciative of the thoughtful commentary by your reviewers which again has improved the quality of our manuscript. Individual responses to your reviewers concerns are below.

I hope these revisions adequately address your concerns. Please do not hesitate to contact me directly if there is anything I can do to help facilitate review and/or revision.

Kind regards,

JEREMY PAMPLIN, MD, FCCM, FACP
LTC, MC, USA
Medical Director, Burn ICU
Chief, Clinical Trials in Burns and Trauma
U.S. Army Institute of Surgical Research
O – 210-916-6603
C – 210-527-7581
Jeremy.c.pamplin.mil@mail.mil

Reviewer #2:

I detected one remaining typo. In the revisions on page 12 there is a statement "....least about of total cognitive work..."

- *Corrected. Thank you.*

I believe the revisions have improved the paper. There will always be criticism of data from a "convenience sample of volunteer subjects" but I believe the alternative of mandating completion of the data forms from every participant in rounds would result in some forms being filled out superficially just "to get them done". The latter option would not necessarily produce more reliable data.

- *Thank you.*

Reviewer #3:

The authors have substantially improved their manuscript. As indicated in my earlier review, I cannot too highly praise this paper. It will have significant impact on the human factors describing burn care and will likely trigger new directions in the organization and delivery of care in burn centers. The suggestions below are meant to improve an already excellent paper and to answer a few questions that continue to intrigue me.

- Thank you.

First, I would like to emphasize that the ICU experience is only a small portion of the total treatment of "burn disease". It appears to me that the authors view burn critical care as an isolated and nearly independent part of burn care, even to the point of suggesting the ICU staff should be formally separated from the post-injury courses of burned patients. Whether increased fragmentation of care proves to be beneficial to patients remains to be seen, but the authors' methodology will likely be the best way to measure these effects.

- We agree that the ICU experience is only a part of a burn patients' overall burn care. We did not intend to imply that the ICU experience nor the clinical teams should be separated. We certainly did not study this nor did we present data about this. In all of the centers involved in this project, clinicians representing all locations of a patient's care (ICU, Ward, Clinic) and all professions contributing to it (intensivist, surgeon, RT, rehabilitation (OT & PT), social work services, nursing, nutrition, etc.) participate in patient care. Unfortunately, our convenience sample was not able to obtain data from all of these clinicians types.

Admittedly, the study sample is quite small, especially in relation to all of the pertinent variables involved. Much more data will be needed to potentially eliminate some of the non-normal distributions. The "other" group is particularly problematic, in that this group is overwhelmingly composed of the more experienced clinicians, particularly for the clinicians who may be "too old" to work on a burn unit. The fact that only one OT represents the Rehab disciplines is eye catching. As one of the pillars of the ABA, the lack of representatives of Rehab group is of concern - does the sample represents the distribution of the "other" in the individual units?

- No, our study distribution of "others" does not represent the actual distribution of "others" caring for patients in the burn center. Our primary target population was nurses and physicians in this protocol since these groups represented the largest proportion of clinicians caring for burn patients on any given day. Also, these groups have the greatest distribution of experience for us to examine.

Table 1 lists n = 11 burn surgeons, but the subgroups add to 31.

- Great catch! Fixed.

Further, are none of the burn surgeons surgical intensivists?

- We did not collect this data specifically in this project, but the majority of the burn surgeons in this project were also surgical intensivists. We have added the term “medical” before intensivist throughout the manuscript for clarification.

In the same table, is "average experience" the duration of medical training or the time spent in burn centers.

- This was the time since graduation from licensing school or year in school for students. We have added this information to the manuscript.

If the figure represents time since medical school graduation, a separate column indicate full time burn center assignment should be provided.

- Unfortunately, we did not collect this information.

Particularly in the military burn center, many of the assignments are short term (less than 5 years) rotators.

- Agree.

I feel strongly that the military burn center should be identified by site - is it site 2 ??

- We are uncertain as to how we should respond to this request. We do not feel that including this information will add additional insight to the manuscript. The military center was not site 2.

In particular, if it is site 2, as is my guess, I think the authors have made a phenomenal observation about the remarkable effectiveness of the military philosophy of education and leadership that would be beneficial to resolution of many of the shortcomings suggested in this study. The authors might consult Dr Gamelli to see if this would be acceptable.

- No response.

This reviewer would be a member of the "> 10 yrs" group. What's wrong with us? Should we leave clinical care or choose a new specialty? Hopefully, in the fullness of this research, you will be able to answer such questions. I should point out that many burn units have only a few daytime physician extenders and no residents. The attendings are on both attending call and resident call 24/7 and these duties certainly constitute workload. Could you better describe this group of clinicians?

- We are unclear as to which group of clinicians this reviewer would like us to describe (attending in general, clinicians with > 10 yrs experience, or attending that perform both “attending” and

“resident” duties during a 24/7 call) and what constitutes “better?” We did not collect additional demographic information for any of these groups.

Please better describe your statistical considerations and methods. You indicate that you used Wilcoxon and Steel-Dwass techniques. What's the difference? What does it mean when one method is significant and the other is not? What does it mean when both are significant?

- We have edited the manuscript to reflect that we used the Wilcoxon Test *with* the Steel-Dwass correction and provided a reference. The Steel-Dwass correction provides a means to make non-parametric data appear more “normal” in distribution before performing statistical comparisons. When you use the Steel Dwass adjustment, you increase the p-value due to increases in the false positive rate. Therefore, Steel-Dwass is conservatively decreasing the chances of a type-1 error. The Steel-Dwass is less likely to be significant over just doing a Wilcoxon Test alone.

Finally, I would suggest further emphasizing the utility of the NASA-TLX instrument as an effective instrument for assessing not just human factors but also for evaluation team effectiveness and maybe even adequacy of staffing in the burn center. For example, was there less frustration at site 2 because it had a lower average census?

- These are thought provoking observations. Unfortunately, we did not collect data to address these considerations in this project, but they represent great ideas for future study!

There are a number of grammatical errors persisting in the manuscript, especially in the latter half of the Discussion. I would like to emphasize that the authors are excellent writers - the manuscript was a pleasure to read. Reference 11 lacks the name of its journal in the citation. As stated above there appears to be a numerical inconsistency in Table 1.

- We have, hopefully, corrected all of the grammatical errors.
- We have added the journal for the citation.
- Table 1 has been corrected as above.

My curiosity has got the best of me. What did Nicole Caldwell do for this paper? Given the potential magnitude of this work, she certainly deserves her recognition.

- She facilitated our regulatory approvals and administrative support of this project. This has been added to the manuscript.

Title: Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs.

Ian McInnis, DO, MS1
Sarah Murray, MSN2
Maria Serio-Melvin, MSN2
Jay K. Aden, PhD2
Elizabeth Mann-Salinas, PhD2
Kevin K. Chung, MD, FCCM, FACP 2
Todd Huzar, MD3
Steven Wolf, MD, FACS4
Christopher Nemeth, PhD5
Jeremy C. Pamplin, MD, FCCM, FACP2

1. William Beaumont Army Medical Center, Fort Bliss, TX
2. United States Army Institute of Surgical Research, Fort Sam Houston, TX;
3. Memorial Hermann Hospital Texas Medical Center, Houston, TX;
4. University of Texas – Southwestern Medical Center, Dallas, TX;
5. Applied Research Associates, Inc., Fairborn, OH

Address for Correspondence and Reprints:

Jeremy C. Pamplin
3698 Chambers Pass, JBSA Ft. Sam Houston, TX 78234
Telephone: 210-916-3301
Fax: 210-271-0830
Email: jeremy.c.pamplin.mil@mail.mil

External funding: This project is supported by a grant from the US Army Medical Research and Materiel Command Telemedicine and Advanced Technology Research Center (TATRC) (W81XWH-13-2-0011).

Conflicts of interest: The authors declare that there are no conflicts of interest.

Disclosure: This study was conducted under a protocol reviewed and approved by the Brooke Army Medical Center Institutional Review Board and in accordance with the approved protocol. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Title: Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs.

Abstract

Introduction:

Multidisciplinary rounds (MDR) in the Burn Intensive Care Unit serve as an efficient means for clinicians to assess patient status and establish patient care priorities. Both tasks require significant cognitive work, the magnitude of which is relevant because increased cognitive work of task completion has been associated with increased error rates. We sought to quantify this workload during MDR using the National Aeronautics and Space Administration Task Load Index ~~survey~~ (NASA-TLX).

Methods:

Research staff at 3 academic regional referral burn centers administered the NASA-TLX to clinicians during MDR. Clinicians assessed their workload associated with 1) “Identify[ing] if the patient is better, same, or worse than yesterday” and 2) “Identify[ing] the most important objectives of care for the patient today.” Data were collected on clinician type, years of experience, and hours of direct patient care.

Results:

Surveys were administered to 116 total clinicians, 41 physicians, 25 nurses, 13 medical students and 37 clinicians in other roles. Clinicians with less experience reported more ~~cognitive~~ work when completing both tasks ($p<.005$). Clinicians in the “others” group (respiratory therapists, dieticians, pharmacists, etc.) reported less ~~cognitive~~ work than all other groups for both tasks ($p<0.05$).

Conclusions:

The NASA-TLX was an effective tool for collecting ~~perceptions of cognitive workload data~~ associated with MDR. Perceived ~~cognitive workload~~ varied by clinician type and experience level when completing 2 key tasks. Less experience was associated with increased perceived work, potentially increasing medical error rates and increasing risk to patients. Creating tools or work processes to reduce cognitive work~~load~~ may improve clinician performance.

Key Words

1. Teaching Rounds
2. Intensive Care
3. ~~Cognitive Workload~~
4. Task Performance and Analysis

Introduction

The burn intensive care unit (ICU) presents a challenging work environment to the healthcare team. Effective care requires the coordinated effort of healthcare professionals from different backgrounds, each member contributing unique perspective toward a common goal. These teams are a necessity given the complexity of managing the medical, administrative, and social aspects of patient care. Multidisciplinary rounds (MDR) is an event that facilitates communication of relevant information among team members and is associated with better outcomes in the ICU.^{1,2,3}

While MDR represent a powerful tool and a logical means for efficient dissemination of information, there is variability in the quality and effectiveness of such rounds. MDR must encompass key aspects of care, to include problems, plans, and goals. The multidisciplinary team must be effectively *managed* to ensure that all important aspects of patient care are addressed. Caution must also be taken to ensure that information is discussed clearly and in such a way that it is accessible to all ~~of the~~ members of the team who need it; ~~thus~~, helping to avoid the need for inefficient clarifications with individual team members after MDR have ended.⁴ Input from all team members must be evaluated and integrated by the team leader, typically a physician, in such a way as to create a cohesive, prioritized care plan.²

A series of decisions are made during MDR, some critical, some routine. Likewise the ~~mental~~ ~~cognitive~~ work that goes into making these decisions is variable. In the same way that physical work is defined in terms of distance and force, ~~mental cognitive~~ work can be imagined as a function of a task and the perceived effort required ~~to completeing~~ it. Different people ~~would~~ have different perceived workloads when completing the same task according to their familiarity with the task and their overall experience with the task or similar tasks.⁵ Calculating workload by this definition requires quantification of perceived mental “force.” Hart and Staveland developed the National Aeronautics and Space Administration Task Load Index (NASA-TLX), a survey composed of six subscales, designed to

assess perceived workload of NASA pilots. These subscales include mental demand, physical demand, temporal demand, frustration, effort, and performance as defined in the NASA-TLX Manuscript.⁵ The NASA-TLX has been used broadly in aviation, the automotive industry, and more recently in medicine.⁶⁻¹¹

Lower perceived workload, as measured by the NASA-TLX, has been linked to better performance in medicine.^{10,11} Proper identification of a patient's illness severity may also be associated with improved clinician performance. Starmer and colleagues demonstrated a significant reduction in preventable adverse events after implementation of a tool that increased communication of patient illness severity during change-of-shift handoffs.¹²

Multidisciplinary rounds present a cognitive challenge: to integrate data, make diagnoses, prioritize care tasks, provide education, and to form shared mental models.^{3,4} The magnitude of this challenge, however, has not been well defined, however. Also unclear are the differences in cognitive workload, if any, experienced by the different types of clinicians who participate in MDR. The current study sought to define the perceived workload associated with the completion of two important tasks during MDR: generating a personal assessment of overall patient condition and identifying the priorities of for the day's care plan.

Methods

We conducted this institutional review board approved survey study in three American Burn Association verified regional referral Burn ICUs. The NASA-TLX surveys were administered by research staff to clinicians at each site who participated in MDR during daily clinical practice according to the process outlined below.

Admission rates to these Burn ICUs varies from 270 to 300 patients per year with an average daily census between 2 (site 2) and 5 (site 1 and 3). At each site, MDR are led by an attending physician who was either a burn surgeon credentialed in critical care (sites 1, 2, and 3), or a medical intensivist with extensive burn experience (site 1). Nurse to patient ratios vary among sites from 1-to-1.5 nurses per patient at site 1, to 1 nurse per patient at sites 2 and 3. Attendance at MDR varies, but typically consists of the bedside nurse, rehabilitation specialists, registered dietitians, clinical pharmacists, respiratory therapists, burn fellows, residents, and students.

Multi-Disciplinary Rounds Conduct

MDR typically consisted of a resident presenting the patient by systems followed by other team members providing additional pertinent details and recommendations as they felt appropriate. The attending physician would direct the information flow and sequence of rounds as he felt necessary and would elicit from the team and/or state a care plan for the patient. Additionally, site 1 used a daily communication tool completed by the unit's "charge nurse" (e.g. nursing shift leader). This communication tool consists of a checklist that confirms use/absence of certain quality metrics (i.e. deep venous thrombosis prophylaxis, gastric ulcer prophylaxis, head-of-bed elevation, etc.) and a list of daily tasks that the charge nurse identified during rounds. At site 1, the charge nurse reviews the checklist and reads the identified tasks as the last activity of a patient's rounds prior to the MDR team moving on to the next patient.

Survey Administration:

Research staff administered the NASA-TLX as ~~a survey using~~ either a paper (sites 1 & 3) or electronic (site 2) ~~means~~ survey to a convenience sample of clinicians present on MDR for at least five discrete rounds events. A “discrete rounds event” was MDR rounds for a single patient. Thus, the NASA-TLX could be administered once per day for five days or five times in a single day for five different patients. Surveys were administered immediately after MDR was complete for a single ICU patient. Each site adjusted the survey collection period to meet the needs of their site and according to patient volume. Surveys were administered during different points in the MDR to account for physical and mental fatigue that may impact workload perception. For example, surveys were administered early in rounds (after the first patient), middle (after the middle patient) or late in rounds (after the last patient).

Prior to administration of the NASA-TLX survey, research staff reviewed a consent document with potential clinician subjects who participated in MDR. Clinicians were not required to participate, but their participation implied consent. Subjects were allowed to quit the study at any time by not completing the survey. Incomplete surveys were not included in the data analysis. Participation was anonymous and no personally identifying information or master list of participants was collected. We collected participant demographics including clinician type, years of experience, and time spent in direct care of the patient.

After the consent was reviewed and a copy provided to subjects, the research staff read directions aloud and then provided the NASA-TLX survey to participants. Time was provided for questions and clarification by the research staff. Clinicians assessed their workload associated with two tasks identified by the following questions: 1) "Identify if the patient is better, same or worse than yesterday" and 2) "Identify the most important objectives of care for the patient today."

Statistical Analysis

For the purpose of this study, we analyzed results according to the following groups: “physician” represents all attending physicians (burn surgeons or **medical** intensivists) who led or participated in MDR, as well as all physicians in training including those physicians who were part of a certified residency training program or fellowship; “nurse” represented all nurse types including licensed vocational nurses, registered nurses, and clinical nurse specialists; “student” represented medical students; and “other” represented clinicians not in any other category including respiratory therapists, dietitians, occupational **or physical** therapists, pharmacists, and psychiatry support personnel.

NASA-TLX Scores were calculated as described by Hart⁵: weighted cognitive load was calculated by first determining individual subscale ratings for mental demand, physical, temporal, performance, effort, and frustration, reported on a scale of 0-100 by asking subjects to place a mark on a line divided into 50 equal parts. A researcher then converted this analog representation into a value from 0-100. Individual subscales ratings were given a weighting factor which represented the perceived importance of a given subscale relative to the other subscales. This weight equaled the number of times a subject selected the subscale over another subscale while completing the “sources of workload” portion of the NASA-TLX survey. During this portion of the survey, subjects were asked to determine which subscales were the more important contributors to workload according to a series of 15 pair-wise comparisons. Thus, a weight could range from 0-5 (0 meaning that a subject never chose the subscale compared to other subscales and 5 meaning the subject always chose a subscale when comparing it to other subscales). Individual raw subscale data were multiplied by this weighting factor and were then summed. This sum was then divided by 15 to yield the weighted (total) **cognitive workload**.

The NASA-TLX survey scores were summarized using medians and interquartile ranges for each of the demographic groups. Due to the small sample size and lack of normality, the scores were analyzed using the nonparametric Wilcoxon's Test ~~alone and~~ with the Steel-Dwass ~~method correction~~ for multiple test correction.¹³ Significance was established when the p-value was less than 0.05. All

analysis was performed using Statistical Analysis System (SAS) v9.2, SAS Institute Inc., Cary, USA or JMP v10.0, SAS Institute Inc., Cary, USA.

Results

Surveys were administered to a total of 119 clinicians across all sites; however three surveys were incomplete and were excluded from the analysis for a final $n = 116$. The median weighted cognitive load for task 1, identifying if a patient was better, same, or worse than the previous day (i.e. identifying the patient's "severity of illness" or SOI), was 40 (IQR 13-67). The median weighted cognitive load rating for task 2, identifying the most important objectives of care for the patient today (i.e. identifying the patient's "priorities of care" or POC) was 43 (IQR 18-68). Mental demand, temporal demand, performance, and effort all contributed more to the overall cognitive load than did physical demand or frustration subscales (table 1, figures 1-2). Perceived workload varied significantly according to clinician experience and professional background (figures 3-6), but not according to clinician location or time spent with the patient (data not shown). Interestingly, the frustration and physical demand subscales were particularly *non-normal* in their distribution across our sample. ~~40%~~Forty percent of respondents reported no frustration and 63% of respondents reported no physical demand for identifying SOI. Numbers were similar for respondents' perception of frustration and physical demand for identifying POCs: 38% and 62% reported zero frustration or physical demand respectively.

Comparing perceived cognitive workload between clinician types revealed several significant differences ($p < 0.05$ for all comparisons, figures 3-4). Students, nurses, and physicians experienced a higher total cognitive workload than did "others" when identifying both SOI and POC. Students experienced more mental demand than "others" when identifying SOI and more than nurses and "others" when identifying POC. Nurses and physicians experienced more temporal demand than the "others" group when identifying SOI, while students, nurses, and physicians experienced more temporal

demand than “others” when identifying POC. Students also reported increased effort associated with identifying SOI as compared to all other groups, and more than physicians and “others” when identifying POC. Students perceived their performance to be worse than “others” when identifying SOI and nurses perceived it to be worse than “others” when identifying POC. Nurses and physicians perceived more frustration than ~~did~~ students and “others” when identifying SOI and more than others when identifying POC. Lastly, there were no significant differences between nurses and physicians for either task (figures 3 and 4).

It should be noted that the three largest groups, physicians (n=41), “other” providers (n=37), and nurses (n=25) had an ~~somewhat~~ unequal distribution of experience levels within their respective professional groups. The physician group included a large proportion (19 surveys, 46%) with <5 years of experience. The “others” group was largely composed of surveys of individuals with 5-10 years of experience (22 surveys, 59%). The nurses were similar to the physicians, in that a large proportion of those surveyed had <5 years of experience (14 surveys, 56%) (table 2).

Significant differences were also discovered when study participants were compared based upon their years of experience ~~since graduation from licensing school (or year in school for students)~~ (P < 0.05 for all comparisons, figures 5-6). Participants with 5-10 years of experience reported the least ~~about~~ amount of total cognitive work for both tasks compared to clinicians with <5 or >10 years of experience. Clinicians with > 10 years of experience perceived more total cognitive work for both tasks compared to clinicians with 5-10 years of experience, but less than clinicians with < 5 years of experience. Finally, clinicians with < 5 years of experience perceived the most total cognitive work, which was significantly more than clinicians with > 4 years of experience.

When examining subscales that contribute to total cognitive work, the following observations were made: Clinicians with ~~more~~ < 5 years of experience perceived more mental demand than those with >4 years of experience when identifying SOI and more than clinicians with 5-10 years of experience

when identifying POC. Clinicians with < 5 years of experience perceived more temporal demand and felt their performance was worse for both tasks when compared to clinicians with 5-10 years of experience. Both tasks were perceived as requiring more effort for clinicians with < 5 years of experience compared to those with > 4 years of experience. Clinicians with 5-10 years of experience perceived significantly less frustration while identifying SOI than did those with < 5 and those with > 10 years of experience, but only less than those with > 10 years of experience when identifying POC. Finally, there was no apparent impact of experience on perceived physical demand. The largest differences in workload perception were consistently between those with < 5 years and those with 5-10 years of experience for all domains except frustration (see figures 5 and 6).

There were no significant differences in perceived total workload when comparing study sites or hours of direct patient care with respect to identifying either SOI or POC. However, there were differences between sites for some subscale domains. Site 3 clinicians perceived significantly more temporal demand than site 1 when identifying SOI (median 150 [IQR 90-210] vs. 60 [40-80], $p < 0.05$). Site 1 clinicians perceived significantly worse performance **when** identifying SOI than did clinicians at site 3 (median 140 [IQR 90-190] vs. 80 [IQR 40-120], $p < 0.05$). Clinicians at site 2 perceived significantly less frustration than clinicians at site 1 or site 3 for both tasks. Site 2 perceived no frustration when identifying SOI or POC, whereas clinicians at site 1 and site 3 perceived frustration as a relatively low contributor to **cognitive workload** when identifying SOI (site 1 & 2 frustration subscale score 20-30 [IQR 75% 0-120]) and when identifying POC (site 1 & 2 frustration subscale score 30 [IQR 75% 0-120]).

Discussions

This manuscript is the first to describe the cognitive work performed by clinicians during MDR in the Burn ICU. The NASA-TLX effectively revealed workload perception differences and similarities in cognitive work associated with completing two critical tasks performed during MDR: identifying a

patient's condition (severity of illness, SOI) and prioritizing associated treatments (priority of care, POC).

Significant findings include:

1. Mental demand, temporal demand, performance, and effort were the primary determinants for the cognitive work performed for the identified tasks on MDR with mental demand being greatest;
2. students, nurses, and physicians all had higher perceived total workload for both SOI and POC than "others";
3. students perceived the most effort on rounds and had the most mental demand when identifying POC and significantly more than "others" when identifying SOI;
4. students, nurses, and physicians experienced significantly more temporal demand when identifying POC than ~~do~~ did "other" healthcare providers while only nurses and physicians perceived this significantly more than others for identifying SOI;
5. clinicians with the least experience ~~have~~ had higher perceived workload when identifying SOI and POC as compared to those with more experience;
6. ~~and~~ some individuals perceived more frustration and physical demand than most others during MDR.

Participation in MDR ~~is~~ was associated with a moderate cognitive workload for ~~the~~ nurses, physicians, students, and other healthcare professionals in attendance when completing two key tasks. Quantification of this workload with the NASA-TLX revealed unequal distribution of workload across the subscales of the task load index. Greater contributors to total load were mental demand, temporal demand, performance, and effort, with mental demand consistently the greatest for both tasks and compared to other subscales. Less substantial contributors were frustration, followed by physical demand. MDR is largely a thought exercise and so it ~~should~~ is not ~~be~~ surprising that participants reported relatively low physical demand. When taken as a whole, the MDR participants reported a low

cognitive workload associated with frustration, particularly at site 2. This low observed frustration level may be a reflection of how well the MDR participants are acclimated to the MDR process, experiencing relatively low levels of stress, annoyance, and irritation when participating in this well-practiced event or that site 2 clinicians work particularly well together, or have some other organizational culture/aspect (e.g. a lower census) that minimizes frustration. Also interesting was that the frustration and physical demand subscales demonstrated a significant right skew effect, driven by a large number of participants reporting little or no load associated with these 2 subscales. This was appreciated at site 2 also. This suggests that certain *individuals* may experience MDR differently than most others. In particular, some individuals perceive remarkably more frustration (figures 1-2).

This study was designed to be descriptive of cognitive workload associated with MDR. We did not seek to find significant ~~among between~~ group differences and yet such differences were found. Why do these differences exist and what impact do they have on patient care? We suspect that clinicians undergo a significant paradigm shift in their approach to medical decision making during the transition from novice to expert. A general stepwise paradigm of skill acquisition has been described previously, and applied to the healthcare setting in prior works.¹⁴⁻¹⁶ ~~A later study~~ One study identified significant macro cognitive differences in the approach of a novice physician in the emergency department as compared to an expert. Junior physicians had difficulty integrating individual aspects of a patient presentation into a cogent picture. The same study showed that less experienced physicians had an overreliance on laboratory data and ~~had~~ difficulty integrating data that was inconsistent with their current diagnosis.¹⁷ Furthermore, differences in professional background, especially those associated with increased specialization with career progression, promote differences in knowledge acquisition and information use that simplifies mental modeling, role based planning, and ultimately decreases cognitive work. These largely isolated professional development pathways may help explain how different healthcare provider types can have significantly different perceived cognitive workloads while

performing the same task under the same conditions. This was a phenomenon we observed when comparing nurses and physician workload to other, more specialized providers (e.g. respiratory therapists, rehabilitation specialist, wound care specialists, pharmacists, and dieticians). Interestingly, this compartmentalization, while helpful in easing cognitive workload may actually be counterproductive with regards to interprofessional communication and goal sharing on MDR.¹⁸

We identified that clinicians with <5 years of experience had an increased total cognitive workload as compared to more senior clinicians in both the 5-10 year and >10 year groups for both tasks. This difference was driven primarily by the effort and mental demand subscales. Our data affirm that the difficulties experienced by junior ED physicians extend to the ICU, where patients have complex presentations, often with significant pathology in multiple organ systems. Additionally, the ICU is data dense: clinicians must find, interpret, and include hundreds of data elements, some of which may be contradictory to each other, ~~from~~ from disparate sources such as vital signs monitors, laboratory reports, imaging studies, devices, other clinicians, protocols, knowledge bases.¹⁹⁻²⁰ Processing all of this data into an accurate assessment of SOI and determination of POC is a highly demanding task for the expert, let alone the novice. Limiting or simplifying these data inputs may be a way to decrease perceived cognitive workload for novice providers. Video projectors have been used successfully to make patient data available to MDR teams.²¹ Such interventions may have a role in easing the burden of gleaning and recalling relevant data and allowing the novice to devote additional mental work to processing this data into important patient assessments.

Medical students are a group with arguably the least experience on the MDR team. They are operating in what is essentially a novel environment while involving themselves in the care of patients with complex pathophysiology. Students had a significantly higher level of workload as measured by the effort subscale when identifying SOI as compared to physicians with more experience. This

difference is not surprising as we might expect medical students to expend more cognitive effort in order to meaningfully participate in MDR.

Overall perceived total cognitive workload associated with identifying SOI and POC was similar when students, nurses, and physicians were compared. This total load reflected a weighted sum of all 6 NASA-TLX sub scores. These groups reported significantly higher workload for both tasks than the “others” group. This difference may be related to the cultural compartmentalization observed in individual healthcare professional communities, mentioned above. The difference may also be a reflection of differences in the duties and responsibilities commonly assumed by the different clinician types. These data should not be taken to mean the “other” providers on the MDR team contribute less because of their lower perceived workload; to the contrary, efforts should be made to achieve optimally low workloads for all MDR team members.

Increased temporal demand, particularly for nurses and physicians, may stem from a perceived need to see other patients, adhere to schedules, and perform other tasks both patient care related (e.g. family meetings, procedures, documentation) and administrative, educational, or research related (e.g. staff meetings, presentations, protocol development). Temporal demand was a particularly high contributor to total cognitive work at site 3, a busy academic and research oriented facility. Perhaps what goes on outside of MDR and/or away from the ICU drives this perceived increase in temporal demand. Limiting responsibilities outside of the ICU might improve perceived workload.

Interestingly, clinicians at site 1 had the lowest perceived temporal demand, at site 1 but this was not associated with improved perception of performance with respect to identifying SOI. The importance of this finding is not clear as one might expect that less time pressure might improve perceived importance performance.

Another interesting trend was observed when examining perceived workload by years of experience. Those in the 5-10 year groups generally reported lower perceived workload than >10 year

groups. This phenomenon was evident in the mental demand, temporal demand, performance, effort, and frustration subscales observed for SOI, as well as the mental demand, temporal demand, and effort, and frustration subscales for POC. Examination of the composition of the experience groups by healthcare professional type revealed an uneven distribution of provider types across the experience groups. The 5-10 year group was composed of a disproportionately large number of “other” providers, as compared to the <5 and >10 years groups. Given that the “other” providers generally have a lower perceived workload as compared to the student, nurse, and physician groups, this differential helps to explain the observed trend. Certainly there may also be some inherent workload decrease in the 5-10 year group. This may represent a sort of cognitive apex effect, where professional development towards expertise is significant, frequency of task performance is optimal, and the clinicians has the advantage of relative youth compared to more senior counterparts as Tsang has demonstrated.²²

Other authors have shown that a higher cognitive workload, as measured with the NASA-TLX, is associated with increased error rates in medical practice.^{10,11} This study identified relative inexperience as a factor associated with higher workload. We also identified medical students, nurses, and physicians as groups with relatively high associated cognitive workload. When taken together, this suggests that novice nurses and doctors may be at particularly high risk for committing mental errors associated with determination of SOI and POC. Incidentally these groups are responsible for providing much of the direct patient care within the ICU. Patient outcomes may be improved when providers are cognizant of an accurate assessment of the patients’ illness severity.¹² A tool which provides these novice providers with an explicit assessment of patient SOI and by extension, the POC for management has the potential to reduce mental errors and improve outcomes in the ICU.²³

These data are particularly valuable because they were collected at the time of the activity and, therefore, are less likely to be distorted by recall errors. Inter-center variability was examined and no significant differences were detected among study sites. This is consistent with the idea that the work

of MDR is not significantly affected by site specific factors. This speaks well to the external validity of the data generated.

This study had several important limitations. Data collection methods were not identical at each study center due to center-specific workflow differences. This may have introduced some confounding variability into the data set. We were not able to detect any significant inter-center workload differences; however, this study was not designed, and therefore not powered, to support such conclusions. The study was designed primarily to be descriptive of workload. Significant differences were discovered, as detailed above, but the failure to detect a significant difference cannot be taken to mean that such a difference does not exist. Another important limitation is that surveys were conducted on multiple occasions at each center over the course of a relatively short period of time. ~~and,~~ As a consequence, some providers were surveyed more than once. This resulted in an increased ~~in~~ number of data points entered into the analysis relative to the number of discrete survey takers. This may have resulted in an underestimation of variability within the study model.

In conclusion, this study provides insight into the workings of a complex MDR process in the ICU. We were able to identify groups within the MDR that experience higher workload as compared to others when completing two essential, yet often unspoken and unsupported, tasks. Furthermore, by describing the groups which experienced the highest perceived workload during MDR, we have identified the groups which stand to benefit the most from interventions targeted at decreasing the ~~mental~~ cognitive work associated with MDR. The study also serves as an affirmation that the NASA-TLX is an efficient tool which can be utilized ~~in-during~~ a time ~~focused pressured environment~~ event like MDR. These findings support future works aimed at reducing perceived workload, as quantified with the NASA-TLX, during MDR in the Burn ICU.

Acknowledgments

We would like to thank Nicole Caldwell, RN for her instrumental assistance in facilitating the regulatory approvals and administrative support of this project.

References

1. Checkley W, Martin GS, Brown SM, et al. Structure, process, and annual ICU mortality across 69 centers: United States Critical Illness and Injury Trials Group Critical Illness Outcomes Study. *Crit Care Med*. 2014;42(2):344-56
2. Zimmerman JE, Shortell SM, Rousseau DM, et al. Improving intensive care: Observations based on organizational case studies in nine intensive care units: A prospective, multicenter study. *Crit Care Med*. 1993;21(10):1443-51
3. Lane D, Ferri M, Lemaire J, et al. A systematic review of evidence-Informed practices for patient care rounds in the ICU. *Crit Care Med*. 2013;41:2015–2029
4. Ten Have ECM, Hagedoorn M, Holman ND, et al. Assessing the quality of interdisciplinary rounds in the intensive care unit. *J Crit Care* 2013;28:476–482
5. Hart SG, Staveland LE. Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In P. A. Hancock and N. Meshkati (Eds.) *Human Mental Workload*. Amsterdam: North Holland Press. 1988.
6. Hart SG. NASA-Task Load Index (NASA-TLX); 20 years later. *Human Factors and Ergonomics Society Annual Meeting Proceedings*. 2006.
7. Ahmed A, Chandra S, Herasevich V, et al. The effect of two different electronic health record user interfaces on intensive care provider task load, errors of cognition, and performance. *Crit Care Med*. 2011;39:1626–1634
8. Horner RD, Szaflarski JP, Ying J, et al. Physician work intensity among medical specialties: emerging evidence on its magnitude and composition. *Med Care*. 2011;49(11):1007-11
9. Rieger A, Fenger S, Neubert S, et al. Psychophysical workload in the operating room: primary surgeon versus assistant. *Surg Endosc*. 2014 Oct 11. Epub ahead of print.

10. Yurko YY, Scerbo MW, Prabhu AS et al. Higher mental workload is associated with poorer laparoscopic performance as measured by the NASA-TLX tool. *Simul Healthc*. 2010;5(5):267-71
11. Mazur LM, Mosaly PR, Hoyle LM et al. Relating physician's workload with errors during radiation therapy planning. *Practical Rad Onc*. 2014;4(2):71-5
12. Starmer AJ, Spector ND, Srivastava R, et al. Changes in medical errors after implementation of a handoff program. *N Engl J Med* 2014;371:1803-12
13. Critchlow, D. E. and Fligner, M. A. On distribution-free multiple comparisons in the one way analysis of variance. *Communications in Statistics – Theory and Methods* 1991;20:127–139
14. Dreyfus SE and Dreyfus HL. A five-stage model of the mental activities involved in directed skill acquisition. University of California at Berkeley. Operations Research Center. Feb 1980.
15. Benner P. From novice to expert. *Am J of Nursing* 1982;82:402-7
16. Burger JL, Parker K, Cason L, et al. Responses to work complexity: the novice to expert effect. *West J Nurs Res* 2010;32:497
17. Schubert CC, Denmark TK, Crandall B, et al. Characterizing novice-expert differences in macrocognition; an explanatory study of cognitive work in the emergency department. *Ann Emerg Med* 2013;61:96-109
18. Ainsworth CR, Pamplin JC, Allen DA, et al. A bedside communication tool did not improve the alignment of a multidisciplinary team's goals for intensive care unit patients. *J Crit Care* 2013;28:112.e7–112.e13
19. Reddy MC, Pratt W, Dourish P, et al. Asking questions: information needs in a surgical intensive care unit. *Proc AMIA Symp*. 2002:647–651
20. Pamplin JC, Anders S, Brown J, et al. Discovering complexities in critical care and their challenges to health IT design in a Burn ICU. *Crit Care Med* 2013;41:12

21. Gurses AP, Xiao Y. A systematic review of the literature on multidisciplinary rounds to design information technology. *J Am Med Inform Assoc.* 2006;13:267–276
22. Tsang PS, Shaner TL. Age, attention, expertise, and time-sharing performance. *Psychology and Aging.* 1998;13:323-347
23. Pamplin, Jeremy C., Sarah J. Murray, and Kevin K. Chung. "Phases-of-illness paradigm: better communication, better outcomes." *Critical Care.* 2011;15(6):1-7

Figures Legends




Figure 1. NASA-TLX subscale scores and total load for all survey takers for completion of task 1 (“Identify if the patient is better, same, or worse than yesterday,” i.e. determine the severity of illness, SOI), represented as median (line within box), inner quartile range (box), and range (capped lines). Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.




Figure 2. NASA-TLX subscale scores and total load for all survey takers for completion of task 2 (“Identify the priorities of care today” or POC), represented as median (line within box), inner quartile range (box), and range (capped lines). Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.




Figure 3. Median values for the NASA-TLX subscales and overall workload for completing task 1

(“Identify if the patient is better, same, or worse than yesterday,” i.e. determine the severity of illness, SOI), as compared among clinician groups. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

♦ $P < 0.05$ by both methods




Figure 4. Median values for the NASA-TLX subscales and overall workload for completing task 2

("Identify the priorities of care today" or POC), as compared among clinician groups. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

♦ $P < 0.05$ by both methods




Figure 5. Median values for the NASA-TLX subscales and overall workload for completing task 1

("Identify if the patient is better, same, or worse than yesterday," i.e. determine the severity of illness, SOI), as compared among subgroups defined by years of experience. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

◆ $P < 0.05$ by both methods




Figure 6. Median values for the NASA-TLX subscales and overall workload for completing task 2

("Identify the priorities of care today" or POC), as compared among subgroups defined by years of experience. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

♦ $P < 0.05$ by both methods

Tables

Table 1. Demographic data for our convenience sample of clinicians participating in multidisciplinary rounds and willing to participate in this research describing the size, constitution, and years of experience **since graduation from licensing school (or year in school for students)** of the four subgroups of survey takers. *CN – charge nurse, CNS – clinical nurse specialist, HN – head nurse, OT – occupational therapist, RT – respiratory therapist.*

Group	n	Average Experience (range)	< 5 yrs	5-10 yrs	>10 yrs
Physician	41		19	11	11
Medical Intensivist	3	16.3 (10-27)	0	1	2
Burn Surgeon	11	11.8 (0-16)	~~22~~ 2	1	8
Resident	18	2.1 (0-13)	17	0	1
Fellow	9	5.9 (5-6)	0	9	0
Nurse	25		14	5	6
CN/CNS/HN	7	4.3 (0-12)	4	2	1
Nurse	18	8.9 (0-35)	10	3	5
Medical Student	13	2.5 (0-4)	13	0	0
Other Provider	37		4	22	11
Dietician	18	7.4 (5-10)	0	18	0
OT	1	1	1	0	0
Pharmacist	7	12.1 (2-24)	1	3	3
Psychiatry Support	1	13	0	0	1
RT	10	12.6 (4-25)	2	1	7
Total	116		50	38	28

Table 2. NASA-TLX subscale scores for all survey takers for completion of task 1 (“Identify if the patient is better, same, or worse than yesterday,” i.e. determine the severity of illness, SOI) and task 2 (“Identify the priorities of care today” or POC), ranked by median subscale score. This constitutes a rank-order list describing subscale contributors to overall workload, denoted as “Total” below.

Rank Order	Task 1: Severity of Illness		Task 2: Priority of Care	
	Subscale	Average Score (IQR)	Subscale	Average Score (IQR)
1	Mental	150 (90-210)	Mental	120 (80-160)
2	Effort	120 (80-160)	Temporal	100 (60-140)
3	Performance	100 (60-140)	Performance	90 (50-130)
4	Temporal	90 (50-130)	Effort	90 (50-130)
5	Frustration	20 (20-20)	Frustration	20 (20-20)
6	Physical	0 (0-0)	Physical	0 (0-0)
Total	40 (13-67)		43 (18-68)	

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



Title: Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs.

Ian McInnis, DO, MS1
Sarah Murray, MSN2
Maria Serio-Melvin, MSN2
Jay K. Aden, PhD2
Elizabeth Mann-Salinas, PhD2
Kevin K. Chung, MD, FCCM, FACP 2
Todd Huzar, MD3
Steven Wolf, MD, FACS4
Christopher Nemeth, PhD5
Jeremy C. Pamplin, MD, FCCM, FACP2

1. William Beaumont Army Medical Center, Fort Bliss, TX
2. United States Army Institute of Surgical Research, Fort Sam Houston, TX;
3. Memorial Hermann Hospital Texas Medical Center, Houston, TX;
4. University of Texas – Southwestern Medical Center, Dallas, TX;
5. Applied Research Associates, Inc., Fairborn, OH

Address for Correspondence and Reprints:

Jeremy C. Pamplin
3698 Chambers Pass, JBSA Ft. Sam Houston, TX 78234
Telephone: 210-916-3301
Fax: 210-271-0830
Email: jeremy.c.pamplin.mil@mail.mil

External funding: This project is supported by a grant from the US Army Medical Research and Materiel Command Telemedicine and Advanced Technology Research Center (TATRC) (W81XWH-13-2-0011).

Conflicts of interest: The authors declare that there are no conflicts of interest.

Disclosure: This study was conducted under a protocol reviewed and approved by the Brooke Army Medical Center Institutional Review Board and in accordance with the approved protocol. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Title: Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs.

Abstract

Introduction:

Multidisciplinary rounds (MDR) in the Burn Intensive Care Unit serve as an efficient means for clinicians to assess patient status and establish patient care priorities. Both tasks require significant cognitive work, the magnitude of which is relevant because increased cognitive work of task completion has been associated with increased error rates. We sought to quantify this workload during MDR using the National Aeronautics and Space Administration Task Load Index (NASA-TLX).

Methods:

Research staff at 3 academic regional referral burn centers administered the NASA-TLX to clinicians during MDR. Clinicians assessed their workload associated with 1) "Identify[ing] if the patient is better, same, or worse than yesterday" and 2) "Identify[ing] the most important objectives of care for the patient today." Data were collected on clinician type, years of experience, and hours of direct patient care.

Results:

Surveys were administered to 116 total clinicians, 41 physicians, 25 nurses, 13 medical students and 37 clinicians in other roles. Clinicians with less experience reported more cognitive work when completing both tasks ($p < .005$). Clinicians in the "others" group (respiratory therapists, dieticians, pharmacists, etc.) reported less cognitive work than all other groups for both tasks ($p < 0.05$).

Conclusions:

The NASA-TLX was an effective tool for collecting perceptions of cognitive workload associated with MDR. Perceived cognitive work varied by clinician type and experience level when completing 2 key tasks. Less experience was associated with increased perceived work, potentially increasing medical error rates and increasing risk to patients. Creating tools or work processes to reduce cognitive work may improve clinician performance.

Key Words

1. Teaching Rounds
2. Intensive Care
3. Cognitive Work
4. Task Performance and Analysis

Introduction

The burn intensive care unit (ICU) presents a challenging work environment to the healthcare team. Effective care requires the coordinated effort of healthcare professionals from different backgrounds, each member contributing unique perspective toward a common goal. These teams are a necessity given the complexity of managing the medical, administrative, and social aspects of patient care. Multidisciplinary rounds (MDR) is an event that facilitates communication of relevant information among team members and is associated with better outcomes in the ICU.^{1,2,3}

While MDR represent a powerful tool and a logical means for efficient dissemination of information, there is variability in the quality and effectiveness of such rounds. MDR must encompass key aspects of care, to include problems, plans, and goals. The multidisciplinary team must be effectively *managed* to ensure that all important aspects of patient care are addressed. Caution must also be taken to ensure that information is discussed clearly and in such a way that it is accessible to all members of the team who need it; thus, helping to avoid the need for inefficient clarifications with individual team members after MDR have ended.⁴ Input from all team members must be evaluated and integrated by the team leader, typically a physician, in such a way as to create a cohesive, prioritized care plan.²

A series of decisions are made during MDR, some critical, some routine. Likewise the cognitive work that goes into making these decisions is variable. In the same way that physical work is defined in terms of distance and force, cognitive work can be imagined as a function of a task and the perceived effort required to complete it. Different people have different perceived workloads when completing the same task according to their familiarity with the task and their overall experience with the task or similar tasks.⁵ Calculating workload by this definition requires quantification of perceived mental “force.” Hart and Staveland developed the National Aeronautics and Space Administration Task Load Index (NASA-TLX), a survey composed of six subscales, designed to assess perceived workload of NASA

pilots. These subscales include mental demand, physical demand, temporal demand, frustration, effort, and performance as defined in the NASA-TLX Manuscript.⁵ The NASA-TLX has been used broadly in aviation, the automotive industry, and more recently in medicine.⁶⁻¹¹

Lower perceived workload, as measured by the NASA-TLX, has been linked to better performance in medicine.^{10,11} Proper identification of a patient's illness severity may also be associated with improved clinician performance. Starmer and colleagues demonstrated a significant reduction in preventable adverse events after implementation of a tool that increased communication of patient illness severity during change-of-shift handoffs.¹²

Multidisciplinary rounds present a cognitive challenge: to integrate data, make diagnoses, prioritize care tasks, provide education, and to form shared mental models.^{3,4} The magnitude of this challenge, however, has not been well defined. Also unclear are the differences in cognitive workload, if any, experienced by the different types of clinicians who participate in MDR. The current study sought to define the perceived workload associated with the completion of two important tasks during MDR: generating a personal assessment of overall patient condition and identifying the priorities for the day's care plan.

Methods

We conducted this institutional review board approved survey study in three American Burn Association verified regional referral Burn ICUs. The NASA-TLX surveys were administered by research staff to clinicians at each site who participated in MDR during daily clinical practice according to the process outlined below.

Admission rates to these Burn ICUs varies from 270 to 300 patients per year with an average daily census between 2 (site 2) and 5 (site 1 and 3). At each site, MDR are led by an attending physician who was either a burn surgeon credentialed in critical care (sites 1, 2, and 3), or a medical intensivist with extensive burn experience (site 1). Nurse to patient ratios vary among sites from 1-to-1.5 nurses per patient at site 1, to 1 nurse per patient at sites 2 and 3. Attendance at MDR varies, but typically consists of the bedside nurse, rehabilitation specialists, registered dietitians, clinical pharmacists, respiratory therapists, burn fellows, residents, and students.

Multi-Disciplinary Rounds Conduct

MDR typically consisted of a resident presenting the patient by systems followed by other team members providing additional pertinent details and recommendations as they felt appropriate. The attending physician would direct the information flow and sequence of rounds as he felt necessary and would elicit from the team and/or state a care plan for the patient. Additionally, site 1 used a daily communication tool completed by the unit's "charge nurse" (e.g. nursing shift leader). This communication tool consists of a checklist that confirms use/absence of certain quality metrics (i.e. deep venous thrombosis prophylaxis, gastric ulcer prophylaxis, head-of-bed elevation, etc.) and a list of daily tasks that the charge nurse identified during rounds. At site 1, the charge nurse reviews the checklist and reads the identified tasks as the last activity of a patient's rounds prior to the MDR team moving on to the next patient.

Survey Administration:

Research staff administered the NASA-TLX as either a paper (sites 1 & 3) or electronic (site 2) survey to a convenience sample of clinicians present on MDR for at least five discrete rounds events. A “discrete rounds event” was MDR rounds for a single patient. Thus, the NASA-TLX could be administered once per day for five days or five times in a single day for five different patients. Surveys were administered immediately after MDR was complete for a single ICU patient. Each site adjusted the survey collection period to meet the needs of their site and according to patient volume. Surveys were administered during different points in the MDR to account for physical and mental fatigue that may impact workload perception. For example, surveys were administered early in rounds (after the first patient), middle (after the middle patient) or late in rounds (after the last patient).

Prior to administration of the NASA-TLX survey, research staff reviewed a consent document with potential clinician subjects who participated in MDR. Clinicians were not required to participate, but their participation implied consent. Subjects were allowed to quit the study at any time by not completing the survey. Incomplete surveys were not included in the data analysis. Participation was anonymous and no personally identifying information or master list of participants was collected. We collected participant demographics including clinician type, years of experience, and time spent in direct care of the patient.

After the consent was reviewed and a copy provided to subjects, the research staff read directions aloud and then provided the NASA-TLX survey to participants. Time was provided for questions and clarification by the research staff. Clinicians assessed their workload associated with two tasks identified by the following questions: 1) "Identify if the patient is better, same or worse than yesterday" and 2) "Identify the most important objectives of care for the patient today."

Statistical Analysis

For the purpose of this study, we analyzed results according to the following groups: “physician” represents all attending physicians (burn surgeons or medical intensivists) who led or participated in MDR, as well as all physicians in training including those physicians who were part of a certified residency training program or fellowship; “nurse” represented all nurse types including licensed vocational nurses, registered nurses, and clinical nurse specialists; “student” represented medical students; and “other” represented clinicians not in any other category including respiratory therapists, dietitians, occupational or physical therapists, pharmacists, and psychiatry support personnel.

NASA-TLX Scores were calculated as described by Hart⁵: weighted cognitive load was calculated by first determining individual subscale ratings for mental demand, physical, temporal, performance, effort, and frustration, reported on a scale of 0-100 by asking subjects to place a mark on a line divided into 50 equal parts. A researcher then converted this analog representation into a value from 0-100. Individual subscales ratings were given a weighting factor which represented the perceived importance of a given subscale relative to the other subscales. This weight equaled the number of times a subject selected the subscale over another subscale while completing the “sources of workload” portion of the NASA-TLX survey. During this portion of the survey, subjects were asked to determine which subscales were the more important contributors to workload according to a series of 15 pair-wise comparisons. Thus, a weight could range from 0-5 (0 meaning that a subject never chose the subscale compared to other subscales and 5 meaning the subject always chose a subscale when comparing it to other subscales). Individual raw subscale data were multiplied by this weighting factor and were then summed. This sum was then divided by 15 to yield the weighted (total) workload.

The NASA-TLX survey scores were summarized using medians and interquartile ranges for each of the demographic groups. Due to the small sample size and lack of normality, the scores were analyzed using the nonparametric Wilcoxon's Test with the Steel-Dwass correction for multiple test correction.¹³ Significance was established when the p-value was less than 0.05. All analysis was

performed using Statistical Analysis System (SAS) v9.2, SAS Institute Inc., Cary, USA or JMP v10.0, SAS Institute Inc., Cary, USA.

Results

Surveys were administered to a total of 119 clinicians across all sites; however three surveys were incomplete and were excluded from the analysis for a final $n = 116$. The median weighted cognitive load for task 1, identifying if a patient was better, same, or worse than the previous day (i.e. identifying the patient's "severity of illness" or SOI), was 40 (IQR 13-67). The median weighted cognitive load rating for task 2, identifying the most important objectives of care for the patient today (i.e. identifying the patient's "priorities of care" or POC) was 43 (IQR 18-68). Mental demand, temporal demand, performance, and effort all contributed more to the overall cognitive load than did physical demand or frustration subscales (table 1, figures 1-2). Perceived workload varied significantly according to clinician experience and professional background (figures 3-6), but not according to clinician location or time spent with the patient (data not shown). Interestingly, the frustration and physical demand subscales were particularly *non-normal* in their distribution across our sample. Forty percent of respondents reported no frustration and 63% of respondents reported no physical demand for identifying SOI. Numbers were similar for respondents' perception of frustration and physical demand for identifying POCs: 38% and 62% reported zero frustration or physical demand respectively.

Comparing perceived cognitive workload between clinician types revealed several significant differences ($p < 0.05$ for all comparisons, figures 3-4). Students, nurses, and physicians experienced a higher total cognitive workload than did "others" when identifying both SOI and POC. Students experienced more mental demand than "others" when identifying SOI and more than nurses and "others" when identifying POC. Nurses and physicians experienced more temporal demand than the "others" group when identifying SOI, while students, nurses, and physicians experienced more temporal

demand than “others” when identifying POC. Students also reported increased effort associated with identifying SOI as compared to all other groups, and more than physicians and “others” when identifying POC. Students perceived their performance to be worse than “others” when identifying SOI and nurses perceived it to be worse than “others” when identifying POC. Nurses and physicians perceived more frustration than students and “others” when identifying SOI and more than others when identifying POC. Lastly, there were no significant differences between nurses and physicians for either task (figures 3 and 4).

It should be noted that the three largest groups, physicians (n=41), “other” providers (n=37), and nurses (n=25) had an unequal distribution of experience levels within their respective professional groups. The physician group included a large proportion (19 surveys, 46%) with <5 years of experience. The “others” group was largely composed of surveys of individuals with 5-10 years of experience (22 surveys, 59%). The nurses were similar to the physicians, in that a large proportion of those surveyed had <5 years of experience (14 surveys, 56%) (table 2).

Significant differences were also discovered when study participants were compared based upon their years of experience since graduation from licensing school (or year in school for students) ($P < 0.05$ for all comparisons, figures 5-6). Participants with 5-10 years of experience reported the least amount of total cognitive work for both tasks compared to clinicians with <5 or >10 years of experience. Clinicians with > 10 years of experience perceived more total cognitive work for both tasks compared to clinicians with 5-10 years of experience, but less than clinicians with < 5 years of experience. Finally, clinicians with < 5 years of experience perceived the most total cognitive work, which was significantly more than clinicians with > 4 years of experience.

When examining subscales that contribute to total cognitive work, the following observations were made: Clinicians with < 5 years of experience perceived more mental demand than those with >4 years of experience when identifying SOI and more than clinicians with 5-10 years of experience when

identifying POC. Clinicians with < 5 years of experience perceived more temporal demand and felt their performance was worse for both tasks when compared to clinicians with 5-10 years of experience. Both tasks were perceived as requiring more effort for clinicians with < 5 years of experience compared to those with > 4 years of experience. Clinicians with 5-10 years of experience perceived significantly less frustration while identifying SOI than did those with < 5 and those with > 10 years of experience, but only less than those with > 10 years of experience when identifying POC. Finally, there was no apparent impact of experience on perceived physical demand. The largest differences in workload perception were consistently between those with < 5 years and those with 5-10 years of experience for all domains except frustration (see figures 5 and 6).

There were no significant differences in perceived total workload when comparing study sites or hours of direct patient care with respect to identifying either SOI or POC. However, there were differences between sites for some subscale domains. Site 3 clinicians perceived significantly more temporal demand than site 1 when identifying SOI (median 150 [IQR 90-210] vs. 60 [40-80], $p < 0.05$). Site 1 clinicians perceived significantly worse performance when identifying SOI than did clinicians at site 3 (median 140 [IQR 90-190] vs. 80 [IQR 40-120], $p < 0.05$). Clinicians at site 2 perceived significantly less frustration than clinicians at site 1 or site 3 for both tasks. Site 2 perceived no frustration when identifying SOI or POC, whereas clinicians at site 1 and site 3 perceived frustration as a relatively low contributor to workload when identifying SOI (site 1 & 2 frustration subscale score 20-30 [IQR 75% 0-120]) and when identifying POC (site 1 & 2 frustration subscale score 30 [IQR 75% 0-120]).

Discussions

This manuscript is the first to describe the cognitive work performed by clinicians during MDR in the Burn ICU. The NASA-TLX effectively revealed workload perception differences and similarities in cognitive work associated with completing two critical tasks performed during MDR: identifying a

patient's condition (severity of illness, SOI) and prioritizing associated treatments (priority of care, POC).

Significant findings include:

1. Mental demand, temporal demand, performance, and effort were the primary determinants for the cognitive work performed for the identified tasks on MDR with mental demand being greatest;
2. students, nurses, and physicians all had higher perceived total workload for both SOI and POC than "others";
3. students perceived the most effort on rounds and had the most mental demand when identifying POC and significantly more than "others" when identifying SOI;
4. students, nurses, and physicians experienced significantly more temporal demand when identifying POC than did "other" healthcare providers while only nurses and physicians perceived this significantly more than others for identifying SOI;
5. clinicians with the least experience had higher perceived workload when identifying SOI and POC as compared to those with more experience;
6. and some individuals perceived more frustration and physical demand than most others during MDR.

Participation in MDR ~~is~~ was associated with a moderate cognitive workload for nurses, physicians, students, and other healthcare professionals in attendance when completing two key tasks. Quantification of this workload with the NASA-TLX revealed unequal distribution of workload across the subscales of the task load index. Greater contributors to total load were mental demand, temporal demand, performance, and effort, with mental demand consistently the greatest for both tasks and compared to other subscales. Less substantial contributors were frustration, followed by physical demand. MDR is largely a thought exercise and so it is not surprising that participants reported relatively low physical demand. When taken as a whole, the MDR participants reported a low cognitive

workload associated with frustration, particularly at site 2. This low observed frustration level may be a reflection of how well the MDR participants are acclimated to the MDR process, experiencing relatively low levels of stress, annoyance, and irritation when participating in this well-practiced event or that site 2 clinicians work particularly well together, or have some other organizational culture/aspect (e.g. a lower census) that minimizes frustration. Also interesting was that the frustration and physical demand subscales demonstrated a significant right skew effect, driven by a large number of participants reporting little or no load associated with these 2 subscales. This was appreciated at site 2 also. This suggests that certain *individuals* may experience MDR differently than most others. In particular, some individuals perceive remarkably more frustration (figures 1-2).

This study was designed to be descriptive of cognitive workload associated with MDR. We did not seek to find significant between group differences and yet such differences were found. Why do these differences exist and what impact do they have on patient care? We suspect that clinicians undergo a significant paradigm shift in their approach to medical decision making during the transition from novice to expert. A general stepwise paradigm of skill acquisition has been described previously, and applied to the healthcare setting in prior works.¹⁴⁻¹⁶ One study identified significant macro cognitive differences in the approach of a novice physician in the emergency department as compared to an expert. Junior physicians had difficulty integrating individual aspects of a patient presentation into a cogent picture. The same study showed that less experienced physicians had an overreliance on laboratory data and had difficulty integrating data that was inconsistent with their current diagnosis.¹⁷ Furthermore, differences in professional background, especially those associated with increased specialization with career progression, promote differences in knowledge acquisition and information use that simplifies mental modeling, role based planning, and ultimately decreases cognitive work. These largely isolated professional development pathways may help explain how different healthcare provider types can have significantly different perceived cognitive workloads while performing the same

task under the same conditions. This was a phenomenon we observed when comparing nurses and physician workload to other, more specialized providers (e.g. respiratory therapists, rehabilitation specialist, wound care specialists, pharmacists, and dieticians). Interestingly, this compartmentalization, while helpful in easing cognitive workload may actually be counterproductive with regards to interprofessional communication and goal sharing on MDR.¹⁸

We identified that clinicians with <5 years of experience had an increased total cognitive workload as compared to more senior clinicians in both the 5-10 year and >10 year groups for both tasks. This difference was driven primarily by the effort and mental demand subscales. Our data affirm that the difficulties experienced by junior ED physicians extend to the ICU, where patients have complex presentations, often with significant pathology in multiple organ systems. Additionally, the ICU is data dense: clinicians must find, interpret, and include hundreds of data elements, some of which may be contradictory to each other, from disparate sources such as vital signs monitors, laboratory reports, imaging studies, devices, other clinicians, protocols, knowledge bases.¹⁹⁻²⁰ Processing all of this data into an accurate assessment of SOI and determination of POC is a highly demanding task for the expert, let alone the novice. Limiting or simplifying these data inputs may be a way to decrease perceived cognitive workload for novice providers. Video projectors have been used successfully to make patient data available to MDR teams.²¹ Such interventions may have a role in easing the burden of gleaning and recalling relevant data and allowing the novice to devote additional mental work to processing this data into important patient assessments.

Medical students are a group with arguably the least experience on the MDR team. They are operating in what is essentially a novel environment while involving themselves in the care of patients with complex pathophysiology. Students had a significantly higher level of workload as measured by the effort subscale when identifying SOI as compared to physicians with more experience. This

difference is not surprising as we might expect medical students to expend more cognitive effort in order to meaningfully participate in MDR.

Overall perceived total cognitive workload associated with identifying SOI and POC was similar when students, nurses, and physicians were compared. This total load reflected a weighted sum of all 6 NASA-TLX sub scores. These groups reported significantly higher workload for both tasks than the “others” group. This difference may be related to the cultural compartmentalization observed in individual healthcare professional communities, mentioned above. The difference may also be a reflection of differences in the duties and responsibilities commonly assumed by the different clinician types. These data should not be taken to mean the “other” providers on the MDR team contribute less because of their lower perceived workload; to the contrary, efforts should be made to achieve optimally low workloads for all MDR team members.

Increased temporal demand, particularly for nurses and physicians, may stem from a perceived need to see other patients, adhere to schedules, and perform other tasks both patient care related (e.g. family meetings, procedures, documentation) and administrative, educational, or research related (e.g. staff meetings, presentations, protocol development). Temporal demand was a particularly high contributor to total cognitive work at site 3, a busy academic and research oriented facility. Perhaps what goes on outside of MDR and/or away from the ICU drives this perceived increase in temporal demand. Limiting responsibilities outside of the ICU might improve perceived workload.

Interestingly, clinicians at site 1 had the lowest perceived temporal demand, but this was not associated with improved perception of performance with respect to identifying SOI. The importance of this finding is not clear as one might expect that less time pressure might improve perceived performance.

Another interesting trend was observed when examining perceived workload by years of experience. Those in the 5-10 year groups generally reported lower perceived workload than >10 year

groups. This phenomenon was evident in the mental demand, temporal demand, performance, effort, and frustration subscales observed for SOI, as well as the mental demand, temporal demand, and effort, and frustration subscales for POC. Examination of the composition of the experience groups by healthcare professional type revealed an uneven distribution of provider types across the experience groups. The 5-10 year group was composed of a disproportionately large number of “other” providers, as compared to the <5 and >10 years groups. Given that the “other” providers generally have a lower perceived workload as compared to the student, nurse, and physician groups, this differential helps to explain the observed trend. Certainly there may also be some inherent workload decrease in the 5-10 year group. This may represent a sort of cognitive apex effect, where professional development towards expertise is significant, frequency of task performance is optimal, and the clinician has the advantage of relative youth compared to more senior counterparts as Tsang has demonstrated.²²

Other authors have shown that a higher cognitive workload, as measured with the NASA-TLX, is associated with increased error rates in medical practice.^{10,11} This study identified relative inexperience as a factor associated with higher workload. We also identified medical students, nurses, and physicians as groups with relatively high associated cognitive workload. When taken together, this suggests that novice nurses and doctors may be at particularly high risk for committing mental errors associated with determination of SOI and POC. Incidentally these groups are responsible for providing much of the direct patient care within the ICU. Patient outcomes may be improved when providers are cognizant of an accurate assessment of the patients’ illness severity.¹² A tool which provides these novice providers with an explicit assessment of patient SOI and by extension, the POC for management has the potential to reduce mental errors and improve outcomes in the ICU.²³

These data are particularly valuable because they were collected at the time of the activity and, therefore, are less likely to be distorted by recall errors. Inter-center variability was examined and no significant differences were detected among study sites. This is consistent with the idea that the work

of MDR is not significantly affected by site specific factors. This speaks well to the external validity of the data generated.

This study had several important limitations. Data collection methods were not identical at each study center due to center-specific workflow differences. This may have introduced some confounding variability into the data set. We were not able to detect any significant inter-center workload differences; however, this study was not designed, and therefore not powered, to support such conclusions. The study was designed primarily to be descriptive of workload. Significant differences were discovered, as detailed above, but the failure to detect a significant difference cannot be taken to mean that such a difference does not exist. Another important limitation is that surveys were conducted on multiple occasions at each center over the course of a relatively short period of time. As a consequence, some providers were surveyed more than once. This resulted in an increased number of data points entered into the analysis relative to the number of discrete survey takers. This may have resulted in an underestimation of variability within the study model.

In conclusion, this study provides insight into the workings of a complex MDR process in the ICU. We were able to identify groups within the MDR that experience higher workload as compared to others when completing two essential, yet often unspoken and unsupported, tasks. Furthermore, by describing the groups which experienced the highest perceived workload during MDR, we have identified the groups which stand to benefit the most from interventions targeted at decreasing the cognitive work associated with MDR. The study also serves as an affirmation that the NASA-TLX is an efficient tool which can be utilized during a time pressured event like MDR. These findings support future works aimed at reducing perceived workload, as quantified with the NASA-TLX, during MDR in the Burn ICU.

Acknowledgments

We would like to thank Nicole Caldwell, RN for her instrumental assistance in facilitating the regulatory approvals and administrative support of this project.

References

1. Checkley W, Martin GS, Brown SM, et al. Structure, process, and annual ICU mortality across 69 centers: United States Critical Illness and Injury Trials Group Critical Illness Outcomes Study. *Crit Care Med*. 2014;42(2):344-56
2. Zimmerman JE, Shortell SM, Rousseau DM, et al. Improving intensive care: Observations based on organizational case studies in nine intensive care units: A prospective, multicenter study. *Crit Care Med*. 1993;21(10):1443-51
3. Lane D, Ferri M, Lemaire J, et al. A systematic review of evidence-Informed practices for patient care rounds in the ICU. *Crit Care Med*. 2013;41:2015–2029
4. Ten Have ECM, Hagedoorn M, Holman ND, et al. Assessing the quality of interdisciplinary rounds in the intensive care unit. *J Crit Care* 2013;28:476–482
5. Hart SG, Staveland LE. Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In P. A. Hancock and N. Meshkati (Eds.) *Human Mental Workload*. Amsterdam: North Holland Press. 1988.
6. Hart SG. NASA-Task Load Index (NASA-TLX); 20 years later. *Human Factors and Ergonomics Society Annual Meeting Proceedings*. 2006.
7. Ahmed A, Chandra S, Herasevich V, et al. The effect of two different electronic health record user interfaces on intensive care provider task load, errors of cognition, and performance. *Crit Care Med*. 2011;39:1626–1634
8. Horner RD, Szaflarski JP, Ying J, et al. Physician work intensity among medical specialties: emerging evidence on its magnitude and composition. *Med Care*. 2011;49(11):1007-11
9. Rieger A, Fenger S, Neubert S, et al. Psychophysical workload in the operating room: primary surgeon versus assistant. *Surg Endosc*. 2014 Oct 11. Epub ahead of print.

10. Yurko YY, Scerbo MW, Prabhu AS et al. Higher mental workload is associated with poorer laparoscopic performance as measured by the NASA-TLX tool. *Simul Healthc*. 2010;5(5):267-71
11. Mazur LM, Mosaly PR, Hoyle LM et al. Relating physician's workload with errors during radiation therapy planning. *Practical Rad Onc*. 2014;4(2):71-5
12. Starmer AJ, Spector ND, Srivastava R, et al. Changes in medical errors after implementation of a handoff program. *N Engl J Med* 2014;371:1803-12
13. Critchlow, D. E. and Fligner, M. A. On distribution-free multiple comparisons in the one way analysis of variance. *Communications in Statistics – Theory and Methods* 1991;20:127–139
14. Dreyfus SE and Dreyfus HL. A five-stage model of the mental activities involved in directed skill acquisition. University of California at Berkeley. Operations Research Center. Feb 1980.
15. Benner P. From novice to expert. *Am J of Nursing* 1982;82:402-7
16. Burger JL, Parker K, Cason L, et al. Responses to work complexity: the novice to expert effect. *West J Nurs Res* 2010;32:497
17. Schubert CC, Denmark TK, Crandall B, et al. Characterizing novice-expert differences in macrocognition; an explanatory study of cognitive work in the emergency department. *Ann Emerg Med* 2013;61:96-109
18. Ainsworth CR, Pamplin JC, Allen DA, et al. A bedside communication tool did not improve the alignment of a multidisciplinary team's goals for intensive care unit patients. *J Crit Care* 2013;28:112.e7–112.e13
19. Reddy MC, Pratt W, Dourish P, et al. Asking questions: information needs in a surgical intensive care unit. *Proc AMIA Symp*. 2002:647–651
20. Pamplin JC, Anders S, Brown J, et al. Discovering complexities in critical care and their challenges to health IT design in a Burn ICU. *Crit Care Med* 2013;41:12

21. Gurses AP, Xiao Y. A systematic review of the literature on multidisciplinary rounds to design information technology. *J Am Med Inform Assoc.* 2006;13:267–276
22. Tsang PS, Shaner TL. Age, attention, expertise, and time-sharing performance. *Psychology and Aging.* 1998;13:323-347
23. Pamplin, Jeremy C., Sarah J. Murray, and Kevin K. Chung. "Phases-of-illness paradigm: better communication, better outcomes." *Critical Care.* 2011;15(6):1-7

Figures Legends




Figure 1. NASA-TLX subscale scores and total load for all survey takers for completion of task 1 (“Identify if the patient is better, same, or worse than yesterday,” i.e. determine the severity of illness, SOI), represented as median (line within box), inner quartile range (box), and range (capped lines). Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.




Figure 2. NASA-TLX subscale scores and total load for all survey takers for completion of task 2 (“Identify the priorities of care today” or POC), represented as median (line within box), inner quartile range (box), and range (capped lines). Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.




Figure 3. Median values for the NASA-TLX subscales and overall workload for completing task 1

("Identify if the patient is better, same, or worse than yesterday," i.e. determine the severity of illness, SOI), as compared among clinician groups. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

◆ $P < 0.05$ by both methods




Figure 4. Median values for the NASA-TLX subscales and overall workload for completing task 2

("Identify the priorities of care today" or POC), as compared among clinician groups. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

♦ $P < 0.05$ by both methods




Figure 5. Median values for the NASA-TLX subscales and overall workload for completing task 1

("Identify if the patient is better, same, or worse than yesterday," i.e. determine the severity of illness, SOI), as compared among subgroups defined by years of experience. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

◆ $P < 0.05$ by both methods




Figure 6. Median values for the NASA-TLX subscales and overall workload for completing task 2

("Identify the priorities of care today" or POC), as compared among subgroups defined by years of experience. Weighted subscale values range from 0-500 (subscale value 0-100 x subscale weight⁵). The total load scales range from 0-100 and are equal to a sum of the weighted subscale values divided by 15.

* $P < 0.05$ by Steel-Dwass Method

+ $P < 0.05$ by Wilcoxon Method

♦ $P < 0.05$ by both methods

Tables

Table 1. Demographic data for our convenience sample of clinicians participating in multidisciplinary rounds and willing to participate in this research describing the size, constitution, and years of experience since graduation from licensing school (or year in school for students) of the four subgroups of survey takers. *CN – charge nurse, CNS – clinical nurse specialist, HN – head nurse, OT – occupational therapist, RT – respiratory therapist.*

Group	n	Average Experience (range)	< 5 yrs	5-10 yrs	>10 yrs
Physician	41		19	11	11
Medical Intensivist	3	16.3 (10-27)	0	1	2
Burn Surgeon	11	11.8 (0-16)	~~22~~ 2	1	8
Resident	18	2.1 (0-13)	17	0	1
Fellow	9	5.9 (5-6)	0	9	0
Nurse	25		14	5	6
CN/CNS/HN	7	4.3 (0-12)	4	2	1
Nurse	18	8.9 (0-35)	10	3	5
Medical Student	13	2.5 (0-4)	13	0	0
Other Provider	37		4	22	11
Dietician	18	7.4 (5-10)	0	18	0
OT	1	1	1	0	0
Pharmacist	7	12.1 (2-24)	1	3	3
Psychiatry Support	1	13	0	0	1
RT	10	12.6 (4-25)	2	1	7
Total	116		50	38	28

Table 2. NASA-TLX subscale scores for all survey takers for completion of task 1 (“Identify if the patient is better, same, or worse than yesterday,” i.e. determine the severity of illness, SOI) and task 2 (“Identify the priorities of care today” or POC), ranked by median subscale score. This constitutes a rank-order list describing subscale contributors to overall workload, denoted as “Total” below.

Rank Order	Task 1: Severity of Illness		Task 2: Priority of Care	
	Subscale	Average Score (IQR)	Subscale	Average Score (IQR)
1	Mental	150 (90-210)	Mental	120 (80-160)
2	Effort	120 (80-160)	Temporal	100 (60-140)
3	Performance	100 (60-140)	Performance	90 (50-130)
4	Temporal	90 (50-130)	Effort	90 (50-130)
5	Frustration	20 (20-20)	Frustration	20 (20-20)
6	Physical	0 (0-0)	Physical	0 (0-0)
Total	40 (13-67)		43 (18-68)	

OPEN ACCESS LICENSE AGREEMENT

This OPEN ACCESS LICENSE AGREEMENT (this “Agreement”), dated as of October 9th , 2015 (the “Effective Date”), by and between Wolters Kluwer Health, Inc., operating as Medical Research / Lippincott Williams & Wilkins, a Delaware corporation, having its principal place of business at Two Commerce Square, 2001 Market Street, Philadelphia, PA 19103 (the “Publisher”), and the corresponding author listed on Schedule A to this Agreement (the “Author”, and together with the Publisher, the “Parties”).

1. Grant of License

The Author hereby grants to the Publisher and its Affiliates the exclusive, worldwide, royalty free, perpetual (for the duration of the applicable copyright) right and license to use the Work for all commercial or educational purposes, including, but not limited to, publishing, reproducing, marketing, distributing (themselves and through distributors), sublicensing, and selling copies of the Work throughout the world for the Term. If the Author is a United States government employee, such license grant shall be limited to the extent the Author is able to grant such license.

2. Warranties, Indemnification, and Limitation of Liability

a. The Author represents and warrants that:

(i) it has the right and power to enter into this Agreement, to grant the rights and licenses granted pursuant to this Agreement, and to perform all of its other obligations contained in this Agreement;

(ii) it has not previously assigned, transferred or otherwise encumbered the rights or licenses granted pursuant to this Agreement; and that the person executing this Agreement on the Author’s behalf is authorized to do so;

(iii) the Work and the licenses granted herein do not and will not infringe upon, violate or misappropriate any intellectual property rights or any other proprietary right, contract or other right or interest of any third party;

(iv) if the Work is a multi-authored Work, the Author has obtained written permission from each author of the Work to enter into this Agreement on behalf such author, and each such author has read, understands and has agreed to the terms of this Agreement; and

(v) the Author has obtained any necessary releases and permissions to quote from other sources in the Work and to include any works and materials in the Work and all such releases and permissions are in full force and effect.

b. The Author hereby indemnifies the Publisher and its directors, officers, employees, agents, and representatives and agrees to defend and hold them harmless from and against any and all liability, damage, loss, costs or expenses (including reasonable

attorney's fees and costs of settlement) incurred by any such party arising out of, or relating to any misrepresentation in, or breach or alleged breach of the Author's representations or warranties in this Agreement. If the Author fails to promptly or diligently pursue any defense of any indemnified party, the indemnified parties, or any of them, may assume such defense at the Author's expense. The obligations of this indemnification will survive any termination or expiration of this Agreement.

c. The Publisher represents and warrants that it has the right and power to enter into this Agreement and to perform its obligations contained in this Agreement, and that the person executing this Agreement on the Publisher's behalf is authorized to do so.

d. The Publisher hereby indemnifies the Author and agrees to defend and hold the Author harmless from and against any and all liability, damage, loss, costs or expenses (including reasonable attorney's fees and costs of settlement) incurred by the Author arising out of, or relating to any misrepresentation in, or breach or alleged breach of the Publisher's representations or warranties in this Agreement. If the Publisher fails to promptly or diligently pursue any defense of the Author, the Author may assume such defense at the Publisher's expense. The obligations of this indemnification will survive any termination or expiration of this Agreement.

e. EXCEPT AS OTHERWISE SET FORTH IN THIS AGREEMENT, NEITHER PARTY MAKES ANY OTHER, AND HEREBY DISCLAIMS ALL OTHER, REPRESENTATIONS AND WARRANTIES OF ANY KIND, WHETHER EXPRESS, IMPLIED, STATUTORY OR OTHERWISE, INCLUDING, WITHOUT LIMITATION, WARRANTIES OF TITLE, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, NONINFRINGEMENT, OR THE ABSENCE OF LATENT OR OTHER DEFECTS, ACCURACY, OR THE PRESENCE OR ABSENCE OF ERRORS, WHETHER OR NOT DISCOVERABLE.

f. EXCEPT TO THE EXTENT REQUIRED BY APPLICABLE LAW, IN NO EVENT WILL EITHER PARTY BE LIABLE TO THE OTHER PARTY BASED UPON ANY LEGAL THEORY FOR ANY SPECIAL, INCIDENTAL, CONSEQUENTIAL, PUNITIVE, OR EXEMPLARY DAMAGES ARISING OUT OF THIS LICENSE OR THE USE OF THE WORK, EVEN IF A PARTY HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES.

3. Creative Commons License.

a. If the Author has not selected either "Research Councils UK (RCUK)" or "Wellcome Trust" in Item 1 of Schedule B, the following shall apply:

The Author acknowledges and agrees that the Work will be published by the Publisher in [INSERT NAME OF JOURNAL] (the "Journal") and made freely available to users under the terms of the Attribution-NonCommercial-NoDerivs 3.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode> (the "CC BY-NC-ND"). The Author acknowledges and agrees that that

Publisher is the exclusive “Licensor”, as defined in the CC BY-NC-ND, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY-NC-ND.

- b. If the Author has selected either “Research Councils UK (RCUK)” or “Wellcome Trust” in Item 1 of Schedule B and has selected “Gold Route” in Item 2 of Schedule B, the following shall apply:

The Author acknowledges and agrees that the Work will be published by the Publisher in [INSERT NAME OF JOURNAL] (the “Journal”) and made freely available to users under the terms of the Attribution 3.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by/3.0/legalcode> (the “CC BY”). The Author acknowledges and agrees that that Publisher is the exclusive “Licensor”, as defined in the CC BY, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY.

- c. If the Author has selected either “Research Councils UK (RCUK)” or “Wellcome Trust” in Item 1 of Schedule B and has selected “Green Route” in Item 2 of Schedule B, the following shall apply:

The Author acknowledges and agrees that the Work will be published by the Publisher in [INSERT NAME OF JOURNAL] (the “Journal”) and made freely available to users under the terms of the Attribution-NonCommercial 3.0 Creative Commons License, as currently displayed at <http://creativecommons.org/licenses/by-nc/3.0/legalcode> (the “CC BY-NC”). The Author acknowledges and agrees that that Publisher is the exclusive “Licensor”, as defined in the CC BY-NC, of the Work and that the Publisher may make the Work freely available to all users under the terms of the CC BY-NC.

4. Royalties.

The Author acknowledges and agrees that this Agreement entitles the Author to no royalties or fees. To the maximum extent permitted by law, the Author waives any and all rights the Author may have to collect royalties or other fees in relation to the Work or in respect of any use of the Work by the Publisher or its sublicensees.

5. Miscellaneous.

- a. Assignment. This Agreement may not be assigned or transferred, in whole or in part, by either party without the prior written consent of the other party. Notwithstanding the above, the Publisher may assign this Agreement without the written consent of the Author (i) to an entity succeeding, whether by sale, merger or other corporate reorganization, to substantially all of the Publisher’s assets and business activity, or (ii) to a corporation or organization that obtains the right to publish the Journal from the Publisher. The Publisher may assign this Agreement to any of its affiliates. This

Agreement will be binding upon and inure to the benefit of the parties hereto and their respective successors and permitted assigns.

b. Counterparts. This Agreement may be executed in two or more counterparts, each of which shall be deemed an original, but all of which together shall constitute one and the same document. Facsimile or Portable Document Format (PDF) signatures will be deemed original signatures for purposes of this Agreement.

c. Entire Agreement; Amendment. This Agreement sets forth the entire agreement of the parties on the subject hereof and supersedes all previous or contemporaneous oral or written representations or agreements relating to the rights and duties provided herein, and may not be modified or amended except by written agreement of the parties.

d. Force Majeure. Neither party shall be liable for any default or delay on its part in performing any obligation under this Agreement if such default or delay is caused by natural disaster, accident, war, civil disorder, strike or any other cause beyond the reasonable control of such party. In the event that either party is prevented by such an occurrence or circumstance for a period of more than ninety (90) days from fulfilling its obligations under this Agreement, the other party may terminate this Agreement upon thirty (30) days' written notice.

e. Governing Law. This Agreement shall be governed in all respects according to the laws of the State of New York without giving effect to the principles of conflict of law thereof.

f. Headings. All headings are for reference purposes only and shall not affect the meaning or interpretation of any provision hereof.

g. Severability. If any provision of this Agreement is held to be illegal, invalid, or unenforceable under the present or future laws, then such provision shall be revised by a court of competent jurisdiction to be enforceable if permitted under applicable law, and otherwise shall be fully severable. In any event, this Agreement shall be construed and enforced as if such illegal, invalid, or unenforceable provision had never comprised a part of this Agreement, and the remaining provisions of this Agreement shall remain in full force and effect and shall not be affected by the illegal, invalid, or unenforceable provision or by its severance from this Agreement.

h. Status of the Parties. The parties are independent contractors. Nothing in this Agreement is intended to or shall be construed to constitute or establish any agency, joint venture, partnership or fiduciary relationship between the parties, and neither party has the right or authority to bind the other party nor shall either party be responsible for the acts or omissions of the other.

i. Waiver; Amendment. The waiver by either party of or the failure by either party to claim a breach of any provision of this Agreement shall not be, or be held to be, a waiver of any subsequent breach or affect in any way the further effectiveness of any such provision. No term or condition of this Agreement may be waived except by an agreement by the parties in writing.

j. Waiver of Jury Trial. EACH PARTY HEREBY WAIVES ITS RIGHT TO A JURY TRIAL IN CONNECTION WITH ANY DISPUTE OR LEGAL PROCEEDING ARISING OUT OF THIS AGREEMENT OR THE SUBJECT MATTER HEREOF.

[Signature Page Follows]

IN WITNESS WHEREOF, each party to this Agreement has caused this Agreement, effective as of the Effective Date, to be signed by its duly authorized representative.

AUTHOR

Print Name: Pamplin, Jeremy

WOLTERS KLUWER HEALTH, INC., OPERATING AS MEDICAL RESEARCH /
LIPPINCOTT WILLIAMS & WILKINS

By: _____

Name:

Title:

Schedule A

This Schedule A must be completed by Author in its entirety. The Publisher is unable to publish the Work unless this Schedule A is completely filled out.

Article Tracking #: JBCR-D-15-00153

Article Title (the “Work”): Comparing the workload perceptions of identifying patient condition and priorities of care among burn providers in three Burn ICUs

Corresponding Author Name (the “Author”) (please print): Jeremy C. Pamplin

Copyright Owner’s Name (please print): U.S. Army

Name of Journal in which Work is to be Published: The Journal of Burn Care & Research

Schedule B

This Schedule B must be completed by Author in its entirety. The Publisher is unable to publish the Work unless this Schedule B is completely filled out.

PUBLIC ACCESS POLICY FUNDING DISCLOSURE

1. Please disclose below if you or any other author of the Work has received funding for research on which the Work is based from any of the following organizations:


- ☐ National Institutes of Health (NIH)
- ☐ Howard Hughes Medical Institute (HHMI)
- ☐ Research Councils UK (RCUK) (Please complete Item 2)
- ☐ Wellcome Trust (Please complete Item 2)

2. If you have selected either Research Councils UK or Wellcome Trust from the above list, please disclose the Open Access option to which the Work will be subject:

- ☐ Gold route
- ☐ Green route



NOTE: If the “Gold” route has been selected, Section 3.b. of the Agreement will apply to the Work, and neither Section 3.a. nor Section 3.c. of the Agreement will apply to the Work. If the “Green” route has been selected, Section 3.c. of the Agreement will apply to the Work, and neither Section 3.a. nor Section 3.b. of the Agreement will apply to the Work.


3. ☐ This Schedule B is inapplicable to the Work.



[Click here to access/download](#)

Copyright Transfer and Disclosure Form
Chung JBCR copyrightTransfer.pdf








[Click here to access/download](#)

Copyright Transfer and Disclosure Form



Mann JBCR copyrightTransfer.pdf






[Click here to access/download](#)



Copyright Transfer and Disclosure Form
McInnis JBCR copyrightTransfer.pdf






[Click here to access/download](#)

Copyright Transfer and Disclosure Form
Murray JBCR copyrightTransfer.pdf








[Click here to access/download](#)

Copyright Transfer and Disclosure Form



Pamplin JBCR copyrightTransfer.pdf






[Click here to access/download](#)

Copyright Transfer and Disclosure Form
SerioMelvin JBCR copyrightTransfer.pdf





[Click here to access/download](#)

Copyright Transfer and Disclosure Form
Wolf JBCR ASAP Copyright form.PDF




CORE Site (USAISR) Tools

1. Original TeamView
2. Original Scales Tool
3. Updated TeamView
4. Updated ECMO TeamView
5. Updated Resident Scales/Checklist/Presentation Tool
6. Updated Nurse Scales/Checklist/Handoff Tool

Team View

Patient Severity of Illness, Status, and Goals

Severity of Illness (transfer from scales tool)



Status

% Open																				
Hrs Sleep Last Night																				
Coordinating Activities																				
See code list behind Scales Tool for codes.																				

Update Checklist items daily!

	NA	Good	Discuss
GI Prophylaxis	☐	☐	☐
CHG Prophylaxis	☐	☐	☐
HOB > 30 degrees	☐	☐	☐
DVT Prophylaxis	☐	☐	☐
Last BM < 48 hrs ago	☐	☐	☐
Respiratory Orders Current	☐	☐	☐
Family has questions	☐		☐
Delirium Positive	☐		☐
Pending Cultures	☐		☐

Antibiotic(s)	NA ☐	Effective Start Date	Planned Stop Date	Diagnosis

Goals (completed during rounds)

Fluid volume goal for today:

- ☐ Positive
- ☐ Negative
- ☐ Even

Major Goal(s) for the next 24-48 hours:

A. Patient’s Current Condition.

Room:_____Date/Time:_____

Please make a noticeable “X” ANYWHERE on ANY of the scales below that indicates your estimate of the patient’s condition right now.

Diagnoses & Problems

Worse = Increasing in number or severity; Better = Decreasing in number or severity

Getting Worse Quickly

Worse

Same

Better

Better Quickly

Organ support

(Mechanical Ventilation, CRRT, Vasopressors, Blood Products, Etc.)

Maximal or Increasing rapidly

High, new, or increasing in amount or number

Stable in amount or Number

Low or Decreasing in amount or number

Permanent or Chronic

None

Mental Status

Paralyzed

Comatose Sedated

Agitated Delirium or Encephalopathy

Hypoactive Delirium or Follows Commands

Alert, Normal, [Baseline]

Acuity Level

Highest/Very Complex

High or Increasing

Standard or Decreasing

Low

Wounds

Fungus/necrosis Wound Fialure (<20% healed @ 3 weeks)

Unexcised full thickness Delayed healing < 50% @ 3 weeks

Stable Healing

Small/Minimal < 10% Open

Risk of Worsening or Sepsis

High

Medium

Low

Severity of Illness (your summary assessment of the above scales)

Most sick, Could Die

Least Sick Could transfer

B. Patient’s Current Treatments

Make a noticeable “X” ANYWHERE the scales below that indicates the patient’s current treatments.

Text in each section is organized by objective/goal in bold type, recommendations in regular type, and considerations in italic type.

Analgesia & Sedation

Deep Sedation +/-Paralyzed

*Decrease O₂ Consumption

Continuous + PRN Narcotics and Sedatives

Moderate Sedation (RASS -1 to -2)

*Safety and Comfort

Scheduled/PRN Narcotics PRN Sedatives

Light Sedation (RASS 0 to -1)

*Patient Comfort, Minmal Pain

PRN/Scheduled Narcotics +/- Scheduled Enteral Narcotics

No Sedation (RASS 0)

*Participatory Patient

PO/Enteral PRN Narcotics Symptomatic Tx of Anxiety

Sleep

Goal: Minimize Delirium

As Able

Day/Night Cycle

Goal 4-6 hours

Avoid awakening 4-6 hrs at night Day/Night Cycle

Use Sleep Protocol

Avoid awakening 6-8 hrs at night Day/Night Cycle Sleep aid

Ventilation

Goal: Minimize VILI, liberation ASAP

Controlled/Assisted

(vent does Work of Breathing)

Low Tidal Volume (Vt) or VDR

Open Lung Approach: ↑PEEP, ↓Vt

Wean/Decrease with Supported Mode

Transition to APRV or CPAP or CPAP/PS

Decrease FIO₂ First, then PEEP/MAP

Spontaneous/Liberated

Transition to CPAP, extubate, or trach collar

Tracheostomy, speaking valves

Monitoring

Standard monitoring (Tele, SpO2, RR, NBP) plus the following:

Maximize knowledge

A-Line, CVP, EV1000, EtCO₂

Continuous SvO₂, Abdominal Pressures, [TTE/IVC measurement]

Assure effective Ventilation & Sedation (Standard ICU)

EtCO₂, A-Line, ±CVP

4E Compatible

Decrease NBP measurements overnight

Remove Foley

Nutrition

Goal: Minimize loss of lean mass

Holding Enteral Feed,

Consider TPN

Enteral Feeds at Trickle (20mL/hr)

Enteral feeds

Increasing to full support

Full Support

Enteral

Full Support

Enteral and PO including supplements

Full support

PO solids and supplements

Fluid Goal

Goal: Maintain organ perfusion; Avoid volume overload

Targeted Resuscitation/No over resuscitation

Give fluids only to achieve defined goal

Lactate decrease by 10% in 4 hrs

ScvO₂ > 70%, UOP > 0.5ml/kg/hr

Blood and Colloids to avoid over resuscitation

Targeted Management for Euvolemia

Assess intravascular volume status daily

Define Goal of Positive, Negative, or even

Fluid challenge

Diuretic challenge

Self-Management of Fluid Balance

No maintenance fluids

CRRT

(Check if NA ☐)

Consider High Dose Therapy

Regular Dose

Breaks for activities and tests

Consider IHD

Labs

Goal: Information availability & minimize blood loss

More Frequent

Q4-Q6: ABG, VBG/SvO₂, Lactate, CBC, Chem

Q12-24: LFT

Pedi Tubes, TEG, Coags, Fibrinogen

Less Frequent

Q12-Q24: ABG, CBC, Chem

Q24-Qweek: LFT, Coags

Weekly Nutrition Labs

As Needed/Intermittent

QOD-QMWF: CBC, Chem

Qweek: LFT, Coags

PRN Labs only

Access

Adequate access

Central line ≥ 3 ports

Arterial Line

Balance access and Infection

Remove arterial line

If on CRRT, Triple lumen dialysis catheter as only central access

Minimize Infection

Peripheral, fewer

Power Wand

PICC

Wound Care

Goal: Minimize wound infection, Suffering, & Heat loss

Complex dressing, NPWD placement/takedown > 40% or lg dressing with shower/staple removal

30-40% mod complex or shower type < 2 hours or no post op dressing change

Smaller or basic dressing or postop with wet downs, <20%

Simple/Self-care

Rehabilitation

Total Care

None or ROM/Positioning as tolerated

Other care is important

As much as tolerated

General Progression: ROM -> Sit/TLC -> Dangle/Tilt/Stand

Rehab likely more important

Minimal Assist

to March/Walk -> Gym & Outside!

Rehab most important, DO NOT DELAY

Medications

Goal: Minimize polypharmacy

More continuous

More IV

Continuous + PRN

More IV + Enteral

PRN + More Scheduled

Mixed IV + Enteral

Some PO

More PO

Some Enteral


Team View

Patient Severity of Illness, Status, and Goals

Severity of Illness *(transfer from condition assessment tool)*

Allergies

Code Status



Status

Place "X" or description where these occur and the days post procedure in the boxes above the blue line. <i>Circle planned events</i>	% Open																				
	OR																				
	Central Line																				
	Arterial Line																				
	Dialysis Cath																				
	Extubation																				
	Intubation																				
	BM																				
	Cultures																				
	Dressings																				
	"Down-Day"																				
	Rehab																				
	Completed																				
	Other Events (e.g. family meeting, etc.)																				

Recommendations/Messages

	Antibiotic(s)	NA ☐	Effective Start Date	Planned Stop Date	Diagnosis

Goals *(completed during rounds)*

Fluid volume goal for today:

☐ Positive _____ L

☐ Negative _____ L

☐ Even = + _____ L

The most important goal(s) for today (what we must do to be successful):

Priorities:

☐ Wound Care ☐ Rehab

☐ Procedure(s) ☐ Imaging

☐ SH/BT ☐ Extubation

☐ Other _____

Team View

Patient Severity of Illness, Status, and Goals

Severity of Illness (transfer from scales tool)

Allergies

Code Status

Date HospDay/ECMO Day	/1	/2	/3	/4	/5	/6	/7	/8	/9	/10	/11	/12	/13	/14	/15	/16	/17	/18	/19	/20	/21
Could Die Most Sick																					
Could Decannulate																					

Status

Flow/ RPM																				
Delta P																				
Circuit PaO2																				
Sweep																				
Patient PaO2																				
Patient PaCO2																				
Compliance																				
PRBC																				
FFP																				
PLT																				
Cryo																				
Amicar/TXA																				
Rehab																				
Other Events (e.g. Procedures)																				

Recommendations/Messages

Antibiotic(s)	NA ☐	Effective Start Date	Planned Stop Date	Diagnosis

Goals (completed during rounds)

Fluid volume goal for today:

☐ Positive _____ L

☐ Negative _____ L

☐ Even = + _____ L

The most important goal(s) for today (what we must do to be successful):

Priorities:

☐ Wound Care

☐ Rehab

☐ Procedure(s)

☐ Imaging

☐ Other _____

Page 1: Nursing Illness Severity Assessment

Bed #:

Time/Date:

Instructions: Make a noticeable “X” ANYWHERE on ANY of the scales below that indicates your estimate of the patient’s condition right now. Use the “average” of these marks to identify the Illness Severity (summary assessment) at the bottom. Transfer that summary assessment to the scale at the top of Page 2.

[illegible]

Diagnoses & Problems

Current or Chronic Problems: Bad <<< Getting Worse <<< Same >>> Getting Better >>> Baseline

Worse = Increasing in number or severity; **Better** = Decreasing in number or severity

Neuro/Pain

Sedation Goal

RASS -4 to -5

RASS -1 to -3

RASS 0 to -1

RASS 0

Paralyzed

Notes:

Respiratory				
Mechanical Ventilation	Low Tidal Volumes	PCV, VDR, or APRV	APRV or CPAP/PSV	Off-Vent
	PEEP>16 or MAP>28	PEEP>12 or MAP>18	Breathing Trials	Chronic vent
FiO2/Oxygenation	iNO		Weaning	
	Rotaprone			
Notes:	100% FiO2,	70-80% FiO2	40-60% FiO2	< 40% FiO2
	P:F < 100 or OI>35	P:F 100-200 or OI > 25	P:F >200 or OI > 14	P:F > 300 or OI < 15
	SpO2 < 90	SpO2>90	SpO2 > 90	SpO2 Normal or Baseline

Cardiac				
Vasopressors	Levophed > 25	Levophed < 25 Vasopressin 0.04	Levophed < 25 Vasopressin 0.04	No Vasopressors
Acidosis/Shock	pH < 7.2 Lactate > 6/increasing Trending ScvO2 often	pH 7.2-7.3 or > 7.5 Lactate decreasing or normal		pH 7.3-7.5 Not checking lactates
Notes:				

GI	Notes
----	-------

Renal/GU	
RRT	High volume CVVH
Goal UF:	CVVH
BUN/Cr:	None or IHD
Notes:	

Endo	Accuchecks:
	Notes:

Heme				
Blood & Fluids	Belmont in the room	Fluid loading	Goal Negative or Even	
Blood Products Given:	PRBC	FFP	PLTS	Cryp

Labs/ABG/pH		
Frequency	> Q6	Q6-Q24 ≤Q24
Electrolytes/Labs	K > 6 or ECG changes DIC with bleeding	DIC Normal
Notes:		

Skin/Wounds

Notes: Bad (IFI or Necrosis) Not Bad Good
Extensive (> 70% open) Medium Sized (20-70% open) Small (< 20% open)

Activity	Number of participants
None	1
ROM	10
Sitting/TLC	10
Stand/Tilt	10
Walking	5

<i>ID</i>	<i>Tmax:</i>	<i>WBC:</i>	<i>ABX:</i>	<i>Day/Reason:</i>
Notes:				

Risk of Worsening			
	High = Any of the Following WBC (> 15/rising or < 4/dropping) Hypothermic (< 36/<96.8) Worsening Mental Status Increasing HR or RR, or decreasing MAP	Medium Febrile Not tolerating rehabilitation Increasing Gastric Residuals Increasing Blood Glucose or Insulin Requirement	Low = All of the following Normal/Stable WBC Normal Temperature Normal Vital Signs Tolerating feeding Stable insulin requirement Tolerating rehabilitation

Severity of Illness

(summary assessment)

****Copy to the Next Page****

Most sick, Could Die

Least Sick Could transfer

If the patient's SOI changes by 2 or more blocks, page the on-call resident or attending to discuss.

Step 1

Step 1

Page 2: Assessment & Recommendations: Patient’s CURRENT Treatments

Step 3. Instructions:

- Mark current treatments.
- Note discrepancies between current treatments & the intended treatments according to patient condition.

Severity of Illness										
---------------------	--	--	--	--	--	--	--	--	--	--

Analgesia & Sedation	Intubated Patients Only		Intubated and Non-Intubated Patients	
	Deep Sedation (RASS -4 to -5) +/-Paralyzed Non-Responsive	Moderate Sedation (RASS -1 to -3) Arousable (opens eyes, withdraws)	Light Sedation (RASS 0 to -1) Interactive (follows commands)	No Sedation (RASS 0) Participatory (expresses self)
	☐ minimize oxygen demand	☐ patient ventilator synchrony	☐ prevent patient harm	☐ maximize patient engagement & participation in care
	☐ maximize perfusion	☐ prevent agitated movements	☐ engage patient in care	☐ maximize rehabilitation
	☐ protect grafts	☐ protect grafts, tube/line/device	☐ more physical/occupational therapy	
	☐ Continuous	☐ Continuous	☐ Scheduled IV/Enteral/PO	☐ Scheduled IV or
	+	+	+	Enteral or PO +
	☐ PRN	☐ PRN	☐ PRN IV or Enteral/PO	☐ PRN Enteral > IV
Notes/Recommendations:				

Delirium	Prevention	☐ Day-Night Light Cycle	☐ Day-Night Light Cycle ☐ Un-Interrupted Sleep, 4-6 hrs ☐ Physical activity ☐ Consider ear plugs	☐ Day-Night Light Cycle ☐ Sleep, 4-8 hrs ☐ Increase mobility ☐ Consider ear plugs, sleep aid
	Treatment	☐ NA	☐ Dexmedetomidine drip ☐ Haloperidol IV Push	☐ Haloperidol IV Push ☐ Quetiapine PO/Enteral
	Notes/Recommendations: Sleep aide dose: Restraints: Y/N Hours of Sleep:			

Monitoring/ Lines	Maximize knowledge Standard ICU Plus:	Assure effective Ventilation & Sedation Standard ICU Plus:	Ward Compatible ☐ Standard ICU
	Trend: ☐ Abdominal Pressures ☐ TTE/IVC size	☐ Continuous CO/SVV/SVR ☐ Foley ☐ EtCO2 ☐ A-Line ☐ ±CVP	Consider: ☐ Decrease NBP measurements overnight ☐ Remove Foley
	Lines: Notes/Recommendations:		

Mechanical Ventilation	☐ Paralysis	☐ VDR Protocol ☐ PCV Protocol ☐ ARDS Algorithm	☐ APRV Protocol ☐ Oscillatory/Demand CPAP protocol ☐ CPAP/PSV Protocol ☐ Daily breathing trials and/or wean	☐ CPAP/CPAP-PS Protocol ☐ Trach collar/speaking valve ☐ NA, not on mechanical ventilation
	Notes/Recommendations:			

Nutrition	☐ Holding Enteral Feeds ☐ Consider TPN	☐ Goal = full enteral support ☐ If unable to achieve, use TPN	☐ Goal is transition to PO solids and supplements
	Notes/Recommendations: Last BM: FMS: Y/N TF Goal:		
	Steroids: Y/N PPI: Y/N		

Labs	Q4-Q6:		Q8-Q24:	Q24-QOD:
	☐ ABG ☐ VBG/SvO2 ☐ Lactate ☐ CBC ☐ Chem	Consider: ☐ TEG ☐ Coags ☐ Fibrinogen Q12-24: ☐ LFT ☐ Drug Levels ☐ Pedi Tubes	☐ ABG ☐ CBC ☐ Chem ☐ LFT ☐ Coags Consider: ☐ Drug Levels	☐ CBC ☐ Chem ☐ Qweek: ☐ LFT ☐ Coags Consider: ☐ Drug Levels ☐ PRN Labs only
	Notes/Recommendations: DVT Prophylaxis: Y/N			

Rehabilitation	None or As Able ☐ None or ☐ ROM/Positioning Q2hrs ☐ Splinting	As Tolerated to Maintain or Improve Function ☐ General Progression: ROM → ☐ Bed-to-Chair Position/Cardiac Chair → ☐ Dangle/Tilt/Stand ☐ Splinting	Improve Function ☐ to March/Walk → ☐ Go Outside! ☐ Splinting
	Notes/Recommendations:		

If all checks are not aligned with the the patient’s SOI, discuss with the physicians.

Page 1. Burn Illness Severity Assessment

[illegible]

If the patient's SOI changes by 2 or more blocks, discuss during a "huddle."

Review QA checklist daily. Discuss problems on rounds.

	NA	Yes	No		NA	Yes	No
Appropriate GI Prophylaxis (PPI if > 20%TBSA)?	☐	☒	☐	Respiratory orders current?	☐	☒	☐
Appropriate DVT Prophylaxis (Medical if > 20%TBSA)?	☐	☒	☐	Wound care order current?	☐	☒	☐
Receiving appropriate oral care (CHG Q6 if intubated, BID teeth brushed if not)?	☐	☒	☐	Family has questions to discuss (ask nursing)?	☐		☐
HOB > 30 degrees (higher if indicated)?	☐	☒	☐	Delirium Positive?	☐		☐
Last BM < 48 hrs ago?	☐	☒	☐	Restraints needed?	☐		☐
Polypharmacy minimized (orders “cleaned”)?	☐	☒	☐	Current treatments match current condition (see next page)?		☒	☐

Page 2. Illness Severity Checklists

Step 3. Instructions: Mark **current** treatments in the white areas. Note discrepancies between current treatments & the intended treatments according to patient's severity of illness. Present information in **grey** on rounds.

One Liner:

24 hr events/rehab tolerance/Subjective:

Severity of Illness

Neuro	Analgesia & Sedation		Intubated Patients Only		Intubated and Non-Intubated Patients	
			Deep Sedation (RASS -4 to -5) +/-Paralyzed Non-Responsive ☐ minimize oxygen demand ☐ maximize perfusion ☐ protect grafts ☐ Continuous + PRN	Moderate Sedation (RASS -1 to -3) Arousable (opens eyes, withdraws) ☐ patient ventilator synchrony ☐ no agitated movements ☐ protect grafts, T/L/D ☐ Continuous + PRN	Light Sedation (RASS 0 to -1) Interactive (follows commands) ☐ prevent patient harm ☐ engage patient in care ☐ more physical/occupational therapy ☐ Scheduled IV/Enteral/PO + PRN IV or Enteral/PO	No Sedation (RASS 0) Participatory (expresses self) ☐ maximize patient engagement & participation in care ☐ maximize rehabilitation ☐ Sched. IV/Enteral/PO + PRN Enteral > IV
	Delirium	Prevention ☐ Day-Night Light Cycle	☐ Day-Night Light Cycle ☐ Un-Interrupted Sleep, 4-6 hrs ☐ Physical activity ☐ Consider ear plugs ☐ Dexmedetomidine drip ☐ Haloperidol IV Push			☐ Day-Night Light Cycle ☐ Sleep, 4-8 hrs ☐ Increase mobility ☐ Ear plugs, sleep aid ☐ Haloperidol IV Push ☐ Quetiapine PO/Enteral
		Treatment ☐ NA				
	Exam (Pupils/MS/Focality)_____		Current Narcotics/Dose last 24 hrs_____		Notes/Plan:	
	GCS:_____CAM ICU:_____Hrs of sleep:_____		Current Sedatives/Dose last 24 hrs_____			
	Imaging:_____		Other meds_____			
Pulm	Mechanical Ventilation	☐ Paralysis	☐ VDR Protocol ☐ PCV Protocol ☐ ARDS Protocol	☐ APRV Protocol ☐ Oscillatory/Demand CPAP protocol ☐ CPAP/PSV Protocol ☐ Daily breathing trial and/or wean	☐ CPAP/CPAP-PS Protocol ☐ Trach collar/speaking valve ☐ No Daily CXR	
	ECMO	Pt ABG_____	VDR: PIP_____PEEP_____OsPEEP_____MAP_____FIO2_____			
	LPM_____RPM_____	Pt SvO2_____	AC: Vt_____mL/kg_____f(set/tot)_____Peep_____FiO2_____Ppk_____Pplat_____			
	VenPres_____ArtPres_____ΔPress_____	Circuit PaO2_____	PC: PIP_____f(set/tot)_____Peep_____FiO2_____Vt_____mL/kg_____			
	SweepFlow_____FIO2_____		PS: PS_____PEEP_____f_____FiO2_____Vt_____mL/kg_____			
			APRV: PH_____PL_____TH_____TL_____FiO2_____Vt_____MVtot/spont_____			
	RR/Sats:_____Exam:_____	ABG_____P:F_____	OI[(MAPxFiO2)/PaO2]_____	Notes/Plan:		
CV	Monitoring	Maximize knowledge	Assure effective Ventilation & Sedation		Ward Compatible	
	Standard monitoring (Tele, SpO2, RR, NBP)	Standard ICU Plus: Trend: ☐ Abdominal Pressures ☐ TTE/IVC size	Standard ICU Plus: ☐ Foley ☐ EtCO2 ☐ A-Line ☐ ±CVP		Standard ICU Consider: ☐ Decrease NBP measurements overnight ☐ Remove Foley	
	HR_____BP_____MAP_____	CVP_____	CO/CI_____SVR_____	Lactate/SvO2: _____	Notes/Plan:	
	Exam_____	Pulses:_____	Meds:_____			
GI	Nutrition	☐ Holding Enteral Feeds ☐ Consider TPN	☐ Goal = full enteral support ☐ If unable to achieve, use TPN		☐ Goal is transition to PO solids and supplements	
	Exam:_____	LFTs/Amy/Lip_____	Notes/Plan:			
	Imaging:_____	PreAlb/UUN/Metabolic Cart:_____				
	Feeding_____	Residuals:_____Last BM_____	Meds:_____			
	Labs	Q4-Q6: ☐ ABG, SvO2, Lactate, CBC, Chem	Consider: ☐ TEG, Coags ☐ Fibrinogen	Q8-Q24: ☐ ABG, CBC, Chem, LFT, Coags Consider: ☐ Drug Levels	Q24-QOD: ☐ CBC, Chem Qweek: ☐ LFT, Coags Consider: ☐ PRN Labs only	
Renal	I_____O_____Net_____UOP/hr_____	CRRT: ☐ Prisma ☐ NxStage BFR_____RFR_____Solution_____UF_____				
	Est Insens loss (1cc/kg/tbsa)_____	Anticoagulation_____	Filter Life (hrs)_____			
	Weight_____	Dose (RFRxHrs on/24/kg)_____	FF_____			
	Drain OP: Gastric_____CT's_____JP's_____NPWD_____Other(s)_____	Notes/Plan:				
	Other Studies: 	C_____M_____P_____	IV Drips:_____			
		Meds:_____				
Endo	FSBGs_____Insulin Tot_____	Meds:_____		Notes/Plan:		
	Other (TSH,Cortisol, etc.)					
Heme		TEG-R_____α_____MA_____Ly30_____RBCs_____FFP_____PLT_____Cryo_____	Notes/Plan:			
		Others:_____	Other_____			
ID	Tmax/TCurrent Cultures	WBC/Diff ABX (Day#/of#) & Levels		Notes/Plan:		
	Rehabilitation	None or As Able ☐ None or ☐ ROM/Positioning Q2hrs ☐ Splinting	As Tolerated to Maintain or Improve Function ☐ General Progression: ROM → ☐ Bed-to-Chair Position/Cardiac Chair → ☐ Danlge/Tilt/Stand ☐ Splinting		Improve Function ☐ to March/Walk → ☐ Go Outside! ☐ Splinting	
Wnds	Exam: _____	Current % Open: _____	Current Wound Care: _____		Notes/Plan:	
	Last OR _____	Next OR: _____	Next Dressings Down Day: _____			
T/L/D	CVC Location/Day _____	ETT (size/depth) _____		Notes/Plan:		
	AL Location/Day _____	Foley(day) _____	FMS(day/last DRE) _____			
	NGT/DHT: _____	Other: _____				

Houston Site Tools


1. Original TeamView
2. Original Scales Tool

[illegible]

C. Team View

Patient Severity of Illness, Status, and Goals

Severity of Illness (transfer from scales tool)



Status

% Open							
% Feeding Achieved							
Biggest Issue							
Coordinating Activities See code list behind Scales Tool for codes.							

Update Checklist items daily!

	NA	Good	Discuss
GI Prophylaxis	☐	☐	☐
CHG Prophylaxis	☐	☐	☐
HOB > 30 degrees	☐	☐	☐
DVT Prophylaxis	☐	☐	☐
Last BM < 48 hrs ago	☐	☐	☐
Respiratory Orders Current	☐	☐	☐
Family has questions	☐		☐
Delirium Positive	☐		☐
Pending Cultures	☐		☐

Antibiotic(s)	NA ☐	Effective Start Date	Planned Stop Date	Diagnosis

Goals (completed during rounds)

Fluid volume goal for today:

- ☐ Positive
- ☐ Negative
- ☐ Even

Major Goal(s) for next 24-48 hours (*Star Primary Goal*)

Priorities of care to achieve these goals

Dallas Site Tools

1. Original Team View
2. Original Scales/Checklist Tool

Team View

Patient Severity of Illness, Status, and Goals

Severity of Illness *(transfer from scales tool)*

Allergies


Code Status

[illegible]

Status

[illegible]

Major Problem List

[illegible]

Draw Current Wounds

Indicate Wound Care

Antibiotic(s)

NA ☐

Start Date/Stop Date

Diagnosis

Recommendations/Messages

Goals (completed during rounds)

Fluid volume goal for today:

☐ Positive L☐ Negative L

☐ Even = + L

The most important goal(s) for today (what we must do to be successful):

Priorities:

___ Wound Care ___ Rehab

___ Procedure(s) ___ Imaging

Other _____

[illegible]

Step 3. For patients with a Green-Yellow Illness Severity. Review each question. Items that are marked No/Discuss should be discussed with a physician as soon as possible (at least during morning or evening rounds).

		Yes/NA	No/Discuss
Analgesia & Sedation	Is the patient interactive/participatory in their care?	☐	☐
	Is the patient awake and participatory in their care? If not, how might you get them so?	☐	☐
	Does the patient have minimal pain?	☐	☐
	Is pain controlled per the patient?	☐	☐
	Is the patient primarily on PO/Enteral pain meds with supplemental IV PRN meds?	☐	☐
	Is the patient on NO sedation?	☐	☐
Mechanical Ventilation	Is the patient off the ventilator?	☐	☐
	If not off the ventilator, does the patient need a tracheostomy?	☐	☐
	If the patient is not off the ventilator, did he/she receive a breathing trial today?	☐	☐
	If not off the ventilator, is the patient on CPAP and/or did he/she receive a breathing trial or trach collar trial?	☐	☐
	What O2 Concentration/Support if applicable?	☐	☐
Labs	Is the frequency of all labs daily or none?	☐	☐
Monitoring	Is the patient on no more than standard ICU monitoring (Telemetry, SpO2, Foley)?	☐	☐
	Can you reduce NBP measurements overnight?	☐	☐
	Can you remove the Foley?	☐	☐
Sleep	Is the patient on schedule to maintain circadian rhythm (Day/Night Cycle)?	☐	☐
	Does he/she need a Sleep Aid?	☐	☐
	How long did this patient sleep?	☐	☐
Rehabilitation	Is the patient able to sit and/or stand at the bedside?	☐	☐
	Can you advance the patient's rehab goal to marching, walking, and possibly going to the gym or outside?	☐	☐
Nutrition	If not on a PO diet, is he/she on full enteral feeds?	☐	☐
	What is the diet?	☐	☐
Medications	Is the patient on PO meds only and possibly some enteral or IV PRN meds?	☐	☐
Transition	Is this patient ready to transfer to the ward?	☐	☐
	What needs to be done for the patient in order to transfer him/her to the ward?	☐	☐

	NA	Good	Discuss		NA	Good	Discuss
GI Prophylaxis	☐	☐	☐	Respiratory orders current	☐	☐	☐
DVT Prophylaxis	☐	☐	☐	Wound care order current	☐	☐	☐
CHG Prophylaxis	☐	☐	☐	Family has questions	☐		☐
HOB > 30 degrees	☐	☐	☐	Delirium Positive	☐		☐
Last BM < 48 hrs ago	☐	☐	☐	Restraints	☐		☐

Step 3. For patients with a Yellow-Orange Illness Severity. Review each question. Items that are marked No/Discuss should be discussed with a physician as soon as possible (at least during morning or evening rounds).

		Yes/NA	No/Discuss
Analgesia & Sedation	Has the goal of sedation been defined? Arousable or more awake?	☐	☐
	If appropriate, limit sedation to promote ventilator weaning.	☐	☐
	Use APRV, CPAP, or normal tidal volume ventilation if decreasing ventilator support.	☐	☐
	Use volume or pressure control ventilation if the patient is not weaning or is getting worse.	☐	☐
	Consider a daily breathing trial.	☐	☐
	Did the patient receive a daily breathing trial or is one scheduled for today?	☐	☐
Labs	Daily labs.	☐	☐
	Is the patient on the correct/best lab frequency? (Q12-24 ABG, CBC, Chem; Q24-QWeek LFT, Coag)	☐	☐
Monitoring	Is the patient on at least standard ICU monitoring only (Telemetry, SpO2, Foley)?	☐	☐
	Routine ICU Monitoring (A-Line, EtCO2, Telemetry, SpO2, RR, NBP, +/- CVP)	☐	☐
Sleep	Is there a schedule to maintain day/night cycles?	☐	☐
	How long did the patient sleep last night?	☐	☐
Rehabilitation	Is the patient “tilting” or going to cardiac chair?	☐	☐
	<i>Consider sitting at bedside.</i>	☐	☐
	<i>Consider standing/marching/or walking if able.</i>	☐	☐
	<i>Consider standing or walking if able.</i>	☐	☐
Nutrition	Is the patient achieving “full support” (100% of caloric and protein goals) via enteral route?	☐	☐
	If unable to achieve “full support”, <i>consider TPN.</i>	☐	☐
Medications	Is the patient receiving mostly IV or Enteral plus IV PRN medications?	☐	☐

	NA	Good	Discuss		NA	Good	Discuss
GI Prophylaxis	☐	☐	☐	Respiratory orders current	☐	☐	☐
DVT Prophylaxis	☐	☐	☐	Wound care order current	☐	☐	☐
CHG Prophylaxis	☐	☐	☐	Family has questions	☐		☐
HOB > 30 degrees	☐	☐	☐	Delirium Positive	☐		☐
Last BM < 48 hrs ago	☐	☐	☐	Restraints	☐		☐

Step 3. For patients with a Orange-Red Illness Severity. Review each question. Items that are marked No/Discuss should be discussed with a physician as soon as possible (at least during morning or evening rounds).

		Yes/NA	No/Discuss
Analgesia & Sedation	Is the patient on mostly continuous IV analgesia and sedation with additional IV PRN medication?	☐	☐
	Is the patient on enough sedation/analgesia to achieve ventilation/oxygenation goals?	☐	☐
	Is the patient on AC Pressure Control with low tidal volumes or the VDR?	☐	☐
Labs	Is the patient on the correct/best lab frequency? (4-6 ABG, VBG/ScvO2, Lactate, CBC, Chem, Coags; Q12-24 LFT)	☐	☐
Monitoring	The patient should have continuous arterial blood pressure monitoring and continuous cardiac output monitoring.	☐	☐
	Goal is to maximize knowledge: Continuous CO, Arterial Line; Consider abdominal pressures, ScvO2 monitoring, Echocardiography, IVC ultrasound measurement	☐	☐
Sleep	<i>Consider day/night cycling if able.</i>	☐	☐
Rehabilitation	Is the patient being positioned and receiving ROM therapy at least every 4 hours?	☐	☐
	Do not prioritize rehabilitation; prioritize other care first.	☐	☐
Nutrition	If shock and holding enteral feeds, consider TPN.	☐	☐
Medications	This patient should be on continuous, titratable IV medications.	☐	☐
	<i>Consider stopping/holding enteral medications.</i>	☐	☐

	NA	Good	Discuss		NA	Good	Discuss
GI Prophylaxis	☐	☐	☐	Respiratory orders current	☐	☐	☐
DVT Prophylaxis	☐	☐	☐	Wound care order current	☐	☐	☐
CHG Prophylaxis	☐	☐	☐	Family has questions	☐		☐
HOB > 30 degrees	☐	☐	☐	Restraints	☐		☐
Last BM < 48 hrs ago	☐	☐	☐				