

TM No. 931133

**REFERENCE COPY**


NAVAL UNDERSEA WARFARE CENTER - DETACHMENT NEW LONDON  
NEW LONDON, CONNECTICUT 06320

Technical Memorandum

STATISTICAL THEOREMS INVOLVING THE COMPLEX  
MULTIVARIATE GAUSSIAN AND WISHART DENSITIES

Date: 13 September 1993

Prepared by:



Douglas A. Abraham

Advanced Engineering Division

Submarine Sonar Department

Approved for public release; distribution is unlimited.

## Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>13 SEP 1993</b>	2. REPORT TYPE <b>Technical Memo</b>	3. DATES COVERED <b>13-09-1993 to 13-09-1993</b>	
4. TITLE AND SUBTITLE <b>Statistical Theorems Involving the Complex Multivariate Gaussian and Wishart Densities</b>		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER <b>0602314N</b>	
6. AUTHOR(S) <b>Douglas Abraham</b>		5d. PROJECT NUMBER <b>RJ14C33</b>	
		5e. TASK NUMBER <b>3</b>	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Undersea Warfare Center Division, Newport, RI, 02841</b>		8. PERFORMING ORGANIZATION REPORT NUMBER <b>TM 931133</b>	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) <b>Office of Naval Research</b>		10. SPONSOR/MONITOR'S ACRONYM(S) <b>ONR 451</b>	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>			
13. SUPPLEMENTARY NOTES <b>NUWC2015</b>			
14. ABSTRACT <b>The complex multivariate Gaussian and Wishart densities are often encountered in the analysis of statistical signal processing algorithms in frequency domain array processing. The derivation of the probability density function of forms involving complex Gaussian random vectors and complex Wishart matrices may be simplified by the use of standard theorems found in multivariate statistics for real Gaussian random vectors and real Wishart matrices. This memorandum contains the extensions to the complex case of a selection of these theorems.</b>			
15. SUBJECT TERMS <b>Gaussian; Wishart; Transient Signal Processing Project; TSP</b>			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	
			18. NUMBER OF PAGES <b>16</b>
			19a. NAME OF RESPONSIBLE PERSON

## Abstract

The *complex* multivariate Gaussian and Wishart densities are often encountered in the analysis of statistical signal processing algorithms in frequency domain array processing. The derivation of the probability density function of forms involving complex Gaussian random vectors and complex Wishart matrices may be simplified by the use of standard theorems found in multivariate statistics for *real* Gaussian random vectors and *real* Wishart matrices. This memorandum contains the extensions to the complex case of a selection of these theorems.

## ADMINISTRATIVE INFORMATION

The work presented in this memorandum was performed under Task 3 of the Transient Signal Processing (TSP) Project as part of the Submarine/Surface Ship USW Surveillance Program sponsored by the Technology Directorate of the Office of Naval Research: Program Element 0602314N, ONR Program UN3B, Project Number RJ14C33, NUWC Job Order No. A60070, Principal Investigator M. W. Gouzie (Code 2121), Program Director, G. C. Connolly (Code 2192). The sponsoring activity's Director for Undersea Surveillance is T. G. Goldsberry (ONR 451).

The author of this memorandum is located at the Naval Undersea Warfare Center, New London Detachment, New London, CT 06320. The technical reviewer for this memorandum was T. Luginbuhl (Code 2121).

The author thanks Tod Luginbuhl for his careful review of this document.

STATISTICAL THEOREMS INVOLVING THE COMPLEX MULTIVARIATE  
GAUSSIAN AND WISHART DENSITIES

INTRODUCTION

The theorems found in this memorandum are extensions to the complex case of theorems commonly found in the analysis of multivariate real Gaussian random variables. The original theorems may be found in either Searle [1] or Muirhead [2]. The extensions are, for the most part, straightforward and result from following the proofs for the real cases. When possible, the proofs have been simplified or deferred. Background and distributional information on multivariate complex Gaussian random variables and complex Wishart matrices may be found in Goodman [3] and to a lesser extent in Anderson [4].

NOTATION

In this memorandum, boldface lower case letters represent vectors, boldface upper case letters represent matrices, and the superscript  $H$  represents the conjugate transpose operation. If the  $n$  dimensional random vector  $\mathbf{x}$  has a complex Gaussian distribution, the notation

$$\mathbf{x} \sim \mathcal{CN}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \quad (1)$$

is used where  $\boldsymbol{\mu}$  is the mean vector and  $\boldsymbol{\Sigma}$  is the covariance matrix. The probability density function of  $\mathbf{x}$  is

$$f(\mathbf{x}) = \frac{1}{\pi^n |\boldsymbol{\Sigma}|} e^{-(\mathbf{x}-\boldsymbol{\mu})^H \boldsymbol{\Sigma}^{-1} (\mathbf{x}-\boldsymbol{\mu})}. \quad (2)$$

If the  $n$ -by- $n$  random matrix  $\mathbf{A}$  has a complex Wishart distribution, the notation

$$\mathbf{A} \sim \mathcal{CW}_n(k, \boldsymbol{\Sigma}) \quad (3)$$

is used where  $k$  is the number of degrees of freedom and  $\boldsymbol{\Sigma}$  is the scale matrix. The probability density function of the complex Wishart distribution is described in Theorem 3. If the scalar random variable  $Z$  has a non-central chi-squared distribution, the notation

$$Z \sim \mathcal{X}_P^2(\delta) \quad (4)$$

is used, where  $P$  is the number of degrees of freedom and  $\delta$  is the non-centrality parameter. The probability density function of such a non-central chi-squared random variable is, as found in Johnson and Kotz [5],

$$f(z) = \sum_{k=0}^{\infty} \frac{e^{-\frac{\delta}{2}} \left(\frac{\delta}{2}\right)^k}{k!} \frac{z^{\frac{P+2k}{2}-1} e^{-\frac{z}{2}}}{2^{\frac{P+2k}{2}} \Gamma\left(\frac{P+2k}{2}\right)}. \quad (5)$$

It should be noted that Searle [1] defines the non-central chi-squared probability density function as having a non-centrality parameter equal to  $\frac{\delta}{2}$  in equation (5).

### THEOREM 1

The following theorem is an extension of Theorem 2, Chapter 2, page 57 of Searle [1] describing the probability density function of certain quadratic forms involving complex Gaussian random vectors.

**Theorem 1** *If  $\mathbf{x} \sim \mathcal{CN}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  and  $\mathbf{A} = \mathbf{A}^H$ , then  $2\mathbf{x}^H \mathbf{A} \mathbf{x} \sim \mathcal{X}_r^2(\delta)$  if and only if  $\mathbf{A}\boldsymbol{\Sigma}$  is idempotent. The non-centrality parameter,  $\delta = 2\boldsymbol{\mu}^H \mathbf{A} \boldsymbol{\mu}$ , and the degrees of freedom parameter,  $r = 2\text{tr}(\mathbf{A}\boldsymbol{\Sigma})$ , where  $\text{tr}(\mathbf{A}\boldsymbol{\Sigma})$  is the trace of  $\mathbf{A}\boldsymbol{\Sigma}$ .*

*Proof:* For the quadratic form

$$Q = 2\mathbf{x}^H \mathbf{A} \mathbf{x}, \quad (6)$$

consider the moment generating function

$$\begin{aligned} M_Q(t) &= \text{E} [e^{tQ}] \\ &= \text{E} [e^{2t\mathbf{x}^H \mathbf{A} \mathbf{x}}] \\ &= \int_{\mathcal{C}^n} \frac{1}{\pi^n |\boldsymbol{\Sigma}|} \exp \left[ 2t\mathbf{x}^H \mathbf{A} \mathbf{x} - (\mathbf{x} - \boldsymbol{\mu})^H \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right] d\mathbf{x} \\ &= \int_{\mathcal{C}^n} \frac{1}{\pi^n |\boldsymbol{\Sigma}|} e^V d\mathbf{x}, \end{aligned} \quad (7)$$

where  $\mathcal{C}^n$  represents  $n$  dimensional complex Euclidean space. The exponent,  $V$ , may be massaged into a single quadratic form in  $\mathbf{x}$ ,

$$V = \mathbf{x}^H (2t\mathbf{A}) \mathbf{x} - \mathbf{x}^H \boldsymbol{\Sigma}^{-1} \mathbf{x} + \boldsymbol{\mu}^H \boldsymbol{\Sigma}^{-1} \mathbf{x} + \mathbf{x}^H \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu} - \boldsymbol{\mu}^H \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}$$

$$= -(\mathbf{x} - \tilde{\boldsymbol{\mu}})^H \tilde{\boldsymbol{\Sigma}}^{-1} (\mathbf{x} - \tilde{\boldsymbol{\mu}}) + \tilde{\boldsymbol{\mu}}^H \tilde{\boldsymbol{\Sigma}}^{-1} \tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}^H \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}, \quad (8)$$

where

$$\tilde{\boldsymbol{\Sigma}}^{-1} = \boldsymbol{\Sigma}^{-1} - 2t\mathbf{A} \quad (9)$$

and

$$\tilde{\boldsymbol{\mu}} = \tilde{\boldsymbol{\Sigma}} \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}. \quad (10)$$

The moment generating function thus becomes,

$$\begin{aligned} M_Q(t) &= \frac{\exp\left[\tilde{\boldsymbol{\mu}}^H \tilde{\boldsymbol{\Sigma}}^{-1} \tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}^H \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}\right]}{|\tilde{\boldsymbol{\Sigma}}|^{-1} |\boldsymbol{\Sigma}|} \int_{\mathcal{C}^n} \frac{\exp\left[-(\mathbf{x} - \tilde{\boldsymbol{\mu}})^H \tilde{\boldsymbol{\Sigma}}^{-1} (\mathbf{x} - \tilde{\boldsymbol{\mu}})\right]}{\pi^n |\tilde{\boldsymbol{\Sigma}}|} d\mathbf{x} \\ &= \frac{\exp\left[\boldsymbol{\mu}^H \left(\boldsymbol{\Sigma}^{-1} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\Sigma}^{-1} - \boldsymbol{\Sigma}^{-1}\right) \boldsymbol{\mu}\right]}{|\tilde{\boldsymbol{\Sigma}}^{-1} \boldsymbol{\Sigma}|} \\ &= \frac{\exp\left[-\boldsymbol{\mu}^H \left(\mathbf{I}_n - \boldsymbol{\Sigma}^{-1} (\boldsymbol{\Sigma}^{-1} - 2t\mathbf{A})^{-1}\right) \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}\right]}{|\mathbf{I}_n - 2t\mathbf{A}\boldsymbol{\Sigma}|} \\ &= \frac{\exp\left[-\boldsymbol{\mu}^H \left(\mathbf{I}_n - (\mathbf{I}_n - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1}\right) \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}\right]}{|\mathbf{I}_n - 2t\mathbf{A}\boldsymbol{\Sigma}|}, \end{aligned} \quad (11)$$

where  $\mathbf{I}_n$  is the  $n$  dimensional identity matrix.

Suppose that  $\mathbf{A}\boldsymbol{\Sigma}$  is idempotent with rank or trace,

$$\text{tr}(\mathbf{A}\boldsymbol{\Sigma}) = m. \quad (12)$$

Note that the eigenvalues of an idempotent matrix are either zero or one, and, by using a singular value decomposition, the matrix  $\mathbf{A}\boldsymbol{\Sigma}$  may be expressed as

$$\mathbf{A}\boldsymbol{\Sigma} = \mathbf{U}\mathbf{V}^H, \quad (13)$$

where the  $n$ -by- $m$  matrices  $\mathbf{U}$  and  $\mathbf{V}$  are orthogonal,

$$\mathbf{V}^H \mathbf{U} = \mathbf{I}_m. \quad (14)$$

Part of the matrix in the quadratic form in the exponent of equation (11) may now be simplified,

$$\mathbf{I}_n - (\mathbf{I}_n - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1} = \mathbf{I}_n - \left(\mathbf{I}_n - \mathbf{U}(2t)\mathbf{V}^H\right)^{-1}$$

$$\begin{aligned}
&= \mathbf{I}_n - \left[ \mathbf{I}_n - \mathbf{U} \left( \mathbf{V}^H \mathbf{U} - \frac{1}{2t} \mathbf{I}_m \right)^{-1} \mathbf{V}^H \right] \\
&= \mathbf{U} \left( 1 - \frac{1}{2t} \right)^{-1} \mathbf{V}^H \\
&= \left( 1 - \frac{1}{1-2t} \right) \mathbf{A} \boldsymbol{\Sigma}.
\end{aligned} \tag{15}$$

The determinant in the denominator of equation (11) is clearly

$$|\mathbf{I}_n - 2t \mathbf{A} \boldsymbol{\Sigma}| = (1 - 2t)^m \tag{16}$$

because  $\mathbf{A} \boldsymbol{\Sigma}$  has  $m$  non-zero eigenvalues, all equal to one. Substituting (15) and (16) into (11) results in

$$\begin{aligned}
M_Q(t) &= \frac{\exp \left[ - \left( 1 - \frac{1}{1-2t} \right) \mu^H \mathbf{A} \mu \right]}{(1 - 2t)^m} \\
&= e^{-\mu^H \mathbf{A} \mu} \frac{\exp \left( \frac{1}{1-2t} \mu^H \mathbf{A} \mu \right)}{(1 - 2t)^{\frac{2m}{2}}}.
\end{aligned} \tag{17}$$

The moment generating function of a non-central Chi-squared random variable with  $r$  degrees freedom and non-centrality parameter  $\delta$ , as found in Muirhead [2], is

$$M_{\chi_{r,\delta}^2}(t) = \frac{e^{-\frac{\delta}{2}} e^{\frac{\delta}{2(1-2t)}}}{(1 - 2t)^{\frac{r}{2}}}. \tag{18}$$

Equating equations (17) and (18), it is seen that the quadratic form,  $Q$ , has a non-central Chi-squared distribution with  $r = 2m = 2\text{tr}(\mathbf{A} \boldsymbol{\Sigma})$  degrees of freedom and non-centrality parameter  $\delta = 2\mu^H \mathbf{A} \mu$ .

If it is assumed that the quadratic form has the described non-central Chi-squared distribution, it can be seen that the matrix  $\mathbf{A} \boldsymbol{\Sigma}$  must be idempotent with rank equal to  $\text{tr}(\mathbf{A} \boldsymbol{\Sigma}) = \frac{r}{2}$  by equating the denominators of equations (11) and (18),

$$\begin{aligned}
(1 - 2t)^{\frac{r}{2}} &= |\mathbf{I}_n - 2t \mathbf{A} \boldsymbol{\Sigma}| \\
&= \prod_{i=1}^n (1 - 2t \lambda_i),
\end{aligned} \tag{19}$$

where  $\lambda_i$  are the eigenvalues of  $\mathbf{A} \boldsymbol{\Sigma}$ . Clearly the left and right sides of equation (19) will be equal only when  $\frac{r}{2}$  of the eigenvalues are equal to one with the remaining equal to zero, which results in an idempotent  $\mathbf{A} \boldsymbol{\Sigma}$  matrix.

## THEOREM 2

The following theorem is the extension of Theorem 4, Chapter 2, page 59 of Searle [1] showing the independence of quadratic forms involving complex Gaussian random vectors.

**Theorem 2** *If  $\mathbf{x} \sim \mathcal{CN}_n(\mu, \Sigma)$ ,  $\mathbf{A} = \mathbf{A}^H$  and  $\mathbf{B} = \mathbf{B}^H$ , then  $\mathbf{x}^H \mathbf{A} \mathbf{x}$  and  $\mathbf{x}^H \mathbf{B} \mathbf{x}$  are independent if and only if  $\mathbf{A} \Sigma \mathbf{B} = \mathbf{B} \Sigma \mathbf{A} = \mathbf{0}$ .*

*Proof:* The proof is identical to that of Searle [1] and, as the result is also identical, will not be repeated.

## THEOREM 3

The following theorem is an extension of Theorem 3.2.10 on page 93 of Muirhead [2] describing the probability density functions of the partitions of a complex Wishart matrix and of a particular function of the partitions that enjoys certain independence properties.

**Theorem 3** *If the  $n$ -by- $n$  matrix  $\mathbf{A} \sim \mathcal{CW}_n(m, \Sigma)$  with  $\mathbf{A}$  and  $\Sigma$  partitioned into two-by-two blocks,*

$$\begin{aligned} \mathbf{A} &= \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} \\ \Sigma &= \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}, \end{aligned} \tag{20}$$

where  $\mathbf{A}_{11}$  and  $\Sigma_{11}$  are  $k$ -by- $k$ , and if

$$\mathbf{A}_{11.2} = \mathbf{A}_{11} - \mathbf{A}_{12} \mathbf{A}_{22}^{-1} \mathbf{A}_{21} \tag{21}$$

and

$$\Sigma_{11.2} = \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21}, \tag{22}$$

then

1.  $\mathbf{A}_{11.2} \sim \mathcal{CW}_k(m - n + k, \Sigma_{11.2})$  and is independent of  $\mathbf{A}_{12}$  and  $\mathbf{A}_{22}$ .

2. The conditional distribution of  $\mathbf{A}_{12}$  given  $\mathbf{A}_{22}$  is

$$\mathcal{CN}_{k \times (n-k)}(\boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} \mathbf{A}_{22}, \boldsymbol{\Sigma}_{11 \cdot 2} \otimes \mathbf{A}_{22}). \quad (23)$$

3.  $\mathbf{A}_{22} \sim \mathcal{CW}_{n-k}(m, \boldsymbol{\Sigma}_{22})$ .

*Proof:* Consider the probability density function of the complex Wishart distributed matrix,  $\mathbf{A}$ , as found in Goodman [3],

$$f(\mathbf{A}) = \frac{|\mathbf{A}|^{m-n}}{\Gamma_{m,n} |\boldsymbol{\Sigma}|^m} \text{etr}(-\boldsymbol{\Sigma}^{-1} \mathbf{A}), \quad (24)$$

where ‘etr’ signifies the matrix exponential trace operation and

$$\Gamma_{m,n} = \pi^{\frac{n(n-1)}{2}} \Gamma(m) \cdots \Gamma(m-n+1) \quad (25)$$

where  $\Gamma(x)$  is the standard Gamma function. Applying the  $k, n-k$  factorization to the determinants of equation (24) yields

$$\begin{aligned} |\mathbf{A}| &= |\mathbf{A}_{22}| |\mathbf{A}_{11} - \mathbf{A}_{12} \mathbf{A}_{22}^{-1} \mathbf{A}_{21}| \\ &= |\mathbf{A}_{22}| |\mathbf{A}_{11 \cdot 2}| \end{aligned} \quad (26)$$

and, similarly,

$$|\boldsymbol{\Sigma}| = |\boldsymbol{\Sigma}_{22}| |\boldsymbol{\Sigma}_{11 \cdot 2}|. \quad (27)$$

The determinant of a partitioned matrix may be found in the Matrix Theory Appendix of Muirhead [2]. The trace of the matrix product  $\boldsymbol{\Sigma}^{-1} \mathbf{A}$  in equation (24) must now be massaged into a form containing the matrix partitions  $\mathbf{A}_{12}$  and  $\mathbf{A}_{22}$  and the matrix  $\mathbf{A}_{11 \cdot 2}$ . This requires substantial algebraic manipulation, beginning with the matrix factorization of the inverse of the matrix  $\boldsymbol{\Sigma}$ ,

$$\boldsymbol{\Sigma}^{-1} = \mathbf{V} = \begin{bmatrix} \mathbf{V}_{11} & \mathbf{V}_{12} \\ \mathbf{V}_{21} & \mathbf{V}_{22} \end{bmatrix}. \quad (28)$$

The following relationships between the partitioned matrix and the partitioned inverse may be easily verified:

$$\mathbf{V}_{11} = \boldsymbol{\Sigma}_{11 \cdot 2}^{-1} \quad (29)$$

$$\Sigma_{22}^{-1} = \mathbf{V}_{22} - \mathbf{V}_{21}\mathbf{V}_{11}^{-1}\mathbf{V}_{12} \quad (30)$$

$$\mathbf{V}_{11}^{-1}\mathbf{V}_{12} = -\Sigma_{12}\Sigma_{22}^{-1}. \quad (31)$$

Applying these to the aforementioned trace results in

$$\begin{aligned} \text{tr}(\Sigma^{-1}\mathbf{A}) &= \text{tr} \left( \begin{bmatrix} \mathbf{V}_{11} & \mathbf{V}_{12} \\ \mathbf{V}_{21} & \mathbf{V}_{22} \end{bmatrix} \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} \right) \\ &= \text{tr} \left( \begin{bmatrix} \mathbf{V}_{11}\mathbf{A}_{11} + \mathbf{V}_{12}\mathbf{A}_{21} & \mathbf{V}_{11}\mathbf{A}_{12} + \mathbf{V}_{12}\mathbf{A}_{22} \\ \mathbf{V}_{21}\mathbf{A}_{11} + \mathbf{V}_{22}\mathbf{A}_{21} & \mathbf{V}_{21}\mathbf{A}_{12} + \mathbf{V}_{22}\mathbf{A}_{22} \end{bmatrix} \right) \\ &= \text{tr}(\mathbf{V}_{11}\mathbf{A}_{11}) + \text{tr}(\mathbf{V}_{12}\mathbf{A}_{21}) + \text{tr}(\mathbf{V}_{21}\mathbf{A}_{12}) + \text{tr}(\mathbf{V}_{22}\mathbf{A}_{22}) \\ &= \text{tr}(\mathbf{V}_{11}\mathbf{A}_{11}) + 2\text{tr}(\mathbf{V}_{12}\mathbf{A}_{21}) + \text{tr}(\mathbf{V}_{22}\mathbf{A}_{22}) \\ &= \text{tr}(\mathbf{V}_{11}[\mathbf{A}_{11.2} + \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}]) + 2\text{tr}(\mathbf{V}_{12}\mathbf{A}_{21}) + \text{tr}(\mathbf{V}_{22}\mathbf{A}_{22}) \\ &= \text{tr}(\mathbf{V}_{11}\mathbf{A}_{11.2}) + \text{tr}(\mathbf{V}_{11}\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}) + 2\text{tr}(\mathbf{V}_{12}\mathbf{A}_{21}) + \text{tr}(\mathbf{V}_{22}\mathbf{A}_{22}) \\ &= \left\{ \begin{array}{l} \text{tr}(\Sigma_{11.2}^{-1}\mathbf{A}_{11.2}) + \text{tr}(\mathbf{V}_{11}\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}) \\ -2\text{tr}(\mathbf{V}_{11}\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{21}) + \text{tr}([\Sigma_{22}^{-1} + \mathbf{V}_{21}\mathbf{V}_{11}^{-1}\mathbf{V}_{12}]\mathbf{A}_{22}) \end{array} \right\} \\ &= \left\{ \begin{array}{l} \text{tr}(\Sigma_{11.2}^{-1}\mathbf{A}_{11.2}) + \text{tr}(\mathbf{V}_{11}\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}) - 2\text{tr}(\mathbf{V}_{11}\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{21}) \\ +\text{tr}(\Sigma_{22}^{-1}\mathbf{A}_{22}) + \text{tr}(\mathbf{A}_{22}\Sigma_{22}^{-1}\Sigma_{21}\mathbf{V}_{11}\Sigma_{12}\Sigma_{22}^{-1}) \end{array} \right\} \\ &= \left\{ \begin{array}{l} \text{tr}(\Sigma_{11.2}^{-1}\mathbf{A}_{11.2}) + \text{tr}(\mathbf{V}_{11}\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}) \\ -\text{tr}(\mathbf{V}_{11}\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{21}) - \text{tr}(\mathbf{V}_{11}\mathbf{A}_{12}\Sigma_{22}^{-1}\Sigma_{21}) \\ +\text{tr}(\Sigma_{22}^{-1}\mathbf{A}_{22}) + \text{tr}(\mathbf{V}_{11}\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{22}\Sigma_{22}^{-1}\Sigma_{21}) \end{array} \right\} \\ &= \left\{ \begin{array}{l} \text{tr}(\Sigma_{11.2}^{-1}\mathbf{A}_{11.2}) + \text{tr}(\Sigma_{22}^{-1}\mathbf{A}_{22}) \\ +\text{tr} \left( \mathbf{V}_{11} \begin{bmatrix} \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21} - \Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{22}\mathbf{A}_{22}^{-1}\mathbf{A}_{21} \\ -\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{22}\Sigma_{22}^{-1}\Sigma_{21} \\ +\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{22}\mathbf{A}_{22}^{-1}\mathbf{A}_{22}\Sigma_{22}^{-1}\Sigma_{21} \end{bmatrix} \right) \end{array} \right\} \\ &= \left\{ \begin{array}{l} \text{tr}(\Sigma_{11.2}^{-1}\mathbf{A}_{11.2}) + \text{tr}(\Sigma_{22}^{-1}\mathbf{A}_{22}) \\ +\text{tr}(\Sigma_{11.2}^{-1}[\Sigma_{12}\Sigma_{22}^{-1}\mathbf{A}_{22} - \mathbf{A}_{12}]\mathbf{A}_{22}^{-1}[\mathbf{A}_{22}\Sigma_{22}^{-1}\Sigma_{21} - \mathbf{A}_{21}]) \end{array} \right\} \end{aligned}$$

$$= \left\{ \begin{array}{c} \text{tr}(\Sigma_{11.2}^{-1} \mathbf{A}_{11.2}) + \text{tr}(\Sigma_{22}^{-1} \mathbf{A}_{22}) \\ + \text{tr}(\Sigma_{11.2}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}] \mathbf{A}_{22}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}]^H) \end{array} \right\} \quad (32)$$

Substituting equations (26), (27), and (32) into the probability density function of equation (24) yields

$$\begin{aligned} f(\mathbf{A}) &= \left\{ \begin{array}{c} \left[ \frac{|\mathbf{A}_{11.2}|^{m-n}}{\Gamma_{m,n} |\Sigma_{11.2}|^m} \text{etr}(-\Sigma_{11.2}^{-1} \mathbf{A}_{11.2}) \right] \left[ \frac{|\mathbf{A}_{22}|^{m-n}}{|\Sigma_{22}|^m} \text{etr}(-\Sigma_{22}^{-1} \mathbf{A}_{22}) \right] \\ \cdot \text{etr}(-\Sigma_{11.2}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}] \mathbf{A}_{22}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}]^H) \end{array} \right\} \\ &= \left\{ \begin{array}{c} \left[ \frac{|\mathbf{A}_{11.2}|^{m-(n-k)-k}}{\Gamma_{m-(n-k),k} |\Sigma_{11.2}|^{m-(n-k)}} \text{etr}(-\Sigma_{11.2}^{-1} \mathbf{A}_{11.2}) \right] \left[ \frac{|\mathbf{A}_{22}|^{m-(n-k)}}{\Gamma_{m,n-k} |\Sigma_{22}|^m} \text{etr}(-\Sigma_{22}^{-1} \mathbf{A}_{22}) \right] \\ \cdot \left[ \frac{\Gamma_{m-(n-k),k} \Gamma_{m,n-k}}{\Gamma_{m,n}} \frac{|\mathbf{A}_{22}|^{-k}}{|\Sigma_{11.2}|^{n-k}} \right] \\ \cdot \text{etr}(-\Sigma_{11.2}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}] \mathbf{A}_{22}^{-1} [\mathbf{A}_{12} - \Sigma_{12} \Sigma_{22}^{-1} \mathbf{A}_{22}]^H) \end{array} \right\} \\ &= f(\mathbf{A}_{11.2}) f(\mathbf{A}_{22}) f(\mathbf{A}_{12} | \mathbf{A}_{22}), \end{aligned} \quad (33)$$

where it is recognized that  $\mathbf{A}_{11.2}$  and  $\mathbf{A}_{22}$  have complex Wishart distributions,

$$\mathbf{A}_{11.2} \sim \mathcal{CW}_k(m - (n - k), \Sigma_{11.2}) \quad (34)$$

and

$$\mathbf{A}_{22} \sim \mathcal{CW}_{n-k}(m, \Sigma_{22}), \quad (35)$$

thus proving the first and third parts of the theorem. The independence of  $\mathbf{A}_{11.2}$  from  $\mathbf{A}_{12}$  and  $\mathbf{A}_{22}$  is seen from the factorization of the probability density function of equation (33). The probability density function of  $\mathbf{A}_{12}$  conditioned on  $\mathbf{A}_{22}$  may be simplified by considering

$$\begin{aligned} \frac{\Gamma_{m-(n-k),k} \Gamma_{m,n-k}}{\Gamma_{m,n}} &= \left\{ \begin{array}{c} \frac{\Gamma(m-n+k) \cdots \Gamma(m-n+1) \Gamma(m) \cdots \Gamma(m-n+k+1)}{\Gamma(m) \cdots \Gamma(m-n+1)} \\ \cdot \pi^{\frac{1}{2} [k(k-1) + (n-k)(n-k-1) - n(n-1)]} \end{array} \right\} \\ &= \pi^{\frac{1}{2} [k^2 - k + n^2 - 2nk + k^2 - n + k - n^2 + n]} \\ &= \pi^{-k(n-k)}. \end{aligned} \quad (36)$$

Substituting equation (36) into the conditional probability density function of  $\mathbf{A}_{12}$  yields

$$f(\mathbf{A}_{12}|\mathbf{A}_{22}) = \frac{\text{etr}\left(-\boldsymbol{\Sigma}_{11\cdot 2}^{-1} [\mathbf{A}_{12} - \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\mathbf{A}_{22}] \mathbf{A}_{22}^{-1} [\mathbf{A}_{12} - \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\mathbf{A}_{22}]^H\right)}{\pi^{k(n-k)} |\mathbf{A}_{22}|^k |\boldsymbol{\Sigma}_{11\cdot 2}|^{n-k}} \quad (37)$$

which is recognized as the probability density function of a complex Gaussian random matrix,

$$\mathbf{A}_{12}|\mathbf{A}_{22} \sim \mathcal{CN}_{k \times (n-k)}(\boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\mathbf{A}_{22}, \boldsymbol{\Sigma}_{11\cdot 2} \otimes \mathbf{A}_{22}), \quad (38)$$

proving the second part of the theorem.

#### THEOREM 4

The following theorem is an extension of Theorem 3.2.11 on page 95 of Muirhead [2] describing the probability density function of the inverse of a matrix quadratic form involving the inverse of a complex Wishart distributed matrix.

**Theorem 4** *If  $\mathbf{A} \sim \mathcal{CW}_n(m, \boldsymbol{\Sigma})$ ,  $\boldsymbol{\Sigma}$  is full rank, and  $\mathbf{P}$  is a  $k$ -by- $n$  matrix with rank  $k$ , then*

$$\left(\mathbf{P}\mathbf{A}^{-1}\mathbf{P}^H\right)^{-1} \sim \mathcal{CW}_k\left(m - n + k, \left(\mathbf{P}\boldsymbol{\Sigma}^{-1}\mathbf{P}^H\right)^{-1}\right). \quad (39)$$

*Proof:* If  $\boldsymbol{\Sigma}$  is full rank, it may be factored into

$$\boldsymbol{\Sigma} = \boldsymbol{\Gamma}\boldsymbol{\Gamma}^H, \quad (40)$$

where  $\boldsymbol{\Gamma}$  is also of full rank. Set

$$\mathbf{B} = \boldsymbol{\Gamma}^{-1}\mathbf{A}\left(\boldsymbol{\Gamma}^H\right)^{-1}. \quad (41)$$

Since  $\boldsymbol{\Gamma}$  is constant and of full rank,  $\mathbf{B}$  is distributed as

$$\mathbf{B} \sim \mathcal{CW}_n(m, \mathbf{I}_n). \quad (42)$$

Define the  $k$ -by- $n$ , rank  $k$  matrix

$$\mathbf{R} = \mathbf{P} \left( \mathbf{\Gamma}^H \right)^{-1}. \quad (43)$$

Substituting equations (43) and (41) into the matrix product described in the theorem results in

$$\begin{aligned} \left( \mathbf{P} \mathbf{A}^{-1} \mathbf{P}^H \right)^{-1} &= \left( \mathbf{R} \mathbf{\Gamma}^H \mathbf{A}^{-1} \mathbf{\Gamma} \mathbf{R}^H \right)^{-1} \\ &= \left( \mathbf{R} \left[ \mathbf{\Gamma}^{-1} \mathbf{A} \left( \mathbf{\Gamma}^H \right)^{-1} \right]^{-1} \mathbf{R}^H \right)^{-1} \\ &= \left( \mathbf{R} \mathbf{B}^{-1} \mathbf{R}^H \right)^{-1}. \end{aligned} \quad (44)$$

Similarly,

$$\begin{aligned} \left( \mathbf{P} \mathbf{\Sigma}^{-1} \mathbf{P}^H \right)^{-1} &= \left( \mathbf{R} \mathbf{\Gamma}^H \mathbf{\Sigma}^{-1} \mathbf{\Gamma} \mathbf{R}^H \right)^{-1} \\ &= \left[ \mathbf{R} \left( \mathbf{\Gamma}^{-1} \mathbf{\Sigma} \left( \mathbf{\Gamma}^H \right)^{-1} \right)^{-1} \mathbf{R}^H \right]^{-1} \\ &= \left( \mathbf{R} \mathbf{R}^H \right)^{-1}. \end{aligned} \quad (45)$$

Using equations (44) and (45), it is seen that the theorem, as stated in equation (39), simplifies to showing that

$$\left( \mathbf{R} \mathbf{B}^{-1} \mathbf{R}^H \right)^{-1} \sim \mathcal{CW}_n \left( m - n + k, \left( \mathbf{R} \mathbf{R}^H \right)^{-1} \right). \quad (46)$$

Using a singular value decomposition, the matrix  $\mathbf{R}$  may be factored into

$$\mathbf{R} = \mathbf{U} \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} \mathbf{V}^H, \quad (47)$$

where  $\mathbf{U}$  is  $k$ -by- $k$  and non-singular and  $\mathbf{V}$  is  $n$ -by- $n$  and orthogonal,

$$\mathbf{V}^{-1} = \mathbf{V}^H. \quad (48)$$

Substituting this factorization into equations (44) and (45) results in

$$\left( \mathbf{R} \mathbf{B}^{-1} \mathbf{R}^H \right)^{-1} = \left( \mathbf{U} \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} \mathbf{V}^H \mathbf{B}^{-1} \mathbf{V} \begin{bmatrix} \mathbf{I}_k \\ \mathbf{0} \end{bmatrix} \mathbf{U}^H \right)^{-1}$$

$$\begin{aligned}
&= (\mathbf{U}^H)^{-1} \left( \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} (\mathbf{V}^H \mathbf{B} \mathbf{V})^{-1} \begin{bmatrix} \mathbf{I}_k \\ \mathbf{0} \end{bmatrix} \right)^{-1} \mathbf{U}^{-1} \\
&= (\mathbf{U}^H)^{-1} \left( \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} \mathbf{C}^{-1} \begin{bmatrix} \mathbf{I}_k \\ \mathbf{0} \end{bmatrix} \right)^{-1} \mathbf{U}^{-1}, \tag{49}
\end{aligned}$$

where

$$\mathbf{C} = \mathbf{V}^H \mathbf{B} \mathbf{V}, \tag{50}$$

and

$$\begin{aligned}
(\mathbf{R} \mathbf{R}^H)^{-1} &= \left( \mathbf{U} \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} \mathbf{V}^H \mathbf{V} \begin{bmatrix} \mathbf{I}_k \\ \mathbf{0} \end{bmatrix} \mathbf{U}^H \right)^{-1} \\
&= \left( \mathbf{U} \begin{bmatrix} \mathbf{I}_k & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{I}_k \\ \mathbf{0} \end{bmatrix} \mathbf{U}^H \right)^{-1} \\
&= (\mathbf{U} \mathbf{U}^H)^{-1}. \tag{51}
\end{aligned}$$

Let

$$\mathbf{C} = \begin{bmatrix} \mathbf{C}_{11} & \mathbf{C}_{12} \\ \mathbf{C}_{21} & \mathbf{C}_{22} \end{bmatrix} \tag{52}$$

and

$$\mathbf{C}^{-1} = \mathbf{D} = \begin{bmatrix} \mathbf{D}_{11} & \mathbf{D}_{12} \\ \mathbf{D}_{21} & \mathbf{D}_{22} \end{bmatrix} \tag{53}$$

be  $k$ -by- $n-k$  partitions of  $\mathbf{C}$  and  $\mathbf{D} = \mathbf{C}^{-1}$ . Then, equation (49) becomes

$$\begin{aligned}
(\mathbf{R} \mathbf{B}^{-1} \mathbf{R}^H)^{-1} &= (\mathbf{U}^H)^{-1} \mathbf{D}_{11}^{-1} \mathbf{U}^{-1} \\
&= (\mathbf{U}^H)^{-1} \mathbf{C}_{11.2} \mathbf{U}^{-1}, \tag{54}
\end{aligned}$$

where  $\mathbf{C}_{11.2}$  is as defined in Theorem 3 and, as seen in Muirhead [2], is equal to  $\mathbf{D}_{11}^{-1}$ .

Since  $\mathbf{V}$  is orthogonal,  $\mathbf{C}$  is distributed as

$$\mathbf{C} \sim \mathcal{CW}_n(m, \mathbf{I}_n). \tag{55}$$

Applying part (i) of Theorem 3, it is seen that

$$\mathbf{C}_{11.2} \sim \mathcal{CW}_k(m - n + k, \mathbf{I}_k), \tag{56}$$

which, when applied to equation (54), results in

$$\begin{aligned} (\mathbf{R}\mathbf{B}^{-1}\mathbf{R}^H)^{-1} &\sim \mathcal{CW}_k\left(m-n+k, (\mathbf{U}\mathbf{U}^H)^{-1}\right) \\ &\sim \mathcal{CW}_k\left(m-n+k, (\mathbf{R}\mathbf{R}^H)^{-1}\right), \end{aligned} \quad (57)$$

which completes the proof.

### THEOREM 5

The following theorem is an extension of Theorem 3.2.12 on page 96 of Muirhead [2] describing the probability density function of the ratio of quadratic forms involving the inverse of the scale matrix and the inverse of a random sample of a complex Wishart distribution. It is interesting to note that the resulting distribution does not depend on the vector in the quadratic forms if it is independent of the complex Wishart distributed matrix.

**Theorem 5** *If  $\mathbf{A} \sim \mathcal{CW}_n(m, \Sigma)$  where  $m > n - 1$  and if  $\mathbf{y}$  is any  $n$ -by-1 random vector independent of  $\mathbf{A}$  such that  $\Pr(\mathbf{y} = \mathbf{0}) = 0$ , then*

$$2\frac{\mathbf{y}^H \Sigma^{-1} \mathbf{y}}{\mathbf{y}^H \mathbf{A}^{-1} \mathbf{y}} \sim \chi_{2(m-n+1)}^2, \quad (58)$$

*and is independent of  $\mathbf{y}$ .*

*Proof:* In Theorem 4, let  $\mathbf{P} = \mathbf{y}^H$ . Then,

$$W = (\mathbf{y}^H \mathbf{A}^{-1} \mathbf{y})^{-1} \sim \mathcal{CW}_1\left(m-n+1, (\mathbf{y}^H \Sigma^{-1} \mathbf{y})^{-1}\right), \quad (59)$$

which has probability density function,

$$\begin{aligned} f_W(w) &= \frac{|w|^{m-n+1-1} \text{etr}\left(-\frac{w}{\theta}\right)}{\Gamma(m-n+1) |\theta|^{m-n+1}} \\ &= \frac{w^{m-n} e^{-\frac{w}{\theta}}}{\Gamma(m-n+1) \theta^{m-n+1}}, \end{aligned} \quad (60)$$

where

$$\theta = (\mathbf{y}^H \Sigma^{-1} \mathbf{y})^{-1}. \quad (61)$$

Performing the transformation

$$Z = \frac{2}{\theta} W = 2 \frac{\mathbf{y}^H \boldsymbol{\Sigma}^{-1} \mathbf{y}}{\mathbf{y}^H \mathbf{A}^{-1} \mathbf{y}}, \quad (62)$$

yields the desired scaled ratio of quadratic forms in equation (58). The probability density function of  $Z$  is found to be

$$\begin{aligned} f_Z(z) &= \frac{\theta \left(\frac{z\theta}{2}\right)^{m-n} e^{-\frac{z}{2}}}{\Gamma(m-n+1) 2\theta^{m-n+1}} \\ &= \frac{z^{m-n+1-1} e^{-\frac{z}{2}}}{\Gamma(m-n+1) 2^{m-n+1}} \\ &= \frac{z^{\frac{r}{2}-1} e^{-\frac{z}{2}}}{\Gamma\left(\frac{r}{2}\right) 2^{\frac{r}{2}}}, \end{aligned} \quad (63)$$

where

$$r = 2(m-n+1), \quad (64)$$

which is a central Chi-squared distribution with  $2(m-n+1)$  degrees of freedom.

## References

- [1] S. R. Searle, *Linear Models*. John Wiley & Sons, 1971.
- [2] R. J. Muirhead, *Aspects of Multivariate Statistical Theory*. John Wiley & Sons, 1982.
- [3] N. R. Goodman, "Statistical Analysis Based on a Certain Multivariate Complex Gaussian Distribution (An Introduction)," *The Annals of Mathematical Statistics*, vol. 34, pp. 152–177, March 1963.
- [4] T. W. Anderson, *An Introduction to Multivariate Statistical Analysis*. John Wiley & Sons, 1984.
- [5] Johnson and Kotz, eds., *Encyclopedia of Statistical Sciences*, vol. 6, pp. 276–284. John Wiley & Sons, 1985.

NUWC Technical Memorandum No. 931133

DISTRIBUTION LIST, EXTERNAL

T. G. Goldsberry    ONR Code 451  
R. Young            ONR Code 451

DISTRIBUTION LIST, INTERNAL

NEW LONDON

D. Abraham (10)    3314  
R. Barton            3314  
G. C. Carter        2192  
W. Chang            3314  
G. Connolly        2192  
R. Dwyer            3331  
J. Fay                3331  
M. Gouzie           2121  
S. Greineder        2121  
F. Khan              3314  
I. Kirsteins         3314  
R. Kneipfer         214  
J. Ianniello         2123  
R. Latourette       2152  
D. Lerro             3314  
R. Lynch             3331  
T. Luginbuhl        2121  
R. Mason             3331  
A. Nuttall            302  
J. Nuttall            2121  
N. Owsley            2123  
K. Peters            3314  
D. Sheldon          3314  
Library (2)

NEWPORT

R. Streit            22101  
Library (2)

WEST PALM

R. Kennedy         3802