



INDUSTRIAL
MATHEMATICS
INSTITUTE

2003:17

The volume estimates and their
applications

B.S. Kashin and V.N. Temlyakov

IMI
Preprint Series

Department of Mathematics
University of South Carolina

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 2003		2. REPORT TYPE		3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE The volume estimates and their applications				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of South Carolina, Department of Mathematics, Columbia, SC, 29208				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We prove new estimates for the entropy numbers of classes of multivariate functions with bounded mixed derivative. It is known that the investigation of these classes requires development of new techniques comparing to the univariate classes. In this paper we continue to develop the technique based on estimates of volumes of sets of the Fourier coefficients of trigonometric polynomials with frequencies in special regions. We obtain new volume estimates and use them to get right orders of decay of the entropy numbers of classes of functions of two variables with a mixed derivative bounded in the L1-norm. This is the first such result for these classes. This result essentially completes the investigation of orders of decay of the entropy numbers of classes of functions of two variables with bounded mixed derivative. The case of similar classes of functions of more than two variables is still open.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

THE VOLUME ESTIMATES AND THEIR APPLICATIONS¹

B.S. KASHIN AND V.N. TEMLYAKOV

ABSTRACT. We prove new estimates for the entropy numbers of classes of multivariate functions with bounded mixed derivative. It is known that the investigation of these classes requires development of new techniques comparing to the univariate classes. In this paper we continue to develop the technique based on estimates of volumes of sets of the Fourier coefficients of trigonometric polynomials with frequencies in special regions. We obtain new volume estimates and use them to get right orders of decay of the entropy numbers of classes of functions of two variables with a mixed derivative bounded in the L_1 -norm. This is the first such result for these classes. This result essentially completes the investigation of orders of decay of the entropy numbers of classes of functions of two variables with bounded mixed derivative. The case of similar classes of functions of more than two variables is still open.

THIS PAPER IS DEDICATED TO THE 70TH BIRTHDAY OF S.A. TELYAKOVSKII

1. INTRODUCTION

We obtain in this paper new estimates of volumes of sets of the Fourier coefficients of trigonometric polynomials of two variables with frequencies in a *hyperbolic layer*. Section 2 is devoted to this kind of estimates. Then, in Section 3 we use results of Section 2 to prove lower estimates for the entropy numbers of classes of functions with bounded mixed derivative. The new volume estimates from Section 2 allowed us to essentially complete the investigation of orders of decay of the entropy numbers $\epsilon_m(W_{q,\alpha}^r, L_p)$ for all $1 \leq q, p \leq \infty$ and large enough r in the case of classes of functions of two variables. In Section 4 we apply results of Section 2 to the following problem of discretization. We look for a set of points with a property: (E) the uniform norm on this set of points is equivalent to the uniform norm for any trigonometric polynomial with frequencies from a given hyperbolic layer. We give in Section 4 other proof of the following surprising result from [KT3], [KT4]: the cardinality of a set with the property (E) must grow at least as N^{1+a} , $a > 0$, where N is the dimension of the corresponding subspace of trigonometric polynomials. The proof of the above result from [KT3], [KT4] made use of extremal properties of the multivariate normal distribution and an inequality for the trigonometric polynomials from [T2]. Later, other inequality for the trigonometric polynomials has been obtained in [T6]. Our proof in

¹The second author was supported by the National Science Foundation Grant DMS 0200187 and by ONR Grant N00014-96-1-1003

this paper uses this new inequality and the volume estimates from Section 2. This allowed us to simplify the original proof (see [KT4]).

We now formulate the main results of the paper. Let $s = (s_1, \dots, s_d)$ be a vector with nonnegative integer coordinates ($s \in \mathbb{Z}_+^d$) and

$$\rho(s) := \{k = (k_1, \dots, k_d) \in \mathbb{Z}_+^d : [2^{s_j-1}] \leq |k_j| < 2^{s_j}, \quad j = 1, \dots, d\}$$

where $[a]$ denotes the integer part of a number a . Denote for a natural number n

$$Q_n := \cup_{\|s\|_1 \leq n} \rho(s); \quad \Delta Q_n := Q_n \setminus Q_{n-1} = \cup_{\|s\|_1 = n} \rho(s)$$

with $\|s\|_1 = s_1 + \dots + s_d$ for $s \in \mathbb{Z}_+^d$. We call a set ΔQ_n *hyperbolic layer*. For a set $\Lambda \subset \mathbb{Z}^d$ denote

$$\mathcal{T}(\Lambda) := \{f \in L_1 : \hat{f}(k) = 0, k \in \mathbb{Z} \setminus \Lambda\}, \quad \mathcal{T}_R(\Lambda) := \{f \in \mathcal{T}(\Lambda) : \hat{f}(k) \in \mathbb{R}, k \in \Lambda\}.$$

For a finite set Λ we assign to each $f = \sum_{k \in \Lambda} \hat{f}(k) e^{i(k,x)} \in \mathcal{T}(\Lambda)$ a vector

$$A(f) := \{(\operatorname{Re} \hat{f}(k), \operatorname{Im} \hat{f}(k)), \quad k \in \Lambda\} \in \mathbb{R}^{2|\Lambda|}$$

where $|\Lambda|$ denotes the cardinality of Λ and define

$$B_\Lambda(L_p) := \{A(f) : f \in \mathcal{T}(\Lambda), \quad \|f\|_p \leq 1\}.$$

The volume estimates of the sets $B_\Lambda(L_p)$ and related questions have been studied in a number of papers of the authors of this paper: the case $\Lambda = [-n, n]$, $p = \infty$ in [K1]; the case $\Lambda = [-N_1, N_1] \times \dots \times [-N_d, N_d]$, $p = \infty$ in [T2], [T3]; the case of arbitrary Λ and $p = 1$ in [KT1]. In particular, the results of [KT1] imply (see Theorem 2.4 of this paper) for $d = 2$ and $1 \leq p < \infty$ that

$$(\operatorname{vol}(B_{\Delta Q_n}(L_p)))^{(2|\Delta Q_n|)^{-1}} \asymp |\Delta Q_n|^{-1/2} \asymp (2^n n)^{-1/2}.$$

We will prove in Section 2 (see Theorem 2.5) that in the case $p = \infty$ the volume estimate is different:

$$(\operatorname{vol}(B_{\Delta Q_n}(L_\infty)))^{(2|\Delta Q_n|)^{-1}} \asymp (2^n n^2)^{-1/2}.$$

We note that in the case $\Lambda = [-N_1, N_1] \times \dots \times [-N_d, N_d]$ the volume estimate is the same for all $1 \leq p \leq \infty$. We discuss this in more detail in Section 2.

In Section 3 we apply the volume estimates to the problem of asymptotic behavior of the entropy numbers of the classes $W_{1,\alpha}^r$. We now give the corresponding definitions. For a natural number m and a compact set F in a Banach space X with the unit ball $B(X)$ we define the m th entropy number as

$$\epsilon_m(F, X) := \inf \{ \epsilon : \exists f_1, \dots, f_{2^m} \in X : F \subset \cup_{j=1}^{2^m} (f_j + \epsilon B(X)) \}.$$

Let $r > 0$ and $\alpha \in \mathbb{R}$. Define

$$F_r(t, \alpha) := 1 + 2 \sum_{k=1}^{\infty} k^{-r} \cos(kt - \alpha\pi/2), \quad t \in [0, 2\pi].$$

We define for $x = (x_1, x_2)$ and $\alpha = (\alpha_1, \alpha_2)$

$$F_r(x, \alpha) := F_r(x_1, \alpha_1)F_r(x_2, \alpha_2).$$

Finally, we define

$$W_{q,\alpha}^r := \{f : f = F_r(\cdot, \alpha) * \varphi, \quad \|\varphi\|_q \leq 1\}$$

where $*$ means convolution. The problem of estimating $\epsilon_m(W_{q,\alpha}^r, L_p)$ has a long history. We will mention some results only in the case $d = 2$. The first result on the right order of $\epsilon_m(W_{q,\alpha}^r, L_p)$ in the case $p = q = 2$ has been obtained by S.A. Smolyak [Sm] in 1960. Here is a list of further contributions: Dinh Dung [D], 1985, the case $1 < q = p < \infty$; V.N. Temlyakov [T1], [T2], 1988, the case $1 < q, p < \infty, r > 1$ and the case $p = 1, 1 < q < \infty$; E.S. Belinsky [B], 1990, the case $p = 1, q = \infty, r > 1/2$; B.S. Kashin, V.N. Temlyakov [KT1], [KT2], 1994–1995, the case $p = 1, q = \infty, r > 0$; J. Kuelbs, W.V. Li [KL], 1993, the case $p = \infty, q = 2, r = 1$; V.N. Temlyakov [T5], 1995, the case $p = \infty, 1 < q < \infty$; V.N. Temlyakov [T6], 1998, the case $p = \infty, q = \infty, r > 1/2$.

In Section 3 we study the case left open: $q = 1, 1 \leq p \leq \infty$. We prove that (see Theorem 3.1):

$$(1.1) \quad \epsilon_m(W_{1,\alpha}^r, L_p) \asymp m^{-r} (\log m)^{r+1/2}, \quad 1 \leq p < \infty, \quad r > \max(1/2, 1 - 1/p)$$

and

$$\epsilon_m(W_{1,0}^r, L_\infty) \asymp m^{-r} (\log m)^{r+1}, \quad r > 1.$$

It is interesting to compare these estimates with the case $1 < q, p < \infty$ where we have

$$(1.2) \quad \epsilon_m(W_{q,\alpha}^r, L_p) \asymp m^{-r} (\log m)^r.$$

We note that when we turn from $q > 1$ to $q = 1$ the exponent for $\log m$ jumps from r in (1.2) to $r + 1/2$ in (1.1).

2. THE VOLUME ESTIMATES

The main goal of this section is to prove new estimates of volumes of sets of the Fourier coefficients of trigonometric polynomials of two variables (dimension $d = 2$) with frequencies in a *hyperbolic layer*. In some cases we will give estimates in the arbitrary dimension d . It is well known (see [K1], [T2], [T4], [KT1]) that the volume estimates of the above mentioned sets can be used in different problems of approximation theory including the problem of estimating the entropy numbers of function classes. We use the notations $\mathcal{T}(\Lambda)$, $A(f)$, and $B_\Lambda(L_p)$ introduced in Section 1. In the case $\Lambda = \Pi(N) := [-N_1, N_1] \times \cdots \times [-N_d, N_d]$, $N := (N_1, \dots, N_d)$, the volume estimates are known. We formulate it as a theorem.

Theorem 2.1. *For any $1 \leq p \leq \infty$ we have*

$$(\text{vol}(B_{\Pi(N)}(L_p)))^{(2|\Pi(N)|)^{-1}} \asymp |\Pi(N)|^{-1/2},$$

with constants in \asymp that may depend only on d .

We note that the most difficult part of Theorem 2.1 is the lower estimate for $p = \infty$. The corresponding estimate was proved in the case $d = 1$ in [K1] and in the general case in [T2] and [T3]. The upper estimate for $p = 1$ in Theorem 2.1 can be easily reduced to the volume estimate for an octahedron (see, for instance [T4]).

The results of [KT1] imply the following estimate.

Theorem 2.2. *For any finite set $\Lambda \subset \mathbb{Z}^d$ and any $1 \leq p \leq 2$ we have*

$$\text{vol}(B_\Lambda(L_p))^{(2|\Lambda|)^{-1}} \asymp |\Lambda|^{-1/2}.$$

The following result of Bourgain-Milman [BM] plays an important role in the volume estimates of finite dimensional bodies.

Theorem 2.3. *For any convex centrally symmetric body $K \subset \mathbb{R}^n$ we have*

$$(\text{vol}(K)\text{vol}(K^\circ))^{1/n} \asymp (\text{vol}(B_2^n))^{2/n} \asymp 1/n$$

where K° is a polar for K , that is

$$K^\circ := \{x \in \mathbb{R}^n : \sup_{y \in K} (x, y) \leq 1\}.$$

Having in mind application of Theorem 2.3 we define some other than $B_\Lambda(L_p)$ sets. Let

$$E_\Lambda^\perp(f)_p := \inf_{g \perp \mathcal{T}(\Lambda)} \|f - g\|_p, \quad E_{\Lambda, R}^\perp(f)_p := \inf_{g \perp \mathcal{T}(\Lambda), \hat{g}(k) \in \mathbb{R}} \|f - g\|_p$$

and

$$B_\Lambda^\perp(L_p) := \{A(f) : f \in \mathcal{T}(\Lambda), \quad E_\Lambda^\perp(f)_p \leq 1\}.$$

It is clear that

$$B_\Lambda(L_p) \subseteq B_\Lambda^\perp(L_p).$$

Moreover, if the orthogonal projector P_Λ onto $\mathcal{T}(\Lambda)$ is bounded as an operator from L_p to L_p then we have

$$(2.1) \quad \text{vol}(B_\Lambda(L_p))^{(2|\Lambda|)^{-1}} \asymp \text{vol}(B_\Lambda^\perp(L_p))^{(2|\Lambda|)^{-1}}.$$

For example it is the case when $\Lambda = \cup_{s \in A} \rho(s)$. We also consider

$$B_{\Lambda, R}(L_p) := \{A(f) : \|f\|_p \leq 1, \quad f \in \mathcal{T}(\Lambda), \quad \hat{f}(k) \in \mathbb{R}\}$$

and

$$B_{\Lambda,R}^\perp(L_p) := \{A(f) : E_{\Lambda,R}^\perp(f)_p \leq 1, \quad f \in \mathcal{T}(\Lambda), \quad \hat{f}(k) \in \mathbb{R}\}.$$

Using the Nikol'skii duality theorem and the basic properties of the de la Vallée Poussin operators one can prove the following relation

$$(2.2) \quad B_{\Lambda,R}(L_p)^\circ = B_{\Lambda,R}^\perp(L_{p'}), \quad p' = \frac{p}{p-1}.$$

Proof of (2.2). This proof uses standard ideas from the duality arguments. We will not give all the details of the proof and, instead, we will outline the main steps of the proof.

First, we note that the relation

$$B_{\Lambda,R}^\perp(L_{p'}) \subseteq B_{\Lambda,R}(L_p)^\circ$$

follows immediately from the inequality that holds for any $f, g \in \mathcal{T}_R(\Lambda)$:

$$\left| \sum_{k \in \Lambda} \hat{f}(k) \hat{g}(k) \right| = \frac{1}{2\pi} \left| \int_0^{2\pi} f \bar{g} \right| \leq E_{\Lambda}^\perp(f)_{p'} \|g\|_p.$$

Second, we prove the inverse inclusion

$$(2.2A) \quad B_{\Lambda,R}(L_p)^\circ \subseteq B_{\Lambda,R}^\perp(L_{p'}).$$

Let X denote the real Banach space

$$L_{p,R} := \{f \in L_p : \hat{f}(k) \in \mathbb{R}\}$$

with $\|f\|_X := \|f\|_p$. Then any $u \in L_{p',R}$ defines a continuous linear functional on X by the formula

$$(u, f) := \frac{1}{2\pi} \left| \int_0^{2\pi} f \bar{u} \right|.$$

For such a functional we have $\|u\|_{X^*} = \|u\|_{p'}$. Indeed, the inequality $\|u\|_{X^*} \leq \|u\|_{p'}$ follows from the Hölder inequality and the opposite one $\|u\|_{X^*} \geq \|u\|_{p'}$ follows from the observation that for any $f \in L_p$ the function $v(x) := \frac{1}{2}(f(x) + \bar{f}(-x))$ is from $L_{p,R}$ and $\|v\|_p \leq \|f\|_p$.

We continue the proof of (2.2A). Let $f \in B_{\Lambda,R}(L_p)^\circ$. Then from the definition of the polar we get that for any $g \in \mathcal{T}(\Lambda)$, $\hat{g}(k) \in \mathbb{R}$, $\|g\|_p \leq 1$ the following inequality holds

$$\left| \sum_{k \in \Lambda} \hat{f}(k) \hat{g}(k) \right| = |(g, f)| \leq 1.$$

For $N \in \mathbb{N}$ denote

$$E_{\Lambda,R}^{\perp,N}(f)_p := \inf_{g \perp \mathcal{T}(\Lambda), g \in \mathcal{T}([-N,N]), \hat{g}(k) \in \mathbb{R}} \|f - g\|_p.$$

Then obviously for any N

$$E_{\Lambda,R}^\perp(f)_{p'} \leq E_{\Lambda,R}^{\perp,N}(f)_{p'}.$$

Next, using the appropriate de la Vallée Poussin operators one can check that for any $g \in \mathcal{T}_R(\Lambda \cup (\mathbb{Z} \setminus [-N, N]))$ with $\|g\|_p \leq 1$ we have

$$\|g_\Lambda\|_p \leq 1 + \epsilon_N, \quad \epsilon_N \rightarrow 0, \quad N \rightarrow \infty$$

where

$$g_\Lambda := \sum_{k \in \Lambda} \hat{g}(k) e^{ikx}.$$

Therefore, by the Nikol'skii Duality Theorem we obtain

$$E_{\Lambda,R}^{\perp,N}(f)_{p'} = \sup_{g \in \mathcal{T}_R(\Lambda \cup (\mathbb{Z} \setminus [-N, N])), \|g\|_p \leq 1} |(f, g)| \leq \sup_{g_\Lambda: \|g_\Lambda\|_p \leq 1 + \epsilon_N, g_\Lambda \in \mathcal{T}_R(\Lambda)} |(f, g)| \leq 1 + \epsilon_N.$$

This completes the proof of (2.2A).

We will now show that the volumes of $B_\Lambda(L_p) \subset \mathbb{R}^{2|\Lambda|}$ and $B_{\Lambda,R}(L_p) \subset \mathbb{R}^{|\Lambda|}$ are closely related. First of all if

$$\left\| \sum_{k \in \Lambda} a_k e^{i(k,x)} \right\|_p \leq 1/2, \quad \left\| \sum_{k \in \Lambda} b_k e^{i(k,x)} \right\|_p \leq 1/2$$

then

$$\left\| \sum_{k \in \Lambda} (a_k + ib_k) e^{i(k,x)} \right\|_p \leq 1.$$

Therefore,

$$(2.3) \quad \left(\frac{1}{2} B_{\Lambda,R}(L_p)\right) \otimes \left(\frac{1}{2} B_{\Lambda,R}(L_p)\right) \subseteq B_\Lambda(L_p).$$

Next, let

$$f(x) = \sum_{k \in \Lambda} (a_k + ib_k) e^{i(k,x)}.$$

Then

$$\sum_{k \in \Lambda} a_k e^{i(k,x)} = \frac{1}{2} (f(x) + \bar{f}(-x))$$

and

$$i \sum_{k \in \Lambda} b_k e^{i(k,x)} = \frac{1}{2} (f(x) - \bar{f}(-x)).$$

This implies that

$$(2.4) \quad B_\Lambda(L_p) \subseteq B_{\Lambda,R}(L_p) \otimes B_{\Lambda,R}(L_p).$$

We get from (2.3) and (2.4) that

$$(2.5) \quad (\text{vol}(B_\Lambda(L_p)))^{(2|\Lambda|)^{-1}} \asymp (\text{vol}(B_{\Lambda,R}(L_p)))^{(|\Lambda|)^{-1}}.$$

Similarly we get

$$(2.6) \quad (\text{vol}(B_\Lambda^\perp(L_p)))^{(2|\Lambda|)^{-1}} \asymp (\text{vol}(B_{\Lambda,R}^\perp(L_p)))^{(|\Lambda|)^{-1}}.$$

This observation, Theorems 2.2 and 2.3 combined with (2.1) imply the following statement.

Theorem 2.4. *Let Λ have the form $\Lambda = \cup_{s \in S} \rho(s)$, $S \subset \mathbb{Z}_+^d$ is a finite set. Then for any $1 \leq p < \infty$ we have*

$$\text{vol}(B_\Lambda(L_p))^{(2|\Lambda|)^{-1}} \asymp |\Lambda|^{-1/2}.$$

We now proceed to the main results of this section. Denote $N := 2|\Delta Q_n|$.

Theorem 2.5. *In the case $d = 2$ we have*

$$(2.7) \quad (\text{vol}(B_{\Delta Q_n}(L_\infty)))^{1/N} \asymp (2^n n^2)^{-1/2};$$

$$(2.8) \quad (\text{vol}(B_{\Delta Q_n}^\perp(L_1)))^{1/N} \asymp 2^{-n/2}.$$

It is interesting to compare the first relation in Theorem 2.5 with the following estimate for $1 \leq p < \infty$ that follows from Theorem 2.4

$$(2.9) \quad (\text{vol}(B_{\Delta Q_n}(L_p)))^{1/N} \asymp (2^n n)^{-1/2}.$$

We see that in the case $\Lambda = \Delta Q_n$ unlike the case $\Lambda = \Pi(N_1, \dots, N_d)$ the estimate for $p = \infty$ is different from the estimate for $1 \leq p < \infty$.

Proof of Theorem 2.5. We begin with the proof of the lower estimate in (2.7). We will formulate and prove it in a more general form.

Lemma 2.1. *Let $\Lambda \subseteq [-2^n, 2^n]^d$ and $N := 2|\Lambda|$. Then*

$$(\text{vol}(B_\Lambda(L_\infty)))^{1/N} \geq C(d)(Nn)^{-1/2}.$$

Proof. We use the following result of E. Gluskin [G].

Theorem 2.6. *Let $Y = \{y_1, \dots, y_M\} \subset \mathbb{R}^N$, $\|y_i\| = 1$, $i = 1, \dots, M$ and*

$$W(Y) := \{x \in \mathbb{R}^N : |(x, y_i)| \leq 1, \quad i = 1, \dots, M\}.$$

Then

$$(\text{vol}(W(Y)))^{1/N} \geq C(\log(M/N))^{-1/2}.$$

Consider the following lattice on the \mathbb{T}^d :

$$G_n := \{x(l) = (l_1, \dots, l_d)\pi 2^{-n-1}, \quad 1 \leq l_j \leq 2^{n+2}, \quad l_j \in \mathbb{N}, \quad j = 1, \dots, d\}.$$

It is clear that $|G_n| = 2^{d(n+2)}$. It is well known that for any $f \in \mathcal{T}([-2^n, 2^n]^d)$ one has

$$\|f\|_\infty \leq C_1(d) \max_{x \in G_n} |f(x)|.$$

Thus, for any $\Lambda \subseteq [-2^n, 2^n]^d$ we have

$$(2.10) \quad \{A(f) : f \in \mathcal{T}(\Lambda), \quad |f(x)| \leq C_1(d)^{-1}, \quad x \in G_n\} \subseteq B_\Lambda(L_\infty).$$

Further

$$|f(x)|^2 = \left| \sum_{k \in \Lambda} \hat{f}(k) e^{i(k,x)} \right|^2 = \left(\sum_{k \in \Lambda} \operatorname{Re} \hat{f}(k) \cos(k, x) - \operatorname{Im} \hat{f}(k) \sin(k, x) \right)^2 + \left(\sum_{k \in \Lambda} \operatorname{Re} \hat{f}(k) \sin(k, x) + \operatorname{Im} \hat{f}(k) \cos(k, x) \right)^2.$$

We associate with each point $x \in G_n$ two vectors $y^1(x)$ and $y^2(x)$ from \mathbb{R}^N :

$$y^1(x) := \{(\cos(k, x), -\sin(k, x)), \quad k \in \Lambda\},$$

$$y^2(x) := \{(\sin(k, x), \cos(k, x)), \quad k \in \Lambda\}.$$

Then

$$\|y^1(x)\|^2 = \|y^2(x)\|^2 = |\Lambda|$$

and

$$|f(x)|^2 = (A(f), y^1(x))^2 + (A(f), y^2(x))^2.$$

It is clear that the condition $|f(x)| \leq C_1(d)^{-1}$ is satisfied if

$$|(A(f), y^i(x))| \leq 2^{-1/2} C_1(d)^{-1}, \quad i = 1, 2.$$

Let now

$$Y := \{y^i(x) / \|y^i(x)\|, \quad x \in G_n, \quad i = 1, 2\}.$$

Then $M = 2^{d(n+2)+1}$ and by Theorem 2.6

$$(2.11) \quad (\operatorname{vol}(W(Y)))^{1/N} \gg (\log(M/N))^{-1/2} \gg n^{-1/2}.$$

Using that the condition

$$|(A(f), y^i(x))| \leq 1$$

is equivalent to the condition

$$|(A(f), y^i(x) / \|y^i(x)\|)| \leq (N/2)^{-1/2}$$

we get from (2.10) and (2.11)

$$(\operatorname{vol}(B_\Lambda(L_\infty)))^{1/N} \gg (Nn)^{-1/2}.$$

This completes the proof of Lemma 2.1

Lemma 2.1 implies immediately the lower estimate in (2.7) because in this case $|\Delta Q_n| \asymp 2^n n$. We emphasize that Lemma 2.1 shows that the lower estimate similar to (2.7) holds for any $\Lambda \subseteq [-2^n, 2^n]^2$ with $|\Lambda| \asymp |\Delta Q_n|$ and therefore it does not depend on the geometry of Λ .

We now proceed to the proof of the upper estimate in (2.7). This proof uses the geometry of ΔQ_n . Comparing the estimate (2.7) with Theorem 2.1 we conclude that the upper estimate in (2.7) cannot be generalized for all $\Lambda \subseteq [-2^n, 2^n]^2$ with $|\Lambda| \asymp |\Delta Q_n|$. We prove first the lower estimate in (2.8). We will use the following lemma that follows directly from Lemma 2.4 in [T6].

Lemma 2.2. *Let $d = 2$. For any $f \in \mathcal{T}(\Delta Q_n)$ satisfying*

$$\|\delta_s(f)\|_\infty \leq 1, \quad \|s\|_1 = n,$$

we have

$$E_{Q_n}^\perp(f)_1 \leq C.$$

Denote

$$H_\infty(\Delta Q_n) := \{f \in \mathcal{T}(\Delta Q_n) : \|\delta_s(f)\|_\infty \leq 1\}$$

and

$$A(H_\infty(\Delta Q_n)) := \{A(f) : f \in H_\infty(\Delta Q_n)\}.$$

Lemma 2.2 implies that ($N = 2|\Delta Q_n|$)

$$(2.12) \quad (\text{vol}(B_{\Delta Q_n}^\perp(L_1)))^{1/N} \gg (\text{vol}(A(H_\infty(\Delta Q_n))))^{1/N}.$$

Using Theorem 2.1 we get

$$(2.13) \quad (\text{vol}(A(H_\infty(\Delta Q_n))))^{1/N} = \left(\prod_{\|s\|_1=n} \text{vol}(A(\mathcal{T}(\rho(s))_\infty)) \right)^{1/N} \gg 2^{-n/2},$$

where

$$\mathcal{T}(\rho(s))_\infty := \{t \in \mathcal{T}(\rho(s)) : \|t\|_\infty \leq 1\}.$$

The lower estimate in (2.8) follows from (2.12) and (2.13).

Using Theorem 2.3 and relations (2.2), (2.5), and (2.6) we complete the proof of Theorem 2.5.

3. ESTIMATES OF THE ϵ -ENTROPY

In this section we use the results from Section 2 for obtaining new lower estimates for the ϵ -entropy of the classes $W_{1,\alpha}^r$ in L_p , $1 \leq p \leq \infty$. We confine ourselves to the case of functions of two variables. We prove the following theorem here.

Theorem 3.1. *The following relations hold*

$$(3.1) \quad \epsilon_m(W_{1,\alpha}^r, L_p) \asymp m^{-r}(\log m)^{r+1/2}, \quad 1 \leq p < \infty, \quad r > \max(1/2, 1 - 1/p);$$

$$(3.2) \quad \epsilon_m(W_{1,0}^r, L_\infty) \asymp m^{-r}(\log m)^{r+1}, \quad r > 1.$$

Proof. We first prove the upper estimates. We will prove the estimates for a bigger class H_1^r . We remind the definition of this class. Let for positive integer l $\Delta_t^l f(y)$, $t, y \in [0, 2\pi]$ denote the l -th difference of f with step t , and

$$\Delta_{(t_1, t_2)}^l f(x_1, x_2) = \Delta_{t_2}^l (\Delta_{t_1}^l f(x_1, x_2))$$

be the mixed l -th difference with the step t_j in the variable $x_j, j = 1, 2$. Define

$$H_1^r = \{f : \|f\|_1 \leq 1, \|\Delta_{t_1}^l f(\cdot, x_2)\|_1 \leq |t_1|^r, \|\Delta_{t_2}^l f(x_1, \cdot)\|_1 \leq |t_2|^r, \\ \|\Delta_{(t_1, t_2)}^l f(x_1, x_2)\|_1 \leq |t_1 t_2|^r\},$$

with $l = [r] + 1$. For the embedding of classes W_1^r into H_1^r see [Te].

It proved to be useful in studying approximation of functions with bounded mixed derivative to consider along with the L_p -norms the Besov type norms. Let $V_n(t)$ be the de la Vallée-Poussin polynomials, $t \in [0, 2\pi]$. We define

$$\mathcal{A}_0(t) := 1, \quad \mathcal{A}_1(t) := V_1(t) - 1, \quad \mathcal{A}_n(t) := V_{2^{n-1}}(t) - V_{2^{n-2}}(t), \quad n \geq 2,$$

and for $x = (x_1, x_2), s = (s_1, s_2)$

$$\mathcal{A}_s(x) := \mathcal{A}_{s_1}(x_1)\mathcal{A}_{s_2}(x_2).$$

Consider the convolution operator A_s with the kernel $\mathcal{A}_s(x)$,

$$A_s(f) := f * \mathcal{A}_s,$$

and define the $B_{p,\theta}$ -norm as follows

$$\|f\|_{B_{p,\theta}} := \left(\sum_s \|A_s(f)\|_p^\theta \right)^{1/\theta}, \quad 1 \leq \theta < \infty.$$

It is proved in [T2] that

$$(3.3) \quad \epsilon_m(H_1^r, B_{\infty,2}) \ll m^{-r} (\log m)^{r+1/2}, \quad r > 1.$$

The corresponding proof from [T2] implies also that

$$(3.4) \quad \epsilon_m(H_1^r, B_{\infty,1}) \ll m^{-r} (\log m)^{r+1}, \quad r > 1.$$

Using the obvious estimate

$$(3.5) \quad \|f\|_\infty \leq \|f\|_{B_{\infty,1}}$$

we get the upper estimate in (3.2) from (3.4) and (3.5).

We now proceed to the upper estimate in (3.1). The proof in [T2] was based on the following known estimate of the entropy numbers of octahedron B_1^n in ℓ_∞^n (see [H], [M]).

Lemma 3.1. *The following estimates hold*

$$\epsilon_m(B_1^n, \ell_\infty^n) \ll \begin{cases} m^{-1} (\log(n/m))^2, & 2m \leq n, \\ n^{-1} 2^{-m/n}, & 2m > n. \end{cases}$$

One can use instead of Lemma 3.1 the following result (see [S]).

Lemma 3.2. *The following estimates hold*

$$\epsilon_m(B_1^n, \ell_p^n) \ll \begin{cases} m^{1/p-1}(\log(1+n/m))^{1-1/p}, & m \leq n, \\ n^{1/p-1}2^{-m/n}, & m > n. \end{cases}$$

Then similarly to the proof in [T2] one gets instead of (3.3) the estimate

$$(3.6) \quad \epsilon_m(H_1^r, B_{\infty,2}) \ll m^{-r}(\log m)^{r+1/2}, \quad r > 1 - 1/p,$$

Next, we use the well known corollary of the Littlewood-Paley inequality (see, for instance [Te])

$$(3.7) \quad \|f\|_p \ll \|f\|_{B_{p,2}}, \quad 2 \leq p < \infty.$$

The upper estimate in (3.1) for $2 \leq p < \infty$ follows from (3.6) and (3.7). The corresponding upper estimate for $1 \leq p < 2$ follows from already considered case $p = 2$.

We now proceed to the lower estimates. We begin with the lower estimate in (3.1) for $p = 2$. We use the following simple well known fact on a minimal ϵ -covering (see [P, p.57]). Let a Banach space E be the \mathbb{R}^d equipped with a norm $\|\cdot\|_E$. Denote the corresponding unit ball by B_E . Let $N_\epsilon(F, E)$ be the minimal number of balls of radius ϵ needed for covering F . Then for any body F with existing $vol(F)$ we have

$$(3.8) \quad N_\epsilon(F, E) \geq \epsilon^{-d} \frac{vol(F)}{vol(B_E)}.$$

For a fixed natural number n we consider the orthogonal projector $S_{\Delta Q_n}$ onto $\mathcal{T}(\Delta Q_n)$. Then for any m

$$(3.9) \quad \epsilon_m(W_{1,\alpha}^r, L_2) \geq \epsilon_m(S_{\Delta Q_n}(W_{1,\alpha}^r), L_2 \cap \mathcal{T}(\Delta Q_n)).$$

Next, it is easy to understand that

$$S_{\Delta Q_n}(W_{1,\alpha}^r) = \{f \in \mathcal{T}(\Delta Q_n) : f = F_r(\cdot, \alpha) * \varphi(\cdot), \quad \varphi \in \mathcal{T}(\Delta Q_n), \quad E_{\Delta Q_n}^\perp(\varphi)_1 \leq 1\}.$$

We observe that the operator of convolution with $F_0(x, \alpha)$ defined on $\mathcal{T}(\Delta Q_n)$ induces an orthogonal operator in the space $\mathbb{R}^{2|\Delta Q_n|}$ of Fourier coefficients $A(f)$. Therefore,

$$vol(\{A(f) : f \in S_{\Delta Q_n}(W_{1,\alpha}^r)\})^{(2|\Delta Q_n|)^{-1}} \gg 2^{-rn} (vol((B_{\Delta Q_n}^\perp(L_1)))^{(2|\Delta Q_n|)^{-1}}.$$

Applying Theorem 2.5 we get

$$(3.10) \quad vol(\{A(f) : f \in S_{\Delta Q_n}(W_{1,\alpha}^r)\})^{(2|\Delta Q_n|)^{-1}} \gg 2^{-n(r+1/2)}.$$

Further,

$$(3.11) \quad (\text{vol}\{A(f) : f \in \mathcal{T}(\Delta Q_n), \quad \|f\|_2 \leq 1\})^{(2|\Delta Q_n|)^{-1}} \ll (2^n n)^{-1/2}.$$

Thus, the relations (3.8)–(3.11) imply

$$(3.12) \quad N_\epsilon(W_{1,\alpha}^r, L_2)^{(2|\Delta Q_n|)^{-1}} \gg \epsilon^{-1} 2^{-rn} n^{1/2}.$$

Specifying $m = 2|\Delta Q_n|$ we get from (3.12)

$$\epsilon_m \gg 2^{-rn} n^{1/2} \asymp m^{-r} (\log m)^{r+1/2}.$$

It is clear that the case of general m follows from the special case $m = 2|\Delta Q_n|$, $n \in \mathbb{N}$ which has been considered above. So, we have established the lower estimate in (3.1) for $p = 2$. It implies the corresponding lower estimate for all $p \geq 2$.

Let us prove the lower estimate in (3.1) for $p = 1$. We use the following interpolation inequality for the entropy numbers (see [Pi])

$$(3.13) \quad \epsilon_{2m-1}(W_{1,\alpha}^r, L_2) \leq 2\epsilon_m(W_{1,\alpha}^r, L_1)^{\frac{p-2}{2(p-1)}} \epsilon_m(W_{1,\alpha}^r, L_p)^{\frac{p}{2(p-1)}}$$

with $p > 2$ such that $1 - 1/p < r$. The lower estimate for the left hand side of (3.13) and the upper estimate for $\epsilon_m(W_{1,\alpha}^r, L_p)$, $r > 1 - 1/p$, have already been proved above. Substituting these estimates into (3.13) we obtain the required lower estimate for the $\epsilon_m(W_{1,\alpha}^r, L_1)$. This completes the proof of the lower estimate in (3.1).

We now proceed to the lower estimate in (3.2). Let $M_\epsilon(F, E)$ denote the maximal number of points $x_i \in F$ such that $\|x_i - x_j\|_E \geq \epsilon$, $i \neq j$. The following simple inequality is well known

$$(3.14) \quad N_\epsilon(F, E) \leq M_\epsilon(F, E) \leq N_{\epsilon/2}(F, E).$$

Alike the above case we will carry out the proof for m of a special form: $m = 2|\Delta Q_n|$. Using Theorem 2.5 and the relation (3.8) we will get the following analog of (3.12):

$$(3.15) \quad N_\epsilon(\mathcal{T}(\Delta Q_n)_1^\perp, L_2)^{(2|\Delta Q_n|)^{-1}} \gg \epsilon^{-1} n^{1/2},$$

where

$$\mathcal{T}(\Delta Q_n)_1^\perp = \{f \in \mathcal{T}(\Delta Q_n) : E_{\Delta Q_n}^\perp(f)_1 \leq 1\}.$$

By (3.14) and (3.15) we conclude that there are 2^m polynomials $\{t_j\}_{j=1}^{2^m}$ from $\mathcal{T}(\Delta Q_n)$ such that

$$(3.16) \quad E_{\Delta Q_n}^\perp(t_j)_1 \leq 1, \quad j = 1, \dots, 2^m;$$

$$(3.17) \quad \|t_i - t_j\|_2^2 \gg n, \quad i \neq j.$$

Let $t_j^\perp \in \mathcal{T}(\Delta Q_n)^\perp$, $j = 1, \dots, 2^m$, be such that

$$(3.18) \quad \|t_j - t_j^\perp\|_1 \leq 2.$$

Consider the following collection of functions

$$\varphi_j := (t_j - t_j^\perp)/2, \quad f_j := F_r(\cdot, 0) * \varphi_j(\cdot), \quad j = 1, \dots, 2^m.$$

Then

$$f_j \in W_{1,0}^r, \quad j = 1, \dots, 2^m.$$

We now estimate from below the quantities $\|f_i - f_j\|_\infty$ for $i \neq j$. Consider the inner products

$$a_{ij} := \langle f_i - f_j, \varphi_i - \varphi_j \rangle.$$

On the one hand by (3.18) we have

$$(3.19) \quad a_{ij} \leq 2\|f_i - f_j\|_\infty.$$

On the other hand

$$(3.20) \quad a_{ij} = \sum_k \hat{F}_r(k, 0) |\hat{\varphi}_i(k) - \hat{\varphi}_j(k)|^2 \gg 2^{-rn} \|t_i - t_j\|_2^2.$$

Thus by (3.17), (3.19), and (3.20) we get

$$\|f_i - f_j\|_\infty \gg 2^{-rn} n, \quad i \neq j.$$

Therefore,

$$\epsilon_m(W_{1,0}^r, L_\infty) \gg 2^{-rn} n \asymp m^{-r} (\log m)^{r+1}.$$

This completes the proof of Theorem 3.1.

4. THE DISCRETE L_∞ -NORM FOR POLYNOMIALS FROM $\mathcal{T}(\Lambda)$

We begin with the following conditional statement.

Theorem 4.1. *Assume that a finite set $\Lambda \subset \mathbb{Z}^d$ has the following properties:*

$$(4.1) \quad (\text{vol}(B_\Lambda(L_\infty)))^{1/N} \leq K_1 N^{-1/2}, \quad N := 2|\Lambda|,$$

and a set $\Omega = \{x^1, \dots, x^M\}$ satisfies the condition

$$(4.2) \quad \forall f \in \mathcal{T}(\Lambda) \quad \|f\|_\infty \leq K_2 \|f\|_{\infty, \Omega}, \quad \|f\|_{\infty, \Omega} := \max_{x \in \Omega} |f(x)|.$$

Then there exists an absolute constant $C > 0$ such that

$$M \geq N e^{C(K_1 K_2)^{-2}}.$$

Proof. Using the assumption (4.2) we derive from Theorem 2.6 in the same way as we proved Lemma 2.1 the following volume estimate

$$(4.3) \quad (\text{vol}(B_\Lambda(L_\infty)))^{1/N} \geq C_1 K_2^{-1} (N \log(M/N))^{-1/2}$$

with an absolute constant $C_1 > 0$. Comparing (4.3) with the assumption (4.1) we get

$$M \geq N e^{C(K_1 K_2)^{-2}}, \quad C = C_1^2.$$

Theorem 4.1 is proved.

We now give some corollaries from Theorem 4.1.

Theorem 4.2. Assume a finite set $\Omega \subset \mathbb{T}^2$ has the following property:

$$(4.4) \quad \forall t \in \mathcal{T}(\Delta Q_n) \quad \|t\|_\infty \leq K_2 \|t\|_{\infty, \Omega}.$$

Then

$$|\Omega| \geq 2|\Delta Q_n| e^{Cn/K_2^2}$$

with an absolute constant $C > 0$.

Proof. By Theorem 2.5 (see (2.7)) we have

$$(\text{vol}(B_{\Delta Q_n}(L_\infty)))^{1/N} \leq C(2^n n^2)^{-1/2} \leq Cn^{-1/2} N^{-1/2}$$

with an absolute constant $C > 0$. Using Theorem 4.1 we obtain

$$|\Omega| \geq 2|\Delta Q_n| e^{Cn/K_2^2}.$$

This proves Theorem 4.2.

Remark 4.1. In a particular case $K_2 = bn^\alpha$, $0 \leq \alpha \leq 1/2$, Theorem 4.2 gives

$$|\Omega| \geq 2|\Delta Q_n| e^{Cb^{-2}n^{1-2\alpha}}.$$

Corollary 4.1. Let a set $\Omega \subset \mathbb{T}^d$ have a property:

$$\forall t \in \mathcal{T}(\Delta Q_n) \quad \|t\|_\infty \leq bn^\alpha \|t\|_{\infty, \Omega}$$

with some $0 \leq \alpha < 1/2$. Then

$$|\Omega| \geq C_3 2^n n e^{Cb^{-2}n^{1-2\alpha}} \geq C_1(b, d, \alpha) |Q_n| e^{C_2(b, d, \alpha)n^{1-2\alpha}}.$$

Corollary 4.2. Let a set $\Omega \subset \mathbb{T}^2$ be such that $|\Omega| \leq C_5 |Q_n|$. Then

$$\sup_{f \in \mathcal{T}(Q_n)} \|f\|_\infty / \|f\|_{\infty, \Omega} \geq Cn^{1/2}.$$

Proof. Denote

$$K_2 := \sup_{f \in \mathcal{T}(Q_n)} \|f\|_\infty / \|f\|_{\infty, \Omega}.$$

Then the condition (4.4) of Theorem 4.2 is satisfied with this K_2 . Therefore, by Theorem 4.2

$$2|\Delta Q_n| e^{Cn/K_2^2} \leq |\Omega| \leq C_5 |Q_n|.$$

This implies that

$$K_2 \gg n^{1/2}.$$

Remark 4.2. One can derive from the known results on recovery of functions from the classes W_∞^r (see [T7], [T8]) that for any n there is a set $\Omega_n \subset \mathbb{T}^d$ such that $|\Omega_n| \leq C|Q_n|$ and

$$\sup_{f \in \mathcal{T}(Q_n)} (\|f\|_\infty / \|f\|_{\infty, \Omega_n}) \ll n^{d-1}.$$

REFERENCES

- [B] E.S. Belinskii, *The asymptotic characteristics of classes of functions with restrictions on the mixed derivative (mixed difference)*, Investigations in the theory of functions on several real variables, Yaroslavskii Univ. (1990), 22–37. (Russian)
- [BM] J. Bourgain and V.D. Milman, *New volume ratio properties for convex symmetric bodies in \mathbb{R}^n* , Invent. Math. **88** (1987), 319–340.
- [D] Dinh Dung, *Approximation of multivariate functions by means of harmonic analysis*, Hab. Dissert., Moscow, MGU, 1985.
- [G] E.D. Gluskin, *Extremal properties of orthogonal parallelepipeds and their applications to the geometry of Banach spaces*, Mat. Sb. **136** (1988), 85–96.
- [H] K. Höllig, *Diameters of classes of smooth functions*, in Quantitative Approximation, Academic Press, New York, 1980, 163–176.
- [K1] B.S. Kashin, *On certain properties of the space of trigonometric polynomials with the uniform norm*, Trudy Mat. Inst. Steklov **145** (1980), 111–116; English transl. in Proc. Steklov Inst. Math. **145** (1981), no. 1.
- [KL] J. Kuelbs and W. V. Li, *Metric entropy and the small ball problem for Gaussian measures*, Preprint (1993), 1–33.
- [KT1] B. S. Kashin and V. N. Temlyakov, *On best m -terms approximations and the entropy of sets in the space L^1* , Mat. Zametki **56** (1994), 57–86; English transl. in Math. Notes **56** (1994), 1137–1157.
- [KT2] B.S. Kashin and V.N. Temlyakov, *On estimating approximative characteristics of classes of functions with bounded mixed derivative*, Mat. Zametki **58** (1995), 922–925; English transl. in Math. Notes **58** (1995).
- [KT3] B.S. Kashin and V.N. Temlyakov, *On a norm and related applications*, Mat. Zametki **64** (1998), 637–640.
- [KT4] B.S. Kashin and V.N. Temlyakov, *On a norm and approximation characteristics of classes of functions of several variables*, Metric theory of functions and related problems in analysis, Izd. Nauchno-Issled. Aktuarno-Finans. Tsentra (AFTs), Moscow, 1999, 69–99.
- [M] V.E. Maiorov, *On various widths of the class H_p^r in the space L_q* , Izv. Akad. Nauk SSSR Ser. Mat. **42** (1978), 773–788; English transl. in Math. USSR-Izv. **13** (1979).
- [P] G. Pisier, *The volume of convex bodies and Banach space geometry*, Cambridge University Press, 1989.
- [Pi] A. Pietsch, *Operator Ideals*, VEB, Berlin, 1978.
- [S] C. Schütt, *Entropy numbers of diagonal operators between symmetric Banach spaces*, J. Approx. Theory **40** (1984), 121–128.
- [Sm] S.A. Smolyak, *The ϵ -entropy of the classes $E_s^{\alpha k}(B)$ and $W_s^\alpha(B)$ in the metric L_2* , Dokl. Acad. Nauk. SSSR **131** (1960), 30–33.
- [Te] V. N. Temlyakov, *Approximation of functions with bounded mixed derivative*, Trudy MIAN **178** (1986); English transl. in Proc. Steklov Inst. Math. (1989), no. 1.
- [T1] V. N. Temlyakov, *On estimates of ϵ -entropy and widths of classes of functions with a bounded mixed derivative or difference*, Dokl. Akad. Nauk SSSR **301** (1988), 288–291.
- [T2] V. Temlyakov, *Estimates of the asymptotic characteristics of classes of functions with bounded mixed derivative or difference*, Trudy Mat. Inst. Steklov **189** (1989), 138–168; English transl. in Proc. Steklov Inst. Math. (1990), no. 4, 161–197.
- [T3] V.N. Temlyakov, *Bilinear Approximation and Related Questions*, Trudy Mat. Inst. Steklov **194** (1992), 229–248; English transl. in Proc. Steklov Inst. of Math. **4** (1993), 245–265.
- [T4] V.N. Temlyakov, *Estimates of Best Bilinear Approximations of Functions and Approximation Numbers of Integral Operators*, Mat. Zametki **51** (1992), 125–134; English transl. in Math. Notes **51** (1992), 510–517.
- [T5] V.N. Temlyakov, *An inequality for trigonometric polynomials and its application for estimating the entropy numbers*, J. Complexity **11** (1995), 293–307.

- [T6] V.N. Temlyakov, *On two problems in the multivariate approximation*, East J. Approx. **4** (1998), 505–514.
- [T7] V.N. Temlyakov, *Approximate reconstruction of periodic functions of several variables*, Mat. Sb. (N.S.) **128(170)** (1985), 256–268; English transl. in Math. USSR Sbornik **56** (1987), 249–261.
- [T8] V.N. Temlyakov, *On approximate recovery of functions with bounded mixed derivative*, J. Complexity **9** (1993), 41–59.