

UNCLASSIFIED

AD NUMBER
ADB209449
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Oct 95. Other requests shall be referred to Commander, U.S. Army Medical Research and Materiel Command, Fort Detrick, Frederick, MD 21702-5012.
AUTHORITY
USAMRMC ltr. 10 Mar 97

THIS PAGE IS UNCLASSIFIED

AD _____

GRANT NO: DAMD17-94-J-4317

TITLE: Cytokines, Neovascularization and Breast Cancer

PRINCIPAL INVESTIGATOR: Scott H. Kurtzman, M.D.

CONTRACTING ORGANIZATION: University of Connecticut
Health Center
Farmington, Connecticut 06030-5355

REPORT DATE: October 1995

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick
Frederick, Maryland 21702-5012


DISTRIBUTION STATEMENT: Distribution authorized to U.S.
Government agencies only; Proprietary Information, October 1995.
Other requests for this document shall be referred to Commander,
U.S. Army Medical Research and Materiel Command, Fort Detrick,
Frederick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

19960508 068

DTIC QUALITY INSPECTED 1

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF COLOR PAGES WHICH DO NOT REPRODUCE LEGIBLY ON BLACK AND WHITE MICROFICHE.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1995		3. REPORT TYPE AND DATES COVERED Annual 1 Oct 94 - 30 Sep 95
4. TITLE AND SUBTITLE Cytokines, Neovascularization and Breast Cancer			5. FUNDING NUMBERS DAMD17-94-J-4317	
6. AUTHOR(S) Scott H. Kurtzman, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Connecticut Health Center Farmington, Connecticut 06030-5355			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick Frederick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only; Proprietary Information, October 1995. Other requests for this document shall be referred to Commander, U.S. Army Medical Research and Materiel Command, Fort Detrick, Frederick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Breast cancers can not grow larger than several millimeters in diameter without the ingrowth of new blood vessels (angiogenesis). Recently, Interleukin 8 (IL-8) has been found to be an angiogenic factor. The general hypothesis for this project is that breast cancer cells are capable of producing angiogenic factors (AF) which support tumor growth and metastasis. In order to test our hypothesis, we are examining the expression of IL-8 antigen in human breast tissue using immunohistochemical methods, examining breast cancer cells grown in culture, and establishing an <i>in vivo</i> model in immunodeficient mice. In the first year of this grant, we have examined 29 breast cancers, 8 <i>in situ</i> cancers, and 11 normal breast tissues for IL-8 expression. In addition, blood vessel staining was done using an antibody to vWF. The results confirmed the expression of IL-8 by breast cancer cells. Currently, we are correlating IL-8 expression with angiogenesis. Additional specimens are being processed. We have begun testing the ability of breast cancer cells <i>in vitro</i> to express IL-8. Nine cell lines have been tested. We have found that in breast cancer cells, IL-1 is a potent inducer of IL-8 production. These results support our hypothesis that breast cancer cells produce AF and likely result in tumor growth and metastasis.				
14. SUBJECT TERMS Breast cancer, angiogenesis, cytokines, Interleukin 8			15. NUMBER OF PAGES 20	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

SK Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

SK In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

SK For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

Where In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

Where In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

Where In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 10/26/95
PI - Signature Date

Table of Contents

INTRODUCTION	5
BODY: EXPERIMENTAL METHODS AND RESULTS	6
Specific Aim I- To characterize IL-8 expression in human breast cancer	6
Study IA- To characterize IL-8 antigen distribution in human breast biopsy specimens	6
Study IB- To demonstrate IL-8 mRNA expression in human breast cancer tissues	7
Study IC- To Correlate IL-8 expression with neovascularization in human breast biopsy specimens	7
Summary of Specific Aim I	8
Goals and Obstacles for Year 02	8
Specific Aim II- To characterize IL-8 expression by breast cancer cell lines in vitro	8
Study IIA- To characterize the expression of IL-8 antigen by cultured human breast cancer cell lines	9
Study IIB- To quantify IL-8 expression in cytokine stimulated breast cancer cells	9
Summary of Specific Aim II	10
Goals and Obstacles for Year 02	10
Specific Methods:	10
CONCLUSIONS	12
Bibliography	13
APPENDIX	14
Statement of Work	14
Table 1 Intensity of Staining for IL-8	15
Table 2 Cytokine stimulation of IL-8 Expression in Human Breast Cancer Cells	15
Figure 1 IL-8 Immunohistochemistry	16
Figure 2 Tissue homogenate RIA	17
Figure 3 Anti vWF staining	18
ABSTRACT 1	19
ABSTRACT 2	20

Introduction

Breast cancer is the most common malignancy in women in the United States. The overall survival for all patients with this diagnosis is approximately 70%. Factors associated with poor prognosis include presentation with advanced stage disease or invasion of neural and vascular structures. Recently, it has become clear that patients whose tumors demonstrate a high degree of angiogenesis have a shorter disease free survival, and that these patients are more likely to develop distant metastases [1]. It is also well accepted that tumor can not grow larger than several millimeters in diameter (approximately 10^6 cells) without ingrowth of additional blood vessels [2]. Angiogenesis has been shown to be critical for the growth and metastasis of human breast cancer [3]. Understanding the underlying mechanisms and factors that control angiogenesis will provide not only insights into the basic mechanisms of tumor growth and metastasis, but also will likely provide novel new therapeutic approaches to invasive breast cancer.

Although the relationship between neovascularization and poor outcome in breast cancer is well established, the mechanism of this relationship is not well understood. Undoubtedly, multiple angiogenic factors (AF) are involved in the recruitment and development of tumor angiogenesis [4]. The possible sources of these factors include the tumor cells themselves, tissue cells or infiltrating leukocytes. Previous studies have demonstrated that AF function by inducing vascular endothelial cell proliferation and/or migration. A variety of AF have been recently identified, and include Interleukin 8 (IL-8), Vascular Endothelial Cell Growth Factor (VEGF), and Fibroblast Growth Factor (FGF).

We and others have hypothesized that tumors behave as wounds that do not heal. Normal wounds heal as a result of infiltration of leukocytes and other cells releasing signaling factors which influence the local environment. As a result, a multitude of reactions occur which result in the promotion of wound healing. These reactions include production of fibrin, collagen and other repair factors, as well as an increase in local blood flow to support these activities. This increased blood flow is a result of both local vasodilatation as well as angiogenesis. We hypothesize that in tumors the normal signals that result in healing associated angiogenesis are overamplified, resulting in support for tumor growth and metastasis. Specifically, we hypothesize that the tumor cells themselves, or the infiltrating lymphocytes are capable of producing angiogenic factors.

In order to investigate this, we examined histologic sections of invasive, non invasive (*in situ*) cancers, as well as normal and non malignant breast tissues to determine if IL-8 antigen was present. In addition, we examined breast tumor homogenates to determine if IL-8 was present in the tissues. To further study the ability of breast cancer cells to produce angiogenic factors, breast cancer cells were grown *in vitro* under control and stimulated conditions. The production of IL-8 in the cell culture supernatant was measured.

Our Statement of Work is presented in the Appendix. Our original target was to complete all the immunohistochemical data gathering during the first year, and develop the *in vitro* model during the second year. We have completed accrual of the immunohistologic specimens and are in the process of analyzing our results. Our goal for the *in situ* hybridization was to directly demonstrate the production of IL-8 by the breast cancer cells. However, our early pilot *in vitro* data so clearly showed production of IL-8

by breast cancer cells, that we chose to pursue this direction first, and postpone the *in situ* hybridization. Thus, we have been quite successful in meeting the timeline specified.

Body: Experimental Methods and Results

Specific Aim I- To characterize IL-8 expression in human breast cancer

Introduction: The goals of Specific Aim 1 are to extend and expand our preliminary observations by: 1) demonstrating the presence of IL-8 in invasive breast cancer but not in non invasive cancer or fibrocystic disease, 2) establishing the production of IL-8 in invasive breast cancer by measuring mRNA; and 3) correlating neovascularization with IL-8 expression and patient outcome. These experiments will support our hypothesis that breast cancer cells produce angiogenic factors. Further studies will examine the regulation of this expression (Specific Aims II and III).

Study IA- To characterize IL-8 antigen distribution in human breast biopsy specimens

Experimental Approach: The purpose of Study IA is to confirm and extend our preliminary observations regarding IL-8 expression in invasive cancer, *in situ* cancer, and non malignant breast tissue. Tissue samples were obtained from patients undergoing breast biopsy. Where possible, clinical information was obtained, which included patient age, TNM stage, histologic grade, and survival information.

In order to further quantify the expression of IL-8 by breast cancer cells, we assayed the levels of IL-8 in tissue homogenates from tissue samples. Homogenates were also made from non malignant breast tissue when this was available. Although not included in the original Statement of Work, it was felt that this data might be used to quantify the level of expression seen in the tissues.

Results

Immunohistochemistry

The results of the immunohistochemical evaluation are presented in Table 1. and Figure 1. Only those specimens examined during Year 01 of the grant and reviewed by the PI's are included in the Table. We have found that for consistency, slides are best processed and read in batches, with standards for staining being used. We have accumulated approximately 30 more specimens which are being processed. This will allow us to reach the target we set in the original proposal.

As observed in our preliminary data, IL-8 antigen expression is clearly seen in the majority of human breast cancer specimens evaluated. There is a lesser degree of inhomogeneous staining seen in the *in situ* cancer specimens. While some staining was

seen in the non malignant breast tissues, the pattern was luminal rather than cellular in contradistinction to that seen in the cancers.

Tissue Homogenates

Tissue homogenates were prepared, or obtained on 102 specimens (these specimens do not necessarily correspond with the specimens used for immunohistochemical analysis). IL-8 levels were determined by RIA for 52 cancers, 6 normal adjacent breast tissue, 1 fibroadenoma and 1 case of DCIS. The results of 15 representative specimens are presented as ng IL-8/mg protein (see Figure 2). As can be seen, elevated levels of IL-8 are seen in tissue homogenates obtained from some breast cancer specimens, but not from the benign tissues. The level in the single case of DCIS appears to be intermediate between the benign tissues and the invasive cancers. This is being further explored with the remainder of the samples.

Study IB- To demonstrate IL-8 mRNA expression in human breast cancer tissues

Experimental Approach: The purpose of Study IB is to establish the exact source of the IL-8 present in breast cancer specimens. Although the presence of IL-8 antigen on the surface of breast cancer cells in human tissue specimens strongly suggests that the breast cancer cells are producing the AF, direct identification using *in situ* RNA methods would further establish that the breast cancer cells are the source of the AF.

However, through the use of *in vitro* techniques using breast cancer cells in culture we will also be able to determine the source of AF (see Specific Aim II). Our early data suggested that breast cancer cells in culture can indeed produce IL-8. Therefore, for strategic reasons, based on the availability of breast cancer cell cultures, and the start up time anticipated for *in situ* hybridization, it was decided that this portion of the experiment be postponed to Year 02 or Year 03, depending on the *in vitro* results.

Study IC- To Correlate IL-8 expression with neovascularization in human breast biopsy specimens

Experimental Approach: Our central hypothesis is that breast cancer cells produce AF that result in neovascularization. Study IA established the presence of IL-8 in breast cancer tissue, the purpose of this Study is to correlate IL-8 expression with neovascularization. Breast tumor specimens obtained in Study IA were stained with antibody to vWF or CD-31 in order to identify blood vessels (Figure 3). In our experience the anti CD-31 better visualized the vessels and was associated with lower background staining thus facilitating vessel counting. For these reasons, anti CD-31 staining will be used for all future studies. Vessels were counted in the manner described by Weidner *et al.* Tumor specimens were reviewed by us and a staff pathologist using a dual head microscope. vessels were identified by specific staining. We did not require that a lumen be seen. The slides were scanned, and the area of highest vessel density selected. Both reviewers counted vessels.

Results

Blood vessel staining was identified in the breast cancer specimens. The level of angiogenesis varied between samples. We are currently correlating IL-8 antigen staining with angiogenesis. Clearly, relating the clinical data to the staining patterns will be critical to understanding the regulation of breast cancer growth.

Summary of Specific Aim I

Our results support our hypothesis. Specific staining for IL-8 antigen was identified in the breast cancer specimens. Not surprisingly, there is a lesser extent of IL-8 antigen in the non malignant breast tissue, although the pattern of staining is different. Angiogenic factors are a normal constituent of wound healing. Therefore, it is not unexpected that normal cells express IL-8. However, it is the exaggerated response and the regulation of the angiogenesis that is the focus of this project. Specific Aim II is directed towards understanding this regulation.

Goals and Obstacles for Year 02

Acquiring adequate tissue samples proved more difficult than anticipated. At this point we have established a more reliable tissue bank and access to more breast tissue. This has been organized in such a manner to allow us to obtain corresponding clinical information. This will allow us to better correlate the IL-8 expression with clinical information and angiogenesis (see Study IC). Our goal is to complete accumulation of tissue samples in the early part of Year 02, and complete both the IL-8 as well as the vessel staining. Correlation with clinical data should be completed during this year.

Specific Aim II- To characterize IL-8 expression by breast cancer cell lines in vitro

The studies in Specific Aim I were directed at establishing the presence and correlation of IL-8 with neovascularization in invasive breast cancer, *in situ* breast cancer and fibrocystic patients. These studies support our hypothesis on the role of IL-8 in neovascularization in breast cancer. We will clearly need to directly demonstrate that tumor derived IL-8 is involved in neovascularization and to identify those factors involved in regulation of this process. To achieve this we propose to utilize a xenograft model of IL-8 POS and IL-8 NEG human breast tumor cells in immunodeficient mice.

In our proposed *in vivo* model it is anticipated that IL-8 expression is regulated by a variety of factors (e.g. cytokines (IL-1/TNF), estrogens, etc.). Since the *in vivo* model is more complicated than an *in vitro* system, we first developed a parallel *in vitro* system to study the regulation of IL-8 expression by the cultured tumor cells. Selected studies were carried out to determine the ability of stimulants of IL-8 expression to regulate IL-8 expression from both the IL-8 POS and IL-8 NEG human breast cancer cells *in vitro*. Clearly, the first step in developing this model is identifying and characterizing IL-8 POS and IL-8 NEG human breast cancer cell lines. Initially, two well established cell lines, MCF-7 and BT-20 were used for these studies. These preliminary experiments were

expanded to include 5 additional breast tumor cell lines, and two lines derived from normal breast tissue. Thus, Specific Aim II will provide the foundations for the *in vivo* model which will allow us to directly test our hypothesis.

Study IIA- To characterize the expression of IL-8 antigen by cultured human breast cancer cell lines

Experimental Approach: The breast cancer lines were obtained from the ATCC catalogue. The normal cell line HMEC was obtained from Clonetics (Clonetics Corp., San Diego, CA) Each cell line is grown in specific media as recommended by ATCC. Cell growth for the individual tumor lines is relatively constant between passages. The lines have distinctive morphological appearances which are confirmed visually to ensure healthy growth as well as identity. Cells are passed upon reaching confluence using trypsin.

In order to directly examine the ability of the breast cancer cells to produce IL-8, we performed immunocytochemical analysis. For these immunocytochemical studies, 10^4 breast cancer cells were plated on tissue culture slides. In selected studies, these cells were treated with the stimulants as described below. The cells become adherent after 24-48 hours. The slides are then exposed to 1cc of 10% formalin at 4°C for at least 2 hours. They are then stained for IL-8 as described in Study IA except that they do not go through the serial alcohol deparaffinization steps.

Study IIB- To quantify IL-8 expression in cytokine stimulated breast cancer cells

Introduction: Cytokine production by tumor cells has been demonstrated in multiple cancer models. Although in Study IIA we showed IL-8 antigen to be present at the cellular level in cultured breast cancer cells, we expect that IL-8 expression is influenced by other mediators either in an autocrine or paracrine fashion. Therefore we treated cultured breast cancer cells with a variety of agents to examine their ability to regulate AF expression. Completion of this study will allow us to proceed to *in vivo* experiments using stimulants, and blockers of cytokine expression and hormonal manipulation.

Experimental Approach: Cells were prepared as described above, and plated on 6 well sterile culture plates. An adequate number of cells was plated in order to produce confluence at 24 hours. The number of cells used was recorded and varied between cell lines. Data is corrected for 10^6 cells. After 24 hours, the culture media was removed, and replaced with the test material in 3 cc of sterile media. Complete, serum containing culture media was used as control. The stimulants used were IL-1 alpha and beta (10 ng/ml), LPS 5ug/ml, TNF alpha and beta (100U/ml), TNF 500U/ml and fibrin. The cell cultures were exposed for 4, 24 or 48 hours. All incubations were carried out at 37°C , under 5% CO_2 . Cells and supernatants are harvested as described above and frozen at -70°C until analysis. Aliquots were analyzed for IL-8 by RIA.

Results

The results of Study IIB are presented in Table 2 and expressed as the stimulated index (ratio of stimulated over media control). The basal (unstimulated) levels in ng/ml are also included in the table. Baseline IL-8 expression was demonstrated by many of the cell lines. The response to the stimulating factors was somewhat variable between the cell lines. However, IL-1 alpha and beta were consistently found to be a potent inducers of IL-8 expression.

Summary of Specific Aim II

In Specific Aim II, we have demonstrated directly the production of IL-8 by breast cancer cells. Further, we have begun to explore the regulation of this expression. Clearly, cytokines such as IL-1 play a significant role in inducing angiogenic factor expression. This information will allow us to develop our *in vivo* model, and plan the studies that will explore inhibition of tumor growth in the immunodeficient mouse model.

Goals and Obstacles for Year 02

Our *in vitro* work is well ahead of schedule, and has been quite successful. As a result, we have been able to expand the number of cell lines being tested. In addition, if time and resources allow, we will expand our survey of angiogenic factors.

Co-culture experiments are also planned. We are considering the feasibility of co-cultures with human vascular endothelial cells.

Specific Methods:

IL-8 Immunohistochemistry

Immunohistochemistry is performed by indirect immunoperoxidase staining. 8 um paraffin embedded tumor sections are deparaffinized in xylene and rehydrated in graded alcohol (100%, 95% and 50%). To inhibit endogenous peroxidases, the sections are immersed in methanol containing 0.01% hydrogen peroxide at 4⁰ C for 15 minutes. Sections are allowed to air dry and then blocked with normal goat serum (Vector, Burlington, CA) at room temperature (RT) for 1 hour. The sections are then washed 3 times with phosphate buffer saline (PBS, pH 7.4) and affinity purified rabbit or chicken anti-human IL-8 (1/500 - 1/9000 dilution in PBS with 0.5% bovine serum albumin), chicken anti-human IL-8 preabsorbed with recombinant human IL-8 specificity control, or buffer control are applied to the sections overnight at 4⁰ C.

After the incubation period, biotinylated goat anti rabbit, or rabbit anti-chicken antibody (Biotin SP Affinipure F(ab)₂ fragment, (Jackson ImmunoResearch Labs, Inc., West Grove, PA); 1/100 dilution in PBS with 0.5% BSA is applied to the sections and allowed to incubate for 1 hour at RT. Sections are washed 3x with PBS in between each

of the following steps. HRP-streptavidin (Zymed, San Francisco, CA) at a 1/250 dilution in PBS is applied to the sections and incubated at RT for 1 hour. The sections are incubated sequentially with 3 amino-9-ethylcarbazole in 0.1M sodium acetate buffer (pH 5.0) and 0.03% H₂O₂ for 30 minutes at RT. Sections are then counterstained in Mayer's hematoxylin (Sigma, St. Louis, MO) for 10 minutes, extensively washed in H₂O₂, and dipped in dilute ammonium hydroxide and mounted in crystal mounting solution (Biomedex, Foster City, CA).

The slides were reviewed by us and by a staff pathologist to confirm the histologic diagnosis, and to grade the degree of staining. Staining was graded as 0-4+. Nonspecific, species matched IgG was used as a negative control.

Tissue Homogenates

Tumor and control tissue samples were stored in a -70°C freezer until processing. They were then thawed, weighed, placed in 1 ml of phosphate buffer saline (PBS) per 0.25 grams of tumor or control tissue, and homogenized in a tissue homogenizer. Triton (0.01 percent final concentration) was then added to the supernatants, which were then re-homogenized. The homogenate was centrifuged twice at 4°C at 10,000 g. The supernatants were then aliquoted and frozen at -20°C.

In addition, residual tissue homogenates from specimens sent to pathology for ER/PR analysis were obtained. Aliquots were frozen at -70°C and then analyzed by RIA for IL-8, and for protein level using the Bradford reagent (see above). ER and PR values were available for these samples and will be correlated with IL-8 levels.

Cytokine Analysis

Cell culture supernatants are analyzed by an IL-8 specific RIA developed in our laboratory. Samples (100 ul) are incubated (25°C, 2h) with chicken anti-IL-8 (100 ul) diluted (1:4,000) in RB buffer (1% BSA, 0.1% Triton X-100 in PBS). Chicken antibody to Interleukin 8 (IL-8) was prepared by intramuscular injection of 100 ug of recombinant human IL-8 (77 amino acids, PreproTech Inc., Rocky Hill, NJ) prepared in Hunger's P₄ Titer Max (CytRx Corp., Norcross, GA). Egg yolks containing antibody were processed as previously described. Antibody titer and specificity were assessed by double immunodiffusion and immunoelectrophoresis, respectively. Human ¹²⁵I-IL 8 (NEN products, Boston, MA) diluted in RB buffer (70-80,000 CPM/ml) is added (100 ul) and the reaction mix incubated overnight (4°C). Immune complexes are precipitated by adding (100 ul) affinity purified goat anti-chicken IgG coupled microspheres (Kirkegaard & Perry Lab Inc., Gaithersburg, MD) diluted in RB buffer (1:20). After incubation (25°C, 2h) beads are pelleted (2500g for 15 min.), blotted and counted (gamma counter 1 min.). Samples are quantified by reference to a standard curve constructed using rIL-8 standards (0.039-10 ng/ml). Samples were assayed in duplicate and results are expressed as the mean +/- S.D. All resulting data are expressed as pg of AF per 10⁶ cells. Determination of statistical significance for AF expression between various culture conditions and times will be done by ANOVA analysis (JMP software program). For ease of analysis, the data were also expressed as cytokine index or CI = cytokine concentration (molar) divided by antagonist concentration (molar) x 10³.

Conclusions

We have been extremely successful in pursuing the studies outlined in the application. Our immunohistochemical results to date have confirmed our preliminary data which showed IL-8 antigen expression in human breast cancer. Early neovascularization data confirms the presence of neovascularization in the breast cancer specimens. Clearly, correlating neovascularization with IL-8 expression will provide an important link in the data. Likewise, correlation with clinical data will be an important step in understanding the significance of our observations.

Our *in vitro* data has been extremely encouraging. Not only have we been able to validate our preliminary data demonstrating IL-8 production by tumor cells, but by expanding our number of cell lines and stimulants, we have shown that different cell lines respond differently to the various stimulants. IL-1 emerges a consistent and strong stimulant of IL-8 expression. This information will be key in establishing the *in vivo* model in Years 03 and 04. We have submitted our work to the Society of Surgical Oncology for presentation March 1996 (see Abstract 1 in Appendix).

We feel that the work done in the first year of this project has supported our central hypothesis. Breast cancer cells are capable of producing angiogenic factors. The implication of this observation is that development of agents that control of angiogenesis will likely lead to treatment resulting in decreased tumor growth and metastasis.

During the latter part of the first year, we developed a more efficient and inexpensive way to obtain breast tumor specimens. This redistribution of resources allowed us to expand the *in vitro* model to include a number of additional breast cell lines. We have observed variability between the cell lines in the ability of cytokines to regulate IL-8 expression. In addition, the tumorigenicity of the various cell lines may vary in the immunodeficient mouse model. Therefore, having a larger group of cell lines to choose from will increase our ability to develop the *in vivo* model.

Further, although this project was directed at examining the role of IL-8 in particular, a variety of other AF exist. If time and resources allow, we plan to examine other angiogenic factors such as VEGF and basic FGF in our system. Our original proposal also did not look at the role of hormones or fibrin in our system. We are currently seeking additional support to extend our experiments in this area. Additionally, we have submitted a proposal on a related project examining the role of fibrin on angiogenic factors.

Our early successes have led to several "spin-off" projects. We are beginning to study other angiogenic factors such as VEGF in both the immunohistochemical studies as well as in the *in vitro* model. We have also generated preliminary data examining antagonists such as IL-1 receptor antagonists (IL-1 RA). If our hypotheses are correct that cytokines such as IL-1 are inducing expression of angiogenic factors, then the role of IL-1 RA will prove important.

Other workers in our laboratory have started similar projects similar to this one in the fields of prostate cancer and colo-rectal cancer. Preliminary work in prostate cancer has been submitted to the American Urologic Association (see Abstract 2 in Appendix). Additional independent funding is being sought for these projects.

Bibliography

1. Weidner, N, Semple, JP, Welch, WR, and Folkman, J, Tumor angiogenesis and metastasis--correlation in invasive breast carcinoma. *New England Journal of Medicine*, 1991. 324 (1): p. 1-8.
2. Weidner, N, Folkman, J, Pozza, F, et al. , Tumor angiogenesis: a new significant and independent prognostic indicator in early-stage breast carcinoma [see comments]. *Journal of the National Cancer Institute*, 1992. 84 (24): p. 1875-87.
3. Gasparini, G and Harris, AL, Clinical importance of the determination of tumor angiogenesis in breast carcinoma: Much more than a prognostic tool. *Journal of Clinical Oncology*, 1995. 13 (3): p. 765-782.
4. Folkman, J and Klagburn, M, Angiogenic factors. *Science*, 1987. 235 : p. 442-447.

Appendix

Statement of Work

Task 1 Year 1- To characterize IL-8 expression in human breast cancer

Study IA- To characterize IL-8 antigen distribution in human breast biopsy specimens

Study IB- To demonstrate IL-8 mRNA expression in human breast cancer tissues

Study IC- To Correlate IL-8 expression with neovascularization in human breast biopsy specimens

Task 2 Year 2- To characterize IL-8 expression by breast cancer cell lines in vitro

Study A- To characterize the expression of IL-8 antigen by cultured human breast cancer cell lines

Study B- To quantify IL-8 expression in cytokine stimulated breast cancer cells

Study C- To quantify IL-8 expression in co cultures of MCF-7 and BT-20 cells

Task 3. Years 3-4- To characterize IL-8 antigen expression and neovascularization in human breast cancer cells grown as tumors in nude mice

Study A- To demonstrate IL-8 antigen expression in subcutaneously implanted human breast cancers

Study B- To establish a dual tumor model using MCF-7 and BT-20 cells

Table 1

Intensity of Staining for IL-8

	0	1+	2+	3+	4+
Invasive cancer	3	2	8	5	1
In situ cancer			1	2	
Benign breast	1	3	1	1	

Table legend: Results of immunohistochemical staining for IL-8 on 29 samples. Rating scale: 0= no staining; 1+ = minimal staining; 2+ = moderate staining; 3+ = strong staining; 4+ = intense staining.

Table 2

Cytokine stimulation of IL-8 Expression in Human Breast Cancer Cells

	MCF-7	BT-20	MDA-231	ZR-75-1	T-47D	HBL-100	HS-578
Media	1.00	1.00	1.00	1.00	1.00	1.00	1.00
TNF- α	1.30	1.84	12.20	0.73	0.91	0.32	9.80
TNF- β	1.00	1.06	1.59	0.68	0.67	0.09	4.10
IL-1 α	0.95	1362	127	107	3.36	0.12	124
IL-1 β	0.81	1292	144	137	1.00	0.24	119

Basal IL-8 Expression (ng/ml)*

Media	0.45	0.31	1.5	0.15	0.1	0.15	0.40
-------	------	------	-----	------	-----	------	------

Table 2: Stimulation index (IL-8 expression in ng/ml stimulated/ IL-8 expression in ng/ml in media control). *The lower portion of the table lists the basal (unstimulated media control) IL-8 expression in ng/ml

Figure 1




Figure 2




Figure 2: RIA for IL-8 demonstrating expression of IL-8 antigen in a number of breast cancers, with no expression seen in non malignant tissue. DCIS appears to be intermediate

Figure 3




Figure 3: Demonstration of anti vWF(A) and control IgG (B) immunohistochemistry for identification of vessels in breast cancer specimen. As can be seen in this figure, both large and small vessels are easily visualized using the anti vWF antibody.

ABSTRACT 1

CYTOKINE REGULATION OF ANGIOGENESIS FACTOR EXPRESSION IN HUMAN BREAST CANCER. SH Kurtzman, L Miller, K Anderson Y Wang, DL Kreutzer, Univ of CT Dept of Surgery, Farmington, CT

Currently, little is known about the regulation of angiogenic factors (AF) expression in breast tumor cells. To fill this gap in our knowledge, we hypothesize that in breast cancer 1) breast cancer cells produce AF, and 2) this expression is regulated by cytokines. In order to investigate this 7 human breast cell lines were grown in continuous culture and exposed to the cytokines IL-1 α , IL-1 β , TNF α , TNF β or control media. Resulting cell culture supernatants were harvested and assayed by RIA for IL-8, a known AF. The results are presented in Table 1 as the ratio of the levels of IL-8 in the stimulated over control condition.

Table 1	MCF-7	BT-20	MDA-231	ZR-75-1	T-47D	HBL-100	HS-578
Media	1.00	1.00	1.00	1.00	1.00	1.00	1.00
TNF- α	1.30	1.84	12.20	0.73	0.91	0.32	9.80
TNF- β	1.00	1.06	1.59	0.68	0.67	0.09	4.10
IL-1 α	0.95	1362	127	107	3.36	0.12	124
IL-1 β	0.81	1292	144	137	1.00	0.24	119

These data clearly indicate that 1) IL-1 appears to be an extremely potent and consistent inducer of IL-8 in a number of breast cancer cell lines and 2) different tumor cell lines respond differently to individual cytokines (high vs. low responders.) Thus, human breast cancer cells are capable of producing AF in response to cytokines such as IL-1 and TNF. Since IL-1 and TNF are produced by tissue cells and leukocytes within the tumor microenvironment, they likely contribute to angiogenesis and resulting tumor growth and metastasis. This information will allow us to design future therapeutic agents targeted at AF expression and their receptors that will aid in the treatment of patients with breast cancer.

**To be presented at the Society of Surgical Oncology
 Spring Meeting, Atlanta, GA March 1996**

Abstract 2

1996 ANNUAL MEETING - ORLANDO, FL
DEADLINE FOR SUBMITTING PAPERS IS OCTOBER 13, 1995

For office use

Do not fold this form. Mail the original and 2 copies in a flat envelope with cardboard to:

Dr. William H. Turner, Jr., AUA Meeting Abstracts, 1120 N. Charles Street, Baltimore, MD 21201-5559

See attached sheet for complete instructions before typing abstract. Each paper submitted by a non-AUA member must be sponsored by an AUA member.

TYPE WITHIN BLOCK - NO MARGINS

TOPIC: (See Topic List)

Choice of Category: 8

Choice of Session:

☒ Research ☐ Clinical

SOURCE OF FUNDING:

UConn Health Center

(NIH, NCI, Industry, Institutional Grants, Foundations, etc.)

CHOICE OF PRESENTATION:

☐ Poster only ☒ Podium or Poster

DIRECT CORRESPONDENCE TO:
(Please type)

FERNANDO FERRER, M.D.

UConn Health Center

Address DIVISION OF UROLOGY

263 FARMINGTON AVENUE

FARMINGTON, CT

City/State

USA 06030-3955

Country

(203) 679-3430

Zip Code

Telephone #

(203) 679-1276

Fax #

STATUS OF PRESENTER

☐ Private practice: ☐ Clinical faculty:☐ Full-time academician: ☒ Resident: ☐ PhD.

UConn Health Center

Name of Institution

☐ AUA member

(category)

☒ Non-member☐ Urologist: ☐ Non-Urologist

UROLOGY RESIDENT

(specialty)

NON-MEMBERS:

APPLICATION FOR NON-MEMBER MUST BE
SPONSORED BY A MEMBER OF THE AUA.
SIGNATURE OF MEMBER IS REQUIRED.

Signature of sponsoring AUA member

PETER C. ALBERTSEN, M.D.

Please print name of sponsoring AUA member

CYTOKINE REGULATION OF ANGIOGENESIS FACTORS IN HUMAN

PROSTATE CANCER. Fernando A. Ferrer, Lauri J. Miller, Ramez I.

Andrawis, Peter C. Albertsen, Scott H. Kurtzman, Donald L. Kreutzer,
Farmington, CT (Presentation by Dr. Ferrer)

Introduction Recently, we have demonstrated immunohistochemically the presence of the potent angiogenesis factor (AF), Interleukin-8 (IL-8) in human prostate cancer (PC). In these studies IL-8 antigen was noted to be consistently associated with tumor cells. Based on this observation, we hypothesized that PC tumor cells 1) produce AF, including IL-8, and 2) that their expression is regulated by cytokines present in the tumor microenvironment. To test this hypothesis, a human PC tumor cell line was exposed to the key cytokines Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) in vitro and the resulting culture supernatants were assayed for IL-8 expression.

Methods The PC cell line DU-145 was grown in continuous culture and exposed to cytokines IL-1 α , IL-1 β , TNF α , TNF β (10ng/ml), as well as a control media. Cell culture supernatants were harvested at 24 Hrs and analyzed by radioimmunoassay for IL-8. The resulting data was initially expressed as ng/ml and then normalized as a stimulation index (S. I.) which represents the ratio of stimulated divided by control IL-8 levels.

Results Our results indicate that IL-1 α and IL-1 β are potent and consistent inducers of IL-8 in the DU-145 PC cell line. Surprisingly, TNF α and TNF β induced little to no significant IL-8 expression (see Table).

Stimulant	Media	TNF - α	TNF - β	IL - 1 α	IL - 1 β
	n=9	n=9	n=9	n=9	n=8
S. I. mean (SEM)	1.00 $\pm$ (0.23)	1.90 $\pm$ (0.48)	1.13 $\pm$ (0.34)	16.73 $\pm$ (5.38)	11.85 $\pm$ (2.12)
p value $\leq$		0.1050	0.7540	0.0100	.0001
IL-8 range (ng/ml)	0.3 - 4.3	1.1 - 4.8	0.7 - 1.9	6.0 - 40.3	6.3 - 44.8

Conclusion In vitro, cytokines such as IL-1 are capable of regulating IL-8 expression in human PC. Cytokines such as IL-1 are known to exist within the tumor microenvironment and by inducing AF, likely contribute to angiogenesis, tumor growth and metastasis. Further in vivo studies will clarify the clinical significance of this observation and suggest treatment strategies targeted at AF and their inducers.

WORD COUNT
(Limited to 300)

260

COMPLETE THE QUESTIONS BELOW:

- Multiple abstracts by the same authors on the same topic will have low priority for presentation at the meeting. These submissions may be combined or denied. Authors would benefit substantially from combining data into a single, meaningful abstract. Is this such a presentation? Yes ☐ No ☒
- Papers accepted for publication or previously published will have low priority for presentation at the meeting. Has this paper been accepted/published? Yes ☐ No ☒
- The American Urological Association requests all speakers at the Annual Meeting disclose any financial support from or business affiliation with industry, or other conflicts of interest, in connection with a product or technique described in their presentation. This disclosure is to be made at the beginning of the presentation. I have read the Disclosure Requirements of the American Urological Association's Annual Meeting. ☒ I have no business affiliation with or relevant financial support from industry in conjunction with this presentation. ☐ I have the following business affiliation or financial support in conjunction with my presentation.

Page 20



DEPARTMENT OF THE ARMY

U.S. ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

10 Mar 97

MEMORANDUM FOR Administrator, Defense Technical Information
Center, ATTN: DTIC-OCF, Fort Belvoir,
VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical report written for Contract Number DAMD17-94-J-4317. Request the limited distribution statement for Accession Document Number ADB209449 be changed to "Approved for public release; distribution unlimited." This report should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322.

FOR THE COMMANDER:


GARY R. GILBERT
Colonel, MS
Deputy Chief of Staff for
Information Management

Completed
2-8-2000
B.W.