

UNCLASSIFIED

AD NUMBER
ADB254454
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Aug 99. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St, Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 23 Aug 2001

THIS PAGE IS UNCLASSIFIED

AD_____

Award Number: DAMD17-99-1-9039

TITLE: Antisense Down-Regulation of Metallothionein:
A Therapeutic Approach for Prostate Cancer

PRINCIPAL INVESTIGATOR: Asim B. Abdel-Mageed, Ph.D.

CONTRACTING ORGANIZATION: Tulane University School of Medicine
New Orleans, Louisiana 70112-2699

REPORT DATE: August 1999

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Aug 99). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20000608 154

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-99-1-9039

Organization: Tulane University Medical Center

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Nmanga chuan Monks
5/25/00

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1999		3. REPORT TYPE AND DATES COVERED Final (15 Jan 99 - 14 Jul 99)
4. TITLE AND SUBTITLE Antisense Down-Regulation of Metallothionein: A Therapeutic Approach for Prostate Cancer			5. FUNDING NUMBERS DAMD17-99-1-9039	
6. AUTHOR(S) Asim B. Abdel-Mageed, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Tulane University School of Medicine New Orleans, Louisiana 70112-2699			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Aug 99). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Despite significant research in the field of prostate cancer (PCa), the disease remains elusive in terms of its etiology, pathophysiology and clinical management. Identification of new genotypic and phenotypic markers underlying initiation, promotion and endocrine therapy failure would be of paramount significance. We have shown that metallothionein (MT) can induce mitogenic response in tumor cells and that its inhibition can lead to growth arrest and induction of apoptotic cell death. The objective of this training grant was to establish an antisense MT adenovirus construct and test its efficacy on growth inhibition and induction of apoptosis <i>in vitro</i> and <i>in vivo</i> . Due to experimental difficulties and the short time-frame specified for this award (6 months) we were able only to generate the antisense MT adenoviral construct. The experiments testing the efficacy of this novel construct are ongoing. The significance of this construct lies on the fact that MT is a mitogenic protein and its abundance, as shown in many human primary tumors, is associated with development of drug resistance and poor prognostic outcome. Thus, it is anticipated that this novel antisense construct will have a significant impact on growth inhibition and enhancement of efficacy of chemotherapeutic agents.				
14. SUBJECT TERMS Prostate Cancer Metallothionein, antisense, adenovirus rector, gene therapy, prostate cancer			15. NUMBER OF PAGES 16	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


✓ ____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

✓ ____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 8/12/99
PI - Signature Date

(3) Table of Contents

Pages

1.	Front Page	
2.	Report Documentation Page (SF 298)	2
3.	Foreword	3
4.	Table of Contents	4
5.	Introduction	5
6.	Body	6
7.	Key Research Accomplishments	13
8.	Reportable Outcome	8
9.	Conclusions	13
10.	References	14
11.	Appendices	15

(5) Introduction:

Prostate cancer (PCa) still poses question regarding its etiology, pathology, pathogenesis and clinical management. It has been estimated that there may be quarter of a million cases of PC worldwide (1) and the incidence of PC is projected to increase significantly (2). While earlier reports estimated that PC account for 35,000 deaths annually in Europe (3) and 38,000 in the USA (4), recent reports demonstrated that the difference between the incidence rate and the death rate is striking; in 1996 there were 317,000 newly diagnosed PC cases and 41,400 deaths in the USA compared to the 1995 figures of 244,000 and 40,400, respectively (5). However, it should be noted that there have been no major changes in mortality and no major advances in treatment parallel with the current rise in diagnosis since the introduction of androgen down-regulation regimen. Recent figures on prostate cancer suggest that the incidence vary considerably by race and age (6, 7). It has been shown that the mortality rates for African-American men were 2.2 times higher than those for white men. In more details, addition, the mortality rate among African-Americans has increased 68% between 1960-62 and 1990-92, and in recent years, have increased approximately 4% annually (8).

Despite significant research in the field of PCa, the disease remains elusive in terms of its pathogenesis and clinical management. Identification of new genotypic and phenotypic markers and characterization of molecular mechanisms underlying promotion of PCa to metastatic phenotypes and endocrine therapy failure would be of paramount significance. Several lines of evidence demonstrate a relationship between metallothionein (MT) expression in various types of human primary tumors and disease progression, metastasis, and poor prognostic outcome. MTs are a set of ubiquitous, low molecular weight (6-7kD) aromatic amino acid-free proteins that have 20 reduced cysteine residues which can chelate seven bivalent heavy metal ions through mercaptide bonds (9). In humans, MTs are encoded by a family of genes located at 16q13 (10,11) that contains 10 functional and 7 nonfunctional isoforms (12,13). The gene sequence and expression of all the functional human MT genes have been identified and well characterized: MT-IIA (14), MT-3 (15), MT-4 (16), MT-IA (17), MT-IB (18), MT-IE, MT-IF and MT-IG (19), and MT-IH and MT-IX (20).

MT overexpression has been shown to be associated with disease progression, metastasis, and poor prognostic outcome in a number of human primary tumors, including malignant melanomas (21), ovarian tumors (22), testicular germ cell tumors (23), bladder transitional cell carcinomas (24), and bladder cancer and renal cell carcinoma (25). However, the functional role of MT in tumor cell growth is poorly understood. To answer this question, we have recently demonstrated in breast carcinoma MCF-7 cells that MT overexpression enhances cellular proliferation and *c-myc* gene expression and suppresses *c-fos* and *p53* transcripts, whereas antisense down-regulation of MT induces growth arrest and apoptotic cell death (26). We also showed that the activation of transcription factor NF-kappa B and its interaction with MT may be required for the MT-mediated mitogenic response (27).

In our laboratory, we demonstrated the isoform-specific expression of MT in PCa and further examined the extent to which they modulate disease progression and metastasis. Specifically, our preliminary data using RT-PCR have shown enhanced MT-IIA, MT-IE, and MT-IF gene expression in human PCa biopsy samples when compared to tissues from benign prostatic hyperplasia (BPH), suggesting that at least the aberrant expression of specific MT isoform(s) may be associated with growth promotion and metastatic disease. This observation was confirmed by IHC localization showing enhanced MT protein expression in cancer cells, basement membrane and lumen secretory products in biopsy specimens from PCa patients when compared to BPH specimens. Our preliminary data also indicated that MT-IIA overexpression alone has enhanced cell multiplication, collagenase (MMP-2) and MSA in a panel of PCa cell lines, LNCaP, PC-3, and DU-

145. These preliminary data have led us to hypothesize that MT plays a pivotal role in the promotion of neoplastic cell growth and in the progression of PCa to metastatic disease. Therefore, in this proposal we seek to ascertain whether the down-regulation of MT gene by antisense adenoviral construct will be able to inhibit growth and metastasis of prostate cancer cells.

(II) Objectives: To further test this hypothesis we proposed the following specific aims: (a) to construct an MT antisense adenoviral vector for achieving high transfection efficiency in human prostate cancer cells, LNCaP, DU-145 and PC-3; (b) to determine *in vitro* growth inhibitory effects of the MT antisense adenoviral construct; (c) to determine the MT antisense induced apoptosis by quantitative DNA fragmentation and TUNNEL methodology; (d) to monitor tumor growth delay and modulation of metastasis in athymic nude mice after transplantation of transfected cells.

(6) Body:

a. Optimization of transfection efficiency of prostate cancer cells with adenoviral vectors.

Adv5- β -galactosidase reporter construct was generously provided by Dr. David T. Curiel, University of Alabama at Birmingham, AL. Subconfluent cultures of prostate cancer cell lines, DU-145, PC-3 and LNCaP, were exposed to various concentrations of the reporter construct (10-100 MOI) and transfection efficiency was determined 24 hr post-transduction by measuring β -galactosidase activity *in situ*. Briefly, the cells were washed in phosphate buffered saline (PBS) and fixed (formaldehyde/formaldehyde solution) at room temperature for 5 min. Cells were then stained (potassium ferricyanide/ferrocyanide, $MgCl_2$ and X-gal) 2 hr to overnight at 37°C, washed in PBS and observed on an inverted microscope. Efficiency of transfection was measured by scoring dark blue cells positive. The results demonstrated that 100 MOI is optimum for achieving 95-100% transfection efficiency in all cell lines. The results are depicted in Figure 1.



Figure 1: Subconfluent cultures of PC-3 cells were transfected with Adv5- β -gal vector (100 MOI) for 24 hr. Cells were then fixed and stained for β -galactosidase activity *in situ* as shown above. The results demonstrated that adenoviral constructs can efficiently express a transgene (95-100%) in prostate epithelial cells.

b. Construction of an MT Antisense Adenoviral Vector.

Plasmids for adenovirus vector construction for transgene insertion in the E1 region of Ad5 vectors was purchased from Mirobix Biosystems Inc, Ontario, Canada. The kit is composed of the shuttle vector, pMH5, the circular replication-defective Ad genome plasmid pJM17A and the low-passage packaging cell line, 293 cells (Figure 2 below).





Figure 2: Adenovirus construction system: shuttle plasmid pMH5 (a) and replication-defective circular viral vector pJM17 (b); intracellular recombination (c)

5'EcoRI /3'XhoI' fragment encompassing the cDNA of mouse MT-I was initially digested with the restriction enzymes and subsequently removed from agarose gel by electroelution using Bio-Rad electroeluter (Figure 3). To subclone the digested fragment in reverse (or antisense) or the sense orientation (control) in the E1 region of shuttle plasmid, 5 micrograms of the pMH5 plasmid (purchased from Microbix Biosystems, Inc, Ontario, Canada, see map below) was initially digested with EcoRI and subsequently dephosphorylated using calf intestinal phosphatase (CIP) enzyme to prevent re-ligation. Ligation was carried out in accord with manufacture's instructions (Promega) using 1X buffer containing dATP and 3 U of T4 DNA ligase in 50 μ l final volume at 16°C overnight. The reaction was stopped by heating at 75°C for 10 min. The overhangs were then filled using Klenow fragment (1 U) (Promega) in presence of dNTPs at 30°C for 10 minutes and reaction was stopped by heating at 75°C for 10 min. Blunt end ligation was carried out as shown above in presence of 6 U of T4 DNA ligase overnight. Reaction was stopped and samples were stored at -20°C until used.

Figure 3:



Lane 1, λ -Hind-III marker; lanes 1-4, 5'EcoRI/XhoI3' digest; Lanes 5-8, mMT-I

b. Transformation.

Competent *E coli* DH₅ α was transformed with plasmids in accord with manufacturer's instructions. Briefly, bacterial cells were preincubated in ice with recombinant or control plasmids for 30 minutes prior to temperature shock at 42°C for 1.5 min. Bacterial cells were then grown in 1 ml LB with (shaking at 200 cpm) in antibiotic-free medium for 1 hr at 37°C. Cultures were then transferred into flasks containing 100 ml LB supplemented with ampicillin (50 μ g/ml) and subsequently grown for an additional 24 hr. Bacterial cell growth was detected spectrophotometrically. Samples of culture media were further cultured in LB agar plates containing ampicillin at 50 μ g/ml. Individual colonies (20) picked and further grown in LB agar plates.

c. Isolation of plasmid DNA .

Mini preps were used for individual clones by culturing bacteria in 10 ml LB media containing ampicillin overnight. DNA was isolated using standard methods. Briefly, Harvested overnight cultures were pelleted, washed in TE buffer, resuspended in GTE buffer containing lysozyme and incubated at 30°C for 20 min. Cell were lysed in presence of SDS and NaOH and plasmid DNA was isolated by centrifugation in presence of 3M K⁺ 5M OAc, pH 5.6. Ribonucleic acids were then removed using RNase A at 37°C for 30 min. DNA was extracted twice using phenol/chloroform/isoamyl alcohol (24:24:1) and precipitated with 1 volume of isopropanol. The pellets were washed once in 70% ethanol and stored at -20°C for further analysis.

d. Analysis of recombinant plasmids by restriction mapping.

To determine whether our DNA inserts (sense and antisense) were successfully cloned in the shuttle plasmid pMH5, we analyzed the DNA plasmids using 0.7% agarose gel containing ethidium bromide.

e. Generation of recombinant shuttle plasmids. Our results showed that among the twenty clones picked up for analysis of DNA inserts, none had shown a band or a band shift in the gel, as opposed to controls. We then decided to repeat the subcloning, transformation and analysis steps shown above without any successful attempt to subclone the mMT-I fragment. Our results have indicated that the bacteria were successfully cloned with the control plasmids, but not with our recombinant shuttle plasmids.

Even though we scrutinized all of our steps to subclone the DNA inserts we were unable to detect why this system have failed (see results below). We concluded that, the shuttle plasmid we purchased may have defective multiple cloning site at the EcoR1 site.




■ ■ ■ ■ ■ ◀ pmMT-I

f. Use of pAd_{vantage} System for construction of MT antisense vector.

Our next approach was to use an alternative system for cloning hMT-II inserts in both orientation. A novel cloning method for recombinant adenovirus construction in *Escherichia coli* has been developed by Genzyme Co, Framingham, MA) (28). The pAd_{vantage} system was a generous gift by Dr. D. W. Souza Genzyme Corporation. The system is composed of the shuttle plasmid pSV-ICEU I (containing the cytomegalovirus promoter (CMV), multiple cloning site (MCS) and polyadenylation site)(Figure 3a) and the viral vector pAd_{vantage} (Figure 3b). Because of close homology between human metallothionein and the use of human prostate cancer cells, we decided to use human MT-IIa cDNA (pBacNEO-sMT-IIa) to generate the antisense adenovirus constructs (Figure 3c).

Figure 3: pAd_{vantage} system: (a) the shuttle plasmid pSV-ICEU I; (b) pAd_{vantage} viral vector; and (c) pBacNEOsMT-IIa.



g. Isolation of hMT-IIa from pBacNEO-sMT-IIa plasmid.

Twenty microgram of pBacNEO-sMT-IIa plasmid was initially digested with 3'XhoI-KpnI5' for 2 hr at 37°C. The products were then fractionated into a 0.7% agarose gel containing ethidium bromide. A DNA fragment (approx 300 bp) encompassing hMT-IIa was then isolated using electroelution technique in accord with the manufacturer instructions (Figure 4). Briefly, gel slices containing the DNA were placed in an apparatus equipped with tubes capped with special membranes and DNA was collected in the membranes by electroelution and subsequently precipitated using standard techniques. The cDNA was then stored at -20°C until used.

and extension at 59°C for 15 sec. The products were then fractionated into a 0.7% agarose gel containing ethidium bromide. A DNA fragment (approx 300 bp) encompassing hMT-IIa was then isolated using electroelution technique in accord with the manufacturer instructions (Figure 4). Briefly, gel slices containing the DNA were placed in an apparatus equipped with tubes capped with special membranes and DNA was collected in the membranes by electroelution and subsequently precipitated using standard techniques. The cDNA was then stored at -20°C until used.




Figure 4: (a) Agarose gel electrophoresis (0.7%) showing 3'XhoI- KpnI5' hMT-IIa fragment (approximately 320 bp); lane 1 3'XhoI- KpnI5 double digest, lane 2 DNA marker ; (b) PCR amplification of hMT-IIa using pBacNEOs-MT-IIa as a template, lane 1, hMT-IIa; lane 2, pBacNEOs-MT-IIa plasmid

h. Cloning of hMT-IIa in the shuttle plasmid pSV-ICEU I.

Preparation of the shuttle plasmid:

The shuttle plasmid pSV-ICEU I was initially digested with *Xho*-I, dephosphorylated with calf intestine phosphatase (CIP) to prevent re-ligation, and precipitated with phenol/chloroform in accord with standard techniques and the stored -20°C until used.

Ligation:

For preparation of the antisense construct, the 3'XhoI- KpnI5' fragment encompassing the human MT-IIa was initially ligated to the shuttle plasmid pSV-ICEU I using 1X ligation buffer, 3 U of T4 DNA ligase, 0.25 µg of shuttle plasmid with three-fold molar excess of the DNA insert at 15°C overnight. The reaction was stopped at 75°C for 10 min. The overhangs were then filled with Klenow fragment (1 U) (Promega) in presence of dNTPs at 30°C for 10 minutes and reaction was stopped by heating at 75°C for 10 min. Blunt end ligation was carried out as shown above in presence of 400 U of T4 DNA ligase (New England Biotechnology) overnight. The reaction was stopped and samples were stored at -20°C. For the preparation of the control sense plasmid, both *Xho*I-digested pSV-ICEU I and 3'XhoI- KpnI5' fragment were initially filled using Klenow fragment as shown above to generate blunt ends and ligation was carried out using high concentration of T4 DNA ligase (New England BioLabs) at 15°C overnight.

Transformation and plasmid Isolation:

Competent *E coli* DH_{5α} was transformed with plasmids in accord with recommended procedure as described above. Mini preps were used for isolation of individual clones in accord with standard protocol as shown above.

Analysis of ligation:

The fidelity of DNA insertion was analyzed using 0.7% agarose gel containing ethidium bromide. The results demonstrated that hMT-IIa was efficiently subcloned in the shuttle vector (Figure 5). Initially, all colonies screened for the control vector have shown no DNA insert and we had repeated the whole cloning techniques before we were successful on isolating a positive clone.




Figure 5: Gel electrophoresis demonstrating a size shift in the recombinant shuttle vector as opposed to control plasmid. M, DNA marker; lane 1-5 MT -IIa antisense recombinant clones; lanes 7-9 control MT-IIa (sense) recombinant clones; lanes 10-12, shuttle plasmid

Analysis of orientation:

The orientation of the DNA insert was performed using restriction mapping of the recombinant plasmids followed by analysis using 0.7% agarose gel electrophoresis. The results demonstrated that the transgene is in the antisense orientation.




Figure 6: Digestion of the recombinant plasmids with *ScaI* and *BamHI*. B. It is anticipated that three (lanes 1-3) fragments will be generated following the double digestion in the positive clones. Lane 3 represents a positive clone for the MT antisense clone. M, DNA marker.

Digestion of recombinant plasmid with I-Ceu I:

A fragment encompassing CMV promoter-antisense MT-IIa cDNA and poly A regions was digested from the recombinant shuttle plasmid pSV-ICEU I using I-Ceu I (New England Biolabs). The DNA insert was then isolated using agarose gel electrophoresis and electroelution as described above. After precipitation, the pellet was stored at -20°C.

J. Subcloning of the antisense MT-IIa from pSV-ICEU I into the I-Ceu I site of the pAd_{vantage} viral vector.

Preparation of pAd_{vantage} for cloning:

Five micrograms of pAd_{vantage} were digested with 30 U of I-Ceu I in a total volume of 500 µl of 1 X I-Ceu I buffer at 37°C for 3 hr. 10 units were added every hr to reach 30 U by the end of the 3 hr period. The plasmid was then dephosphorylated using CIP as shown above for an additional 20 min at room temperature and DNA was extracted using phenol/chloroform and precipitated with ethanol as shown above.

Subcloning of the I-Ceu I fragment into the I-Ceu I-site of the pAd_{vantage}:

The viral vector (0.25 µg) and three-fold molar excess of the I-Ceu I DNA insert were incubated at 16°C overnight in presence of 400 U of T4 DNA ligase and 1 X ligase buffer in a total volume of 50 µl. The fidelity of the DNA insert in the viral vector was verified by transformation, screening of putative colonies, and by agarose gel electrophoresis when compared to the viral vector alone as we showed above.




Figure 7: Agarose gel electrophoresis (0.7%, TAE Buffer) for analysis of MT antisense adenovirus construct. A representative clone (lanes 1) was analyzed for efficient ligation of I-Ceu I fragment into the I-Ceu I-site of the pAd_{vantage} and compared to the control plasmid (lane 2). A shift in band size in lane 1 as opposed to lane 2 indicates successful ligation.

k. 293 Cell Culture.

Low passage 293 cells, a human embryo kidney cells transformed by sheared Adenovirus type 5 DNA, were obtained from Microbix Biosystems Inc., (Ontario, Canada). They contain and express the early region (E1) of Ad 5 virus and can complement the E1 defective Adv mutants and vectors. The cells were grown in MEM supplemented with Earle's salts, glutamine, 10% heat-inactivated FBS, penicillin and streptomycin and incubated at 37°C with 5% CO₂.

I. Viral Packaging in 293 cells.

Preparation of the viral constructs (by first digesting the construct with *Sna*BI to expose the inverted terminal repeats) for packaging in 293 cells is now underway and it is anticipated the viral plaques will become visible in about 7-10 days.

(7) Key Research Accomplishments:

- Generation of novel recombinant antisense MT adenovirus vector.

(8) Reportable outcomes:

- Construction of metallothionein MT adenovirus construct.
- Applied for funding for a DOD grant in 1999 entitled "Antisense down-regulation of metallothionein: a combined therapeutic approach for prostate cancer."
- The data will be also used for funding application to the NIH, ACS and CaPCure.

(9) Conclusions.

The aim of this training grant was to construct a metallothionein antisense adenovirus vector and test its efficacy on enhancing growth arrest and inducing apoptosis in cancer cells using *in vitro* and *in vivo* approaches. - Like other drug resistance genes, metallothionein has been shown to play a role in protecting cancer cells against chemotherapeutic agents and radiotherapy— thus this vector can also be used to enhance sensitization of cancer cells to chemo/radiotherapy. Because this application was a *training* grant (no funds for research supplies or other personnel), and due to the unexpected research problems, I wasn't able to accomplish all specific aims within the short-time frame specified for the grant (6 months). One recommendation, at least, is to include funding for research supplies and to extend the time-frame for these training grants to 1-2 years, so that the research will be completed.

(10) References.

1. Parkin, D.M., Pisani, P., and Ferlay, J. Estimates of the worldwide incidence of 18 major cancers in 1985. *Int. J. Cancer*, 54:594-606, 1993.
2. Boyle, P., Maisonneuve, P., Napalkov, P. Incidence of prostate cancer will double by year 2030: the argument for. *Eur Urol.*, 29 (Suppl 2):3-9, 1996.
3. Jensen, O.M., Esteve, J., Moller, H., and Renard, H. Cancer in the European community and its member states. *Eur. J. Cancer*, 25:1167-1256, 1990.
4. Boring, C.C., Squires, T.S., and Tong, T. cancer Statistics, CA. *Cancer J. Clin.*, 44:7-26, 1994.
5. Parker, S.L., Tong, T., Bolden, S., and Wingo, P.A. Cancer Statistics 1996., CA. *Cancer J. Clin.*, 1996.
6. Davis KJ, Martin L, Parker L, et al. Cancer Facts & Figures for African Americans. American Cancer Society, Inc., 1996.
7. Kosary, CL, Ries LAG, Miller BA et al: SEER Cancer Statistics Review, 1973-192. National cancer Institute. NIH Publication No. 95-2789. Bethesda, MD, 1995.
8. Plepys C, Klein R Health Status Indicators: Differentials by Race and Hispanic Origins. Statistical Notes; No. 10, Hyattsville, MD: National Center for Health Statistics. 1995.
9. Kägi J.H.R. and Schäffer. Biochemistry of metallothionein. *Biochemistry*. 27:8509-15, 1988.
10. Schmidt CJ. Hamer DH. McBride OW. Chromosomal location of human metallothionein genes: implications for Menkes' disease. *Science*. 224(4653):1104-6, 1984.
11. Karin M., Eddy R.L., Nienry M. et al. Human metallothionein genes are clustered on chromosome 16. *Proc. Natl. Acad. Sci., USA*, 81:5494-98, 1984.
12. West, A.K., Stallings, R., Hildebrand, C.E., Chiu, R., Karin, M. and Richards, R. Human metallothionein genes: structure of the functional locus at 16q 13. *Genomics* 8, 513-518, 1990.
13. Stennard, F.A., Holloway, A.F., Hamilton, J. and West, A.K. (1994) Characterization of six additional human metallothionein genes. *Biochim. Biophys. Acta* 1218, 357-365.
14. Karin, M. and Richards, R.I. Human metallothionein genes-primary structure of the metallothionein-II gene and a related processed gene. *Nature* 299, 797-802, 1982.
15. Palmiter, R.D., Findley, S.D., Whitmore, T.E. and Durnam, D.M. MT-III, a brain-specific member of the metallothionein gene family. *Proc. Natl. Acad. Sci.* 89, 6333-6337, 1992.
16. Quaife, C.J., Findley, S.D., Erickson, J.C., Froelick, G.J., Kelly, E.J., Zambrowicz, B.P. and Palmiter, R.D. Induction of a new metallothionein isoform (MT-IV) occurs during differentiation of stratified squamous epithelia. *Biochemistry* 33, 7250-7259, 1994.
17. Richards, R.I., Heguy, A. and Karin, M. Structural and functional analysis of the human metallothionein-I_A gene: differential induction by metal ions and glucocorticoids. *Cell* 37, 263-272, 1984.
18. Heguy, A., West, A., Richards, R.I. and Karin, M. Structure and tissue-specific expression of the human metallothionein I_B gene. *Mol. Cell. Biol.* 6, 2149-2157, 1986.
19. Sadhu, C. and Gedamu, L. Regulation of human metallothionein (MT) genes. Differential expression of MTI-F, MTI-G, and MTII-A genes in the hepatoblastoma cell line (HEPG2). *J. Biol. Chem.* 263, 2679-2684, 1988 .

20. Jahroudi, N., Foster, R., Price-Haughey, J., Beitel, G. and Gedamu, L. Cell-type specific and differential regulation of the human metallothionein genes. *J. Biol. Chem.* 265, 6506-6511, 1990.
21. Zelger, B. Hittmair, A., Schir, M. et al. Immunohistochemically demonstrated metallothionein expression in malignant melanoma. *Histopathol.*, 23:257-64, 1993.
22. Murphy D., McGown A.T., Crowther D., Mander A., and Fox B.W. Metallothionein levels in ovarian tumors before and after chemotherapy. *Br. J. Cancer*, 63:711-14, 1991.
23. Chin J.L., Banerjee D., Kadhim S.A., et al. Metallothionein in testicular germ cell tumors and drug resistance. *Clinical Correlation Cancer*, 72:3029-35, 1993.
24. Bahnson R.R., Banner B.F., Ernstoff et al. Immunohistochemical localization of metallothionein in transitional cell carcinoma of the bladder. *J. Urology*, 146:1518-20, 1991.
25. Saika T., Tsushima T., Nasu Y., et al. Histopathological study of metallothionein in bladder cancer and renal cell carcinoma. *Nippon Hinyokika Gakkai Zasshi-Japanese J. of Urology*, 83:636-42, 1992.
26. Abdel-Mageed A.B., and Agrawal K.C. Antisense down-regulation of metallothionein induces growth arrest and apoptosis in breast carcinoma cells. *Cancer Gene Therapy*, 4:199-207, 1997.
27. Abdel-Mageed, A.B., and Agrawal, K.C. Activation of NF-kappa B: A potential mechanism for metallothionein-induced mitogenic response. *Cancer Res*, 58:2335-2338, 1998.
28. Souza, D.W., and Armentano, D. Novel Cloning method for recombinant adenovirus construction in *Escherichia coli*. *Biotechniques*. 26:502-508, 1999.

11. Appendices: N/A



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

23 Aug 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to the technical reports listed at enclosure. Request the limited distribution statement for these reports be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Reports to be Downgraded to Unlimited Distribution

ADB241560	ADB253628	ADB249654	ADB263448
ADB251657	ADB257757	ADB264967	ADB245021
ADB263525	ADB264736	ADB247697	ADB264544
ADB222448	ADB255427	ADB263453	ADB254454
ADB234468	ADB264757	ADB243646	
ADB249596	ADB232924	ADB263428	
ADB263270	ADB232927	ADB240500	
ADB231841	ADB245382	ADB253090	
ADB239007	ADB258158	ADB265236	
ADB263737	ADB264506	ADB264610	
ADB239263	ADB243027	ADB251613	
ADB251995	ADB233334	ADB237451	
ADB233106	ADB242926	ADB249671	
ADB262619	ADB262637	ADB262475	
ADB233111	ADB251649	ADB264579	
ADB240497	ADB264549	ADB244768	
ADB257618	ADB248354	ADB258553	
ADB240496	ADB258768	ADB244278	
ADB233747	ADB247842	ADB257305	
ADB240160	ADB264611	ADB245442	
ADB258646	ADB244931	ADB256780	
ADB264626	ADB263444	ADB264797	