

UNCLASSIFIED

AD NUMBER
ADB268358
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Mar 2001. Other requests shall be referred to US Army Medical Research and Materiel Comd., 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 21 Feb 2003

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-98-1-8487

TITLE: Exploiting Novel Polyamine Regulatory Responses to a
Therapeutic Advantage in Human Prostatic Carcinoma: A
Preclinical Study

PRINCIPAL INVESTIGATOR: Carl W. Porter, Ph.D.

CONTRACTING ORGANIZATION: Health Research, Incorporated
Buffalo, New York 14263

REPORT DATE: March 2001

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government
agencies only (proprietary information, Mar 01). Other requests
for this document shall be referred to U.S. Army Medical Research
and Materiel Command, 504 Scott Street, Fort Detrick, Maryland
21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

20010723 066

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-98-1-8487

Organization: Health Research, Incorporated

Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Andrew S. Clemens
0 1/05/2001

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 074-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE March 2001	3. REPORT TYPE AND DATES COVERED Annual (1 Sep 99 - 28 Feb 01)		
4. TITLE AND SUBTITLE Exploiting Novel Polyamine Regulatory Responses to a Therapeutic Advantage in Human Prostatic Carcinoma: A Preclinical Study		5. FUNDING NUMBERS DAMD17-98-1-8487		
6. AUTHOR(S) Carl W. Porter, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Health Research, Incorporated Buffalo, New York 14263 E-Mail: porter@sc3101.med.buffalo.edu		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Mar 01). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) We have previously reported that two polyamine antagonists currently undergoing clinical evaluation: the polyamine analog, N ¹ , N ¹¹ -diethylnorspermine (DENSPM) and the biosynthetic enzyme inhibitor, 4-amidinoindan-1-one 2'-amidinohydrazone (CGP-48664), appear to exert unusual regulatory effects in prostate carcinoma cell lines relative to other cell lines. We have proposed to develop strategies <i>in vitro</i> and <i>in vivo</i> for therapeutically exploiting this observation. In pursuit of this goal, we have discovered that CGP-48664 potently induces apoptosis via a p53 / caspase-3 dependent pathway that is unrelated to its intended mode of action as an inhibitor of polyamine biosynthesis. Such an effect has not been previously reported and may have relevance to ongoing clinical trials with the compound. Future studies will investigate the therapeutic significance of this finding <i>in vivo</i> . In other developments, we have determined that a polyamine analog related to DENSPM exerts meaningful antitumor activity against DU145 human prostate carcinoma xenografts and are seeking proof-of-principle for a novel polyamine gene-based antiproliferative strategy. The former will be optimized with respect to drug effects on polyamine regulatory responses and on the antitumor activity of the compound. The overall goal of both research efforts is to identify and develop polyamine-based / prostate-directed therapies as rapidly as possible.				
14. SUBJECT TERMS Apoptosis, analogs, antitumor, CGP-48664, DENSPM, inhibitors, Polyamines, prostate			15. NUMBER OF PAGES 28	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

✓ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Carl W. Porter 2/14/01
PI - Signature Date

TABLE OF CONTENTS

<u>PAGE</u>	<u>DESCRIPTION</u>
	Front Cover
2	Report Documentation
3	Foreword
4	Table of Contents
5	A. Introduction
5	B. Tasks
6	C. Progress
10	D. Key Research Accomplishments
10	E. Reportable Outcomes
11	F. Conclusions
11	G. References
12	Appendices (Figure Legends and Figures 1-14)

REPORT BODY (Figures & Figure Legends located in Appendix 1)**A. Introduction:**

Polyamines are organic cations found in all cells and known to be essential for the initiation and maintenance of cell growth. Intracellular levels are sensitively maintained by a series of regulatory responses. The relationship of polyamines to the prostate is unique among all tissues since in addition to synthesizing these molecules for cell growth, the gland produces massive quantities for export into semen. It might, therefore, be expected that prostatic tumors could exhibit atypical polyamine-related regulatory responses. We have recently observed that in contrast to other normal and tumor cell types, two of three prostate carcinoma lines displayed an altered ability to regulate polyamine transport in responses to polyamine analog or inhibitors (1). More specifically, two lines could not down-regulate uptake in response to analogs and one could not up-regulate it in response to polyamine depletion caused by inhibitors. The findings have clear therapeutic implications for two polyamine antagonists that have recently undergone clinical evaluation: the polyamine analog, N¹, N¹¹-diethylnorspermine (2, DENSPM; Parke-Davis) and the biosynthetic enzyme inhibitor, 4-amidinoindan-1-one 2'-amidinohydrazone (3-4, CGP-48664, also known as SAM486A Novartis). Thus, it should be possible to identify more effective analogs of these drugs and/or to develop drug treatment schedules capable of optimizing selective delivery of DENSPM (which utilizes the transporter) to prostate tumors and in the case of CGP-48664, to minimize antiproliferative escape via salvage of exogenous polyamines. Herein, we propose to further examine polyamine regulatory responses in prostate carcinoma cell lines treated with analogs and inhibitors; to examine the underlying molecular mechanism(s) responsible for alterations in those responses; and most importantly, to develop *in vitro* and *in vivo* strategies for therapeutically exploiting these fundamental observations. We believe that we are particularly well-poised in terms of long term involvement with both polyamine analogs and inhibitors. Given the current clinical status of both drugs, promising findings should be readily translatable into experimental therapies for prostate carcinoma patients.

B. Originally Proposed Tasks

- Task 1. To further determine the extent to which altered polyamine homeostatic regulatory responses are characteristic of prostate cancer cell lines.
- Task 2. To develop effective *in vitro* strategies based on observations made in Task 1 which optimize the antiproliferative effects of DENSPM, CGP-48664 or related inhibitors and analogs.
- Task 3. To investigate mechanisms underlying aberrant regulatory responses identified in Task 1.
- Task 4. To extend strategies developed in Task 2 to *in vivo* model systems involving appropriate human prostatic carcinoma lines growing in scid mice.
- Task 5. To further optimize the choice of analog and/or inhibitor using compounds available to us from various collaborating laboratories.

- **Task Modifications:**

The ordered pursuit of the above proposed tasks has been altered by three significant developments: (i) our observation that CGP-48664 exerts a significant apoptotic effect in LNCaP prostate carcinoma cells and that this response seems to be unrelated to its intended activity as an inhibitor of polyamine biosynthesis; (ii) clinical trials indicated that schedule modification of DENSPM (i.e. splitting a single daily dose to twice or three times daily, as we proposed in Tasks 1 and 2) is not well tolerated in patients and (iii) the current halt in clinical trials of CGP-48664 and DENSPM. In both cases, the decision was based on priority relative to alternative drugs available in the pipeline of Novartis and Pfizer following the take over of Parke Davis. At present, Novartis is seeking to license SAM-486 to a smaller company with expertise in polyamines such as Oridigm (Seattle, OR) or Ilex (San Antonio, TX). Similarly, Parke Davis has returned the DENSPM license to Geltex Pharmaceuticals (Waltham, MA) who is now raising funding for clinical trials. While these issues are being resolved, we have placed greater emphasis on Task 5 as it relates to identification of new polyamine analogs having greater antitumor activity and/or reduced host toxicity (in collaboration with S'LIL Pharmaceuticals, Madison, WI). In addition, we have begun the investigation of a novel gene-based strategy that could have important implications in prostate cancer.

C. Progress:

1. Mode of CGP-48664 Drug Action in LNCaP Cells:

As originally proposed in Task 1b, we began this project by attempting to confirm variant polyamine regulatory responses to CGP-48664 and DENSPM by examining the effects of other known polyamine inhibitors, and analogs. The goal was to determine that the responses could be generalized to compounds having comparable modes of action as CGP-48664 and DENSPM before moving on to examine the generality of altered polyamine responses in additional prostate carcinoma cell lines. An initial comparison of growth effects between CGP-48664 and MDL-46811, a well known mechanism-based specific inhibitor of S-adenosylmethionine decarboxylase (2, SAMDC), revealed that at 50 μ M, the former inhibitor induced an immediate and complete cessation of LNCaP cell growth while at 100 μ M, the latter had virtually no effect on the growth these cells (Figure 1). As expected for SAMDC inhibitors (2-4), both drugs similarly increased putrescine and decreased spermidine and spermine pools (data not shown). Our immediate suspicion was that CGP-48664 was exerting unexpected drug effects that were unrelated to inhibition of SAMDC. This possibility was further indicated by the observation that growth inhibition was not preventable with exogenous spermidine.

In the first year report, we indicated that CGP-48664 induces apoptosis in LNCaP cells (subG1 peak and DNA fragmentation) and this cellular response appears to be wholly unrelated to drug effects on SAMDC. Additional findings indicated that the apoptotic effect of CGP-48664 may be related to induction of p53 and more clearly, to activation of caspase-3 dependent apoptosis. The possible protein response interrelationships as they relate to caspase 3-induced apoptosis were defined. A comparison of cytochrome c levels in mitochondrial versus the S-100 fractions of LNCaP treated cells revealed that CGP-48664 caused the mitochondrial protein to be released into the cytosolic fraction; a typical feature of p53/caspase-mediated apoptosis. Indeed, early activation of caspase-3, and caspase-mediated cleavage of poly(ADP-ribose) polymerase (PARP) were seen in treated cells. The individual who performed the original studies has since

accepted a postdoctoral position at San Antonio. We subsequently discovered that the LNCaP cells used in his experiments were contaminated with mycoplasma. In attempting to accommodate the recommendations of reviewers using uncontaminated cells, we were unable to repeat the caspase-3 induction, PARP cleavage or DNA fragmentation reported in the manuscript. We were however, able to demonstrate that CGP-48664 does cause apoptosis by Annexin V staining and that it was not preventable with spermidine. Induction of p53 and p21 are also repeatable. Thus, CGP-48664 induces apoptosis in LNCaP cells via a polyamine unrelated pathway that remains to be defined. This development constitutes a major set-back in progress. Future studies will attempt to identify the pathway responsible for the apoptotic response to CGP-48664.

2. Antitumor Studies with Recently Available Polyamine Analogs:

Last year, we reported CGP-48664 at 50 mg/kg per day i.p. (commonly used dose for this drug) showed absolutely no antitumor activity (data not shown) while DENSPM at 80 mg/kg per day i.p. at the maximally tolerated dose for 5 days yielded a modest antitumor activity which was apparent as a slowing in tumor growth followed by a full growth rate recover (data not shown). During the past two years, we collaborated with S'LIL Pharmaceuticals (5, Madison, WI) in accessing a novel polyamine analogs that were found by company, to have *in vitro* activity against prostate carcinoma cells lines and *in vivo* activity in a benign prostate hyperplasia model (*PROPRIETARY INFORMATION from S'LIL*). In previous reports, we indicated that SL-11147, an unsaturated analog of DENSPM having a double bond linking the two center carbons of the molecule, produced antitumor activity nearly identical to DENSPM in the DU145 model. We now report on the antitumor activity of two additional compounds SL-11144 and SL-10093 (Figure 1). The first, SL-11144 (Figure 1), is a diethylated polymer of homospermine units. More specifically, SL-11144 is a terminally N-diethylated molecule with an intra-amine carbon arrangement of [4x(4-4-4)]-2-db-2-[4x(4-4-4)] where db indicates an unsaturated double bond in the middle of a central homospermine moiety. The second, SL-11093 (Figure 1), is an analog of N^1 , N^{11} -diethylhomospermine also known as DEHSPM where DE refers to diethyl and where *homo* refers to the intra-amine carbon spacings as 4-4-4 as opposed to 3-4-3 with spermine. The analog is rendered particularly novel by a unique cyclopropyl group connecting the two center carbons of the molecule. This SL-11093 was selected on the basis of its close structural similarity to DEHSPM which has undergone a partial Phase I clinical trial and on its *in vitro* antiproliferative activity against the three human prostate carcinoma cell lines. Against human prostate carcinoma lines, ID₅₀ values were in the range of 10-15 μ M.

In vivo studies were preceded by *in vitro* growth inhibition studies involving DU-145 cells, SL-11144 was much more cytotoxic than SL-11093 with respect to both dose (Figure 2) and time (Figure 3). Whereas SL-11144 affected growth kinetics in a manner entirely consistent with high cytotoxicity, SL-11093 was weakly antiproliferative and cytostatic at best. For example, the dose-response curves shown in Figure 2 show that growth falls off rapidly with increasing doses of SL-11144, this does not occur with SL-11144 where following an initial fall-off in growth, cells seem to be unaffected by increasing concentrations of analog (i.e. 1 to 100 μ M). As further evidence for the cytotoxicity caused by SL-11144, we note that by annexin staining, 48% of DU-145 cells treated for 48 h with the analog were in late apoptosis while by comparison, only 3.6% of cells treated with SL-11093 showed signs of apoptotic staining.

Antitumor studies began with the treatment of established DU-145 prostate carcinoma tumors (~ 200 mm³) at analog doses approaching the maximally tolerated dose as determined by Dr. Cyrus Bacchi (Pace University, NYC). More specifically, SL11144 was administered

intraperitoneally at 5-10 mg/kg/d for 5 days per dosing round and SL-11093 was administered at 25-50 mg/kg/d for 5 days per dosing round. Control animals were injected intraperitoneally with saline on the identical schedule. Owing to simplicity of structure, analog availability and lower potential for host toxicity, initial studies were carried out with SL-11093.

A pilot study was conducted with a single dosing round of SL-11093 at 25 and 50 mg/kg/d for 5 days. As indicated in Figure 4, tumor growth was modestly affected in a dose-dependent fashion. Because there was minimal host toxicity and because the analog seemed to be most effective during treatment, we evaluated SL-11093 in a follow-up experiment where the dosing schedule was repeated three times with a 9 day rest period in between dosing rounds. As shown in Figure 5, the growth of DU-145 tumors in treated mice was markedly suppressed (Figure 5B) relative to the saline treated mice (Figure 5A). The drug effect seemed tumoristatic since tumor growth remained relatively flat and there was no indication of tumor regressions. For uncertain reasons, one tumor seemed to escape the antitumor effects of the drug (Figure 5B). When the median values for both groupings are compared in Figure 5C, it is apparent that tumor growth suppression by SL-11093 was highest during treatment and that tumors rapidly recovered growth shortly after treatment ceased. As indicated in Figure 5D, there was a minor loss in body mass during treatment after which mice attained weights similar to those of saline-treated mice.

We then examined the antitumor activity of SL-11144 (at 5 mg/kg/d) under the schedule of multiple dosing rounds used with SL-11093. As shown in Figure 6, the analog initially suppressed tumor growth during the first two rounds of treatment but then lost effectiveness during the third round. Animal weights indicated minimal host toxicity during the first three rounds of treatment after which there appeared to be a latent weight loss effect.

The multiple (5) dosing experiment with SL-11093 (at 50 mg/kg/d) was then repeated and compared with SL-11144 (at 5 mg/kg/d). Antitumor effects during and shortly following treatment with either analog (days 20-98) are shown in Figure 7. Once again, the growth of DU-145 tumors in SL-11093-treated mice was effectively suppressed (Figure 7B) relative to the saline-treated mice (Figure 7A). By comparison, SL-11144 seemed less active in suppressing tumor growth since several tumors escaped its effects (Figure 7C) and those tumors that responded were less affected than those treated with SL-11093. Although there was virtually no decrease in body weight during treatment with either analog (Figure 7D), saline mice gained weight during the course of the experiment while those treated with analog did not. As with the initial study (Figure 4), the second experiment strongly suggested that the drug effect was tumoristatic. This is most apparent in Figure 8 which shows tumor volumes (within the same experiment) beyond the treatment period shown in Figure 7. Except for two anecdotal regressions in saline (Figure 8A) and SL-11144 (Figure 8C) treated mice, there were no tumor regressions that could be reliably attributed to drug treatment. The strongest indication for the tumoristatic nature of SL-11093 derives from the finding that after being suppressed for ~70 days, tumor growth sharply resumed about 20 days following the last round of treatment with the analog. Taken together, this experiment confirms earlier indication that multiple rounds of dosings with SL-11093 produce meaningful antitumor responses with minimal host toxicity. The antitumor effect is tumoristatic and SL-11093 is more effective than SL-11144 against the DU-145 prostate carcinoma. As a result of these experiments, SL-11093 is being developed towards clinical trials for prostate cancer.

In the most recent experiment, we examined the antitumor activity of SL-11093 against Panc-1 pancreatic carcinoma xenografts. The rationale was based on the clinical need for effective therapies against this disease and the fact that the exocrine pancreas typically contains high levels of polyamines, perhaps because it is heavily involved in protein synthesis. As shown in Figure 9, the experiment was complicated by heterogeneous growth characteristics of the

untreated tumor (Figure 9A). Despite beginning at similar starting volumes, tumors tended to either grow rapidly or slowly. As shown in Figure 9B, this same pattern of tumor growth was also seen in the SL-11093 treated tumors. Due to this tumor scatter, the tumors were not subjected to multiple dosing rounds. However, when the saline and analog-treated groups were represented within the same plot (Figure 9C), it appears that a single round of therapy is virtually without effect on tumor growth. Although inconclusive, the data suggest that SL-11093 is much less active against the Panc-1 pancreatic carcinoma than DU-145 prostate carcinoma (Figure 4). Efforts to reduce the growth heterogeneity of the model via passaging are underway before a repeat the experiment using multiple rounds of analog therapy is attempted. If negative, the data would be indicate possible disease specificity of SL-11093 for prostate carcinoma.

Overall, we believe that the data with SL-11093 are especially encouraging. Both we and S'LIL Pharmaceuticals have interest in developing this particular analog this toward clinical trial. Ongoing studies are examining of oral availability of the analog. In addition, we propose to study the biochemical polyamine parameter profile of tumors (focusing on drug accumulation) treated with these novel compounds in order to gain insight into their mode of *in vivo* drug action and how it compares to DENSPM (6-8). Finally, we will attempt to optimize the biochemical and antitumor effects using analogs of SL-11144 found by investigators at S'LIL to have promising *in vitro* activity against prostate carcinoma cell lines.

3. Antiproliferative potential of a polyamine catabolic enzyme:

In the last report, we alluded to studies in which conditional expression transfection strategies (tetracycline repression) were used to unequivocally demonstrate that (a) overexpression of the polyamine catabolic enzyme, spermidine/spermine N¹-acetyltransferase (SSAT) leads to polyamine pool depletion and inhibition of cell growth (Figure 10) and (b) basal SSAT activity levels determine sensitivity to DENSPM (Figure 11), (c) overexpression of the enzyme greatly increases sensitivity to DENSPM (Figure 12) and (d) modest expression of SSAT prevents uptake of polyamines (data not shown) and hence, circumvention of growth inhibition by polyamine inhibitors. The findings are reported in reference 10. The implications of these findings are multifold. At a minimum, regulated expression of SSAT can influence sensitivity to certain analogs, inhibit cell growth and/or augment antitumor activity of polyamine inhibitors by preventing salvage of exogenous polyamines.

By way of convenience, these experiments were conducted in MCF-7 breast carcinoma cells (already containing the tetracycline-repressible trans-activator system). Given the demonstrated novelty of polyamine homeostasis in prostate carcinoma (1), we feel that it would be highly worthwhile to use the same technology (conditional regulation of SSAT) to make similar determinations in a prostate carcinoma cell line. Anticipated findings would be used as the basis for a possible gene therapy strategy to enhance the therapeutic potency and selectivity DENSPM towards prostate cancer (this effort is not covered by any other funding mechanism). Towards this goal, we have undertaken the following approaches: (a) to prepare adenovirus vectors engineered to express SSAT under a CMV and PSA promoter and (b) to conditionally express SSAT using a tetracycline expression system similar to that used in our earlier studies. Both studies are underway. With regard to the latter, a LNCaP subline containing the tetracycline-repressible trans-activator system has been obtained from Dr. Thomas Powell (Cleveland Clinic, 9). Human SSAT cDNA cloned into pTRE expression vector containing a tetracycline responsive element (TRE) has been transfected into the above mentioned LNCaP subline. Three high conditional SSAT overexpression clone, SSAT/LNGK9-17, were isolated and found to have express 28-, 34- and 35-fold more SSAT activity in the absence of Doxycycline

than in the presence of the antibiotic (Figure 13). Biological characterization of these cells with respect to growth, polyamine pool effects and sensitivity to the polyamine analog DENSPM is underway.

In anticipation of adenoviral gene therapy type studies, we have cloned human SSAT cDNA into a replication-deficient adenoviral AdCMV5-IRES-GFP vector with an internal ribosomal entry site (IRES) for green fluorescent protein (GFP), both under the control of a CMV5 promoter for maximum expression (Q-Biogene, Montreal, Canada). Recently, we selected an adenoviral clone which expresses the SSAT and GFP dicistronic message (Figure 14). The selected clone is now undergoing three rounds of plaque purification to obtain the highest SSAT expressing recombinant adenovirus.

D. Key Research Accomplishments:

- Characterization of the antitumor activity of new and novel polyamine analogs against human DU145 prostate carcinoma xenografts. In particular, SL-11093 displays antitumor activity on a multi-treatment schedule that would seem to warrant further preclinical drug development and this possibility is being pursued.
- Experimental evidence that the polyamine catabolic enzyme SSAT is a determinant of drug action for polyamine analogs and when it is overexpressed in MCF-7 human breast carcinoma cells, the enzyme depletes polyamine pools, inhibits cell growth and prevent uptake of exogenous polyamines.
- Consistent with the previous finding and our interest in prostate carcinoma, we have developed a LNCaP prostate carcinoma in which SSAT is conditionally expressed under a tetracycline promoter and for the purpose of gene therapy modeling, we are well-along in developing adenovirus particles capable of transducing SSAT.

E. Reportable Outcomes:

- Mi, Z., Vredenberg, M.R., Kramer, D.L., Diegelman, P. and Porter, C.W. Induction of caspase-3 dependent apoptosis by the S-adenosylmethionine decarboxylase inhibitor CGP-48664 in human prostate carcinoma cells. Cancer Res. (Provisional acceptance; see letter in Appendix)
- Siu, L.L., Rowinsky, E.K. Weiss, G.R., Hammond, L., Kraynak, M., Moczygemba, J., DeMoor, C., Choi, L., Linnartz, R., Barbet, N.C., Capdeville, R., Von Hoff, D.D., Porter, C.W., and Eckhardt, S.G., A phase I and pharmacokinetic study of SAM486A, a novel polyamine biosynthesis inhibitor, administered on a daily schedule to patients with solid tumors. J. Nat'l Cancer Inst.. J. Clinical Oncol. (submitted).
- Vujcic, S., Halmekytö, M., Jänne, J., Porter, C.W. Conditional overexpression of spermidine/spermine N¹-acetyltransferase inhibits cell growth and enhances sensitivity to the polyamine analog N¹, N¹¹-diethylnorspermine (DENSPM). J. Biol. Chem. 275:38309-38328, 2000.

F. Conclusions (& Future Directions):

Conclusions of the studies to date are two-fold. Firstly, certain new polyamine analogs express meaningful antitumor activity in a human prostate carcinoma xenografts system. In collaboration with S'LIL Pharmaceuticals, we propose to pursue further preclinical evaluation and development towards clinical trial. We further propose to examine the basis for this activity *in vitro* and *in vivo* and to continue to evaluate the antitumor activity of new analogs rationally designed and synthesized by S'LIL on the basis of findings to date. Secondly, the polyamine catabolic enzyme SSAT has potential as a gene modulator of polyamine analog antiproliferative activity and as a direct inhibitor of cell growth. We propose to examine whether these properties offer novel potential for achieving pharmacological selectivity and potency in prostate model systems.

G. References:

1. Mi, Z., Kramer, D.L., Miller, J.T., Bergeron, R.J., Bernacki, R., and Porter, C.W. Human prostatic carcinoma cell lines display altered regulation of polyamine transport in response to polyamine analogs and inhibitors. *The Prostate* 34:51-60, 1998
2. Danzin, C., Marchal, P. and Casara, P., Irreversible inhibition of rat S-adenosylmethionine decarboxylase by 5' {[Z]-4-amino-2-butenyl}methylamino}-5'-deoxyadenosine. *Biochem. Pharmacol.* 40:1499-1503, 1990.
3. Regenass, U., Caravatto, G., Mett, H., Stanek, J., Schneider, P., Muller, M., Matter, A., Vertino, P., and Porter, C.W. New S-adenosylmethionine decarboxylase inhibitor with potent antitumor activity. *Cancer Res.* 52:4712-4718, 1992
4. Regenass, U., Mett, H., Stanek, J., Muller, M., Kramer, D., and Porter, C.W. CGP-48664, a new S-adenosylmethionine decarboxylase inhibitor with broad-spectrum antiproliferative and antitumor activity. *Cancer Res.* 54:3210-3217, 1994
5. Marton, L.J., and Pegg, A.E. Polyamines as targets for therapeutic intervention. *Annu. Rev. Pharmacol.*, 35: 55-91, 1995
6. Bernacki, R.J., Bergeron, R.J. and Porter, C.W. Antitumor activity of N¹,N¹²-bis(ethyl)spermine homologs against human MALME-3 melanoma xenografts. *Cancer Res.* 52:2424-2430, 1992.
7. Porter, C.W., Bernacki, R.J., Miller, J. and Bergeron, R.J. Antitumor activity of N¹,N¹¹-bis(ethyl)-norspermine against human melanoma xenografts and possible biochemical correlates of drug action. *Cancer Res.* 53:581-586, 1993.
8. Schipper, R.G., Deli, G., Deloyer, P., Lange, W.P.H.J., Schalken, J.A., and Verhofstad, A.A.J., Antitumor activity of the polyamine analog N¹, N¹¹-diethylnorspermine against human prostate carcinoma. *The Prostate* 44:313-321, 2000.
9. Gschwend, J.E., Fair, W.R., and Powell, C.T., Evaluation of the tetracycline-repressible transactivator system for inducible gene expression in human prostate cancer cell lines. *The Prostate* 33:166-176, 1997.
10. Vujcic, S., Halmekyto, M., Jänne, J., Porter, C.W. Conditional overexpression of spermidine/spermine N¹-acetyltransferase inhibits cell growth and enhances sensitivity to the polyamine analog N¹, N¹¹-diethylnorspermine (DENSPM). *J. Biol. Chem.* 275:38309-38328, 2000.

APPENDIX 1

FIGURE LEGENDS & FIGURES

Figure 1. Structure of the S-adenosylmethionine decarboxylase inhibitor, CGP-48664, and the polyamine analogs, DENSPM, SL-11093 and SL-11144.

Figure 2. Dose-response determination of IC₅₀ values for SL-11144 and SL-11093. Note that SL-0093 reaches a maximum inhibition at 1 μ M.

Figure 3. Time-dependence of growth inhibition by SL-11144 and SL-11093. Note that the former analog is much more toxic than SL-11093 which appears to be wholly cytotoxic at the doses used.

Figure 4. Initial experiment showing the effect of single round of SL-11093 treatment on the growth of DU-145 human prostate carcinoma xenografts. The analog was administered intraperitoneally at 25 and 50 mg/kg/d x 5 days (minimally toxic dose). Thin lines represent individual tumor volumes and thick lines represent median tumor volumes.

Figure 5. Experiment showing the effect of multiple (3) round of SL-11093 treatment on the growth of DU-145 human prostate carcinoma xenografts. The analog was administered intraperitoneally at 50 mg/kg/d x 5 days (minimally toxic dose) with a 9 day respite between treatment rounds. Thin lines represent individual tumor volumes and thick lines represent median tumor volumes. Panel D shows animal weight loss during and following treatment.

Figure 6. Experiment showing the effect of multiple (3) round of SL-11144 treatment on the growth of DU-145 human prostate carcinoma xenografts. The analog was administered intraperitoneally at 5 mg/kg/d x 5 days (minimally toxic dose). Thin lines represent individual tumor volumes and thick lines represent median tumor volumes. Panel D shows animal weight loss during and following treatment.

Figure 7. Repeat experiment showing the effect of multiple (3) round of SL-11093 or SL-11144 treatment on the growth of DU-145 human prostate carcinoma xenografts. The analog was administered intraperitoneally at 50 or 5 mg/kg/d x 5 days (respectively) with a 9 day respite between treatment rounds. Thin lines represent individual tumor volumes and thick lines represent median tumor volumes. Panel D shows animal weight loss during and following treatment. *This Figure emphasized Days 20-98.*

Figure 8. Same experiment as in Figure 7 showing data during Days 20-175.

Figure 9. Experiment showing the effect of a single round of SL-11093 treatment on the growth of PANC-1 human pancreatic carcinoma xenografts. The analog was administered intraperitoneally at 50 mg/kg/d x 5 days (minimally toxic dose). Thin lines represent individual tumor volumes and thick lines represent median tumor volumes. Panel D shows animal weight loss during and following treatment.

Figure 10. DENSPM dose-response for H-10 and M-3 cells grown $\pm$ Dox . Cells were grown $\pm$ Dox for 2 days and then exposed to increasing concentrations of DENSPM for an additional 2 days. Data points represent mean values $\pm$ standard deviation.


Figure 11. Effects of conditional overexpression of SSAT on DENSPM-induced growth inhibition of H-10 cells (left panel) and M-3 cells (right panel). Dox was removed at -2 days and 10 μ M DENSPM (Rx) treatment began at day 0 for both transfectants. These experiments were performed simultaneously with those shown in Figure 8 where the growth effects of -Dox alone are presented. Data points represent mean values $\pm$ standard deviation.

Figure 12. Effects of conditional overexpression of SSAT on clone H-10 (left panel) and clone M-3 (right panel) cell growth. Data points represent mean values $\pm$ standard deviation.

Figure 13. Northern blot analysis of SSAT mRNA expression in LNGK9 clones transfected with a Tetracycline regulatable human SSAT cDNA. Note that overexpression ranges from 5- to 35-fold in the absence of Doxycycline.

Figure 14. Northern blot analysis of SSAT-IRES-GFP mRNA expression in 293 cells transduced with SSAT/AdCMV5-IRES-GFP adenoviral clones. Note that clone 1 alone transduces high levels of SSAT-IRES-GFP mRNA expression.

Figure 1. Structures of Polyamine Analogs & Inhibitors




CGP-48664



DENSPM



SL-11093



SL-11144

**Figure 2. Dose-Response Determination in DU-145
Prostate Carcinoma Cells**




Figure 3. Growth-Response Determination in DU-145 Prostate Carcinoma Cells




Figure 4. DU-145 Prostate Carcinoma Xenografts, Rx SL-11093




Figure 5. DU-145 Prostate Carcinoma Xenografts, Rx SL-11093




Figure 6. DU-145 Prostate Carcinoma Xenografts, Rx SL-11144




Figure 8. DU-145 Prostate Carcinoma Xenografts (Days 20 to 175)




Figure 9. Panc-1 Pancreatic Carcinoma Xenografts, Rx SL-11093






Figure 10. SSAT Inhibition of Cell Growth



**Figure 11. Dose-Dependent Effects of SSAT Expression
on DENSPM Cell Growth Inhibition**



**Figure 12. Effects of SSAT Expression on
DENSPM Cell Growth Inhibition**




Figure 13. *SSAT RNA expression in LNGK9 clones transfected with tet-regulatable human SSAT cDNA*





Figure 14. SSAT-IRES-GFP mRNA in 293 cells transduced with SSAT/AdCMV5-IRES-GFP adenoviral clones





DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 Feb 03

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

ADB263458	ADB282838
ADB282174	ADB233092
ADB270704	ADB263929
ADB282196	ADB282182
ADB264903	ADB257136
ADB268484	ADB282227
ADB282253	ADB282177
ADB282115	ADB263548
ADB263413	ADB246535
ADB269109	ADB282826
ADB282106	ADB282127
ADB262514	ADB271165
ADB282264	ADB282112
ADB256789	ADB255775
ADB251569	ADB265599
ADB258878	ADB282098
ADB282275	ADB232738
ADB270822	ADB243196
ADB282207	ADB257445
ADB257105	ADB267547
ADB281673	ADB277556
ADB254429	ADB239320
ADB282110	ADB253648
ADB262549	ADB282171
ADB268358	ADB233883
ADB257359	ADB257696
ADB265810	ADB232089
ADB282111	ADB240398
ADB273020	ADB261087
ADB282185	ADB249593
ADB266340	ADB264542
ADB262490	ADB282216
ADB266385	ADB261617
ADB282181	ADB269116
ADB262451	
ADB266306	
ADB260298	
ADB269253	
ADB282119	
ADB261755	
ADB257398	
ADB267683	
ADB282231	
ADB234475	
ADB247704	
ADB258112	
ADB267627	