

UNCLASSIFIED

AD NUMBER	
ADC008372	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	confidential
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; JUN 1971. Other requests shall be referred to U.S. Naval Oceanographic Office, Washington, DC 20390.
AUTHORITY	
30 Jun 1983, Group-4, per document marking, DoDD 5200.10; ONR ltr, 31 Jan 2006	

THIS PAGE IS UNCLASSIFIED

000393

CONFIDENTIAL

IR NO. 71-2

GOV LIBRARY COPY

INFORMAL REPORT

MOST Project -2

ADC008372

SOUND VELOCITY AND BOTTOM CHARACTERISTICS FOR LRAPP ATLANTIC AREAS I, II, AND III(U)

See 1473

COPY AVAILABLE TO DDC DOES NOT PERMIT FULLY LEGIBLE PRODUCTION

JUNE 1971

NOV 11 1973

The opinions and assertions contained in this report are solely those of the author(s) and should not be taken as an official, or inferred official opinion of the Naval Oceanographic Office, Department of the Navy, Department of Defense, or United States Government.

This document may be further distributed by any holder only with specific prior approval of the Naval Oceanographic Office

AD No.

DDC FILE COPY

"Unauthorized Disclosure Subject to Criminal Sanctions"

GROUP 4
DOWNGRADED AT 3 YEAR INTERVALS;
DECLASSIFIED AFTER 12 YEARS

NAVAL OCEANOGRAPHIC OFFICE
WASHINGTON, D. C. 20390

Classified by D.D. Form 257

CONFIDENTIAL

001373

1977

710726 0193

CONFIDENTIAL

INFORMAL REPORT

The Informal Report (IR) as produced at the Naval Oceanographic Office is a means for personnel to issue timely scientific and technical preliminary reports of their investigations. These are primarily informal documents used to report preliminary findings or useful byproducts of investigations and work to members of the scientific and industrial communities.

Informal Reports are assigned sequential numbers for each calendar year; the digits preceding the dash indicate the year.

The distribution made of this report is determined primarily by the author. Information concerning obtaining additional copies or being placed on a distribution list for all future Informal Reports in a given area of interest or specialty field, should be obtained from:

Field Management and
Dissemination Department
Code 4420
Naval Oceanographic Office
Washington, D.C. 20390

ACCESSION FOR	
NTIS	☒
DDP	☐
<i>Letter on file</i>	
9	

CONFIDENTIAL

UNCLASSIFIED

CONFIDENTIAL

ABSTRACT (U)

Inventories and summaries of sound velocity, bathymetry, and bottom characteristics have been compiled for three areas in the North Atlantic Ocean between 15° and 65° N latitude and east of 50° to 65° W longitude. The following information is contained in this report: seasonal inventories of sound velocity profiles extending deeper than deep axial depth, charts of the seasonal extent and average axial depth of the upper sound channel, charts of the annual 'strength' of the upper sound channel; charts of the annual extent and average depth of the subsurface sound velocity maximum, annual contour charts of deep axial depth; charts showing bathymetry shoaler than critical depth for summer and winter; an index of the best available bathymetric contour charts; inventories of surficial bottom sediment samples; charts of surficial bottom sediment analysis by grain size classes; an inventory of bottom cores; and a partial inventory of continuous seismic profiles. In addition, a brief analysis is given of water masses that effect sound velocity structures.

DON F. FENNER
PAUL J. BUCCA
CHARLES L. DAVIS

Undersea Surveillance Oceanographic Project
Ocean Science Department
Science and Engineering Center

This report has been reviewed and approved for release as a
CONFIDENTIAL Informal Report.

W B Randlett

W. B. RANDLETT

Director

Undersea Surveillance Oceanographic Project

DATE: 15 June 1971

CONFIDENTIAL

UNCLASSIFIED

iii

(This page 1)

62-0.26 0193

UNCLASSIFIED

TABLE OF CONTENTS (U)

	Page
INTRODUCTION	1
SOUND VELOCITY PROFILE INVENTORIES	3
UPPER SOUND CHANNEL	6
SUBSURFACE SOUND VELOCITY MAXIMUM	9
DEEP SOUND CHANNEL	11
BATHYMETRY SHOALER THAN CRITICAL DEPTH	12
BATHYMETRIC INDEX CHARTS	13
SURFICIAL BOTTOM SEDIMENTS	13
BOTTOM CORE INVENTORIES	14
CONTINUOUS SEISMIC PROFILE INVENTORIES	14
REFERENCES	117

FIGURES (U)

1. Location of LRAPP Atlantic Priority Areas	2
2. Inclusive Dates for NODC Seasonal Sound Velocity Summaries	4
I-1. Location of LRAPP Atlantic Area I Subareas	19
I-2A - I-2C. Number of Winter (Jan-Mar) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	20
I-3A - I-3C. Number of Spring (Apr-Jun) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	23
I-4A - I-4C. Number of Summer (Jul-Sep) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	26

FIGURES (CONTINUED)

	Page
I-5A - I-5C. Number of Autumn (Oct-Dec) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	29
I-6. Areal Extent and Average Axial Depth of Upper Sound Channel for Winter (Jan-Mar)	32
I-7. Areal Extent and Average Axial Depth of Upper Sound Channel for Spring (Apr-Jun)	33
I-8. Areal Extent and Average Axial Depth of Upper Sound Channel for Summer (Jul-Sep)	34
I-9. Areal Extent and Average Axial Depth of Upper Sound Channel for Autumn (Oct-Dec)	35
I-10. Annual Average "Strength" of Upper Sound Channel (if present) Relative to Subsurface Sound Velocity Maximum	36
I-11. Annual Areal Extent and Average Axial Depth of Subsurface Sound Velocity Maximum	37
I-12. Annual Average Depth of Deep Sound Axis (in meters X 100)	38
I-13. Bathymetry Shoaler than Average Critical Depth for Winter (Nov-Apr) (Contour interval = 500 fathoms)	39
I-14. Bathymetry Shoaler than Average Critical Depth for Summer (May-Oct) (Contour interval = 500 fathoms)	40
I-15. Index of Best Existing Bathymetric Charts	41
I-16A - I-16C. Number of Bottom Sediment Samples per One-degree Square	42
I-17. Distribution of Surficial Bottom Sediments	45
I-18A - I-18C. Number of Bottom Cores per One-degree Square	46
I-19A - I-19C. Partial Inventory of Continuous Seismic Profiles	49
II-1. Location of LRAPP Atlantic Area II Subareas	55

UNCLASSIFIED

FIGURES (CONTINUED)

	Page
II-2A - II-2C. Number of Winter (Jan-Mar) Oceanographic Observations per One-Degree Square Deeper than Deep Axial Depth	56
II-3A - II-3C. Number of Spring (Apr-Jun) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	59
II-4A - II-4C. Number of Summer (Jul-Sep) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	62
II-5A - II-5C. Number of Autumn (Oct-Dec) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	65
II-6. Areal Extent and Average Axial Depth of Upper Sound Channel for Winter (Jan-Mar)	68
II-7. Areal Extent and Average Axial Depth of Upper Sound Channel for Spring (Apr-Jun)	69
II-8. Areal Extent and Average Axial Depth of Upper Sound Channel for Summer (Jul-Sep)	70
II-9. Areal Extent and Average Axial Depth of Upper Sound Channel for Autumn (Oct-Dec)	71
II-10. Annual Average "Strength" of Upper Sound Channel (if present) Relative to Subsurface Sound Velocity Maximum	72
II-11. Annual Areal Extent and Average Axial Depth of Subsurface Sound Velocity Maximum	73
II-12. Annual Average Depth of Deep Sound Channel Axis (in meters X 100) .	74
II-13. Bathymetry Shoaler than Average Critical Depth for Summer (Nov-Apr) (Contour interval = 500 fathoms)	75
II-14. Bathymetry Shoaler than Average Critical Depth for Summer (May-Oct) (Contour interval = 500 fathoms)	76
II-15. Index of Best Existing Bathymetric Charts	77

UNCLASSIFIED

(FIGURES (CONTINUED))

	Page
II-15A - II-16C. Number of Bottom Sediment Samples per One-degree Square	78
II-17. Distribution of Surficial Bottom Sediments	81
II-18A - II-18C. Number of Bottom Cores per One-degree Square	82
II-19A - II-19C. Partial Inventory of Continuous Seismic Profiles	85
III-1. Location of LRAPP Atlantic Area III Subareas	91
III-2A - III-2B. Number of Winter (Jan-Mar) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	92
III-3A - III-3B. Number of Spring (Apr-Jun) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	94
III-4A - III-4B. Number of Summer (Jul-Sep) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	96
III-5A - III-5B. Number of Autumn (Oct-Dec) Oceanographic Observations per One-degree Square Deeper than Deep Axial Depth	98
III-6. Areal Extent and Average Axial Depth of Upper Sound Channel for Winter (Jan-Mar)	100
III-7. Areal Extent and Average Axial Depth of Upper Sound Channel for Spring (Apr-Jun)	101
III-8. Areal Extent and Average Axial Depth of Upper Sound Channel for Summer (Jul-Sep)	102
III-9. Areal Extent and Average Axial Depth of Upper Sound Channel for Autumn (Oct-Dec)	103
III-10. Annual Average "Strength" of Upper Sound Channel (if present) Relative to Subsurface Sound Velocity Maximum	104
III-11. Annual Areal Extent and Average Axial Depth of Subsurface Sound Velocity Maximum	105

UNCLASSIFIED

FIGURES (CONTINUED)

	Page
III-12. Annual Average Depth of Deep Sound Channel Axis (in meters X 100)	106
III-13. Bathymetry Shoaler than Average Critical Depth for Winter (Nov-Apr) (Contour interval = 500 fathoms)	107
III-14. Bathymetry Shoaler than Average Critical Depth for Summer (May-Oct) (Contour interval = 500 fathoms)	108
III-15. Index of Best Existing Bathymetric Charts	109
III-16A - III-16B. Number of Bottom Sediment Samples per One-degree Square	110
III-17. Distribution of Surficial Bottom Sediments	112
III-18A - III-18B. Number of Bottom Cores per One-degree Square	113
III-19A - III-19B. Partial Inventory of Continuous Seismic Profiles	115

UNCLASSIFIED

INTRODUCTION

U By a Memorandum for Undersea Surveillance Oceanographic Center dated 6 July 1970, ONR Code 102-OS requested that data inventories and summaries of sound velocity, bathymetry, and bottom characteristics be provided for three priority areas in the North Atlantic Ocean (see Figure 1). NAVOCEANO letter ser 3768 of 4 August 1970 stated that the following information would be provided:

- Sound velocity profile inventories per one-degree square per standard three-month season
- Areal contours of the deep sound channel axis on an annual basis
- Areal contours of the upper sound channel axis and depth of the subsurface sound velocity maximum on a seasonal and/or annual basis along with an indication of the "strength" of the upper sound channel
- Charts showing the height and extent of bottom features shallower than critical depth for summer and winter
- Index charts showing the most accurate available bathymetric charts and their compilation dates
- Charts showing the distribution of surficial sediments classified by principle grain size constituents
- Charts showing a partial inventory of bottom cores.

In addition, a partial inventory of continuous seismic profiles is included.

(U) Information on the parameters cited above has been organized in three sections, each corresponding to one of the three priority areas shown on Figure 1. In order to maintain a page size presentation, the three areas have been broken into a total of eight subareas. Figure designations are consistent according to the following system:

- Roman numerals designate the priority area (e.g., I, II, or III)
- Arabic numerals designate the parameter within each priority area (e.g., the number of winter oceanographic observations per one-degree square deeper than deep axial depth is Figure I-2, II-2, or III-2)

UNCLASSIFIED




FIGURE 1: LOCATION OF LRAPI
ATLANTIC PRIORITY AREAS

1 of 2

2

UNCLASSIFIED

- Letters designate subareas within priority areas (e.g., Figures I-2A, I-2B, and I-2C).

(U) The locations of these subareas are shown on Figures I-1, II-1, and III-1. For purposes of this report, deep axial depth is defined as the deepest sound velocity minimum (usually absolute minimum). Critical depth is defined as that depth where the sound velocity is equal to the maximum sound velocity found at the surface or in the surface mixed layer.

SOUND VELOCITY PROFILE INVENTORIES

(U) Figures I-2A through I-5C, II-2A through II-5C, and III-2A through III-5B show the number of existing sound velocity observations per one-degree square for the four standard seasons of winter (January through March), spring (April through June), summer (July through September), and autumn (October through December). All observations shown on these figures extend deeper than the deep sound channel (see Figures I-12, II-12, and III-12). The four standard seasons were chosen since the area includes tropical to subarctic conditions. These figures are based upon the following data sources:

- National Oceanographic Data Center (NODC) seasonal sound velocity summaries containing all Nansen cast data extending deeper than 50 meters processed as of the dates shown on Figure 2
- Sound velocimeter and Nansen cast data from the following NAVOCEANO Marine Geophysical Survey (MGS) Atlantic Task Areas:
 - Alpine Area SF (NAVOCEANO, Dec 1966)
 - Alpine Area ST (NAVOCEANO, Aug 1968)
 - Alpine Area I (NAVOCEANO, Sep 1966a and Feb 1967a)
 - Texas Instruments (TI) Area 2 (NAVOCEANO, Feb 1968)
 - TI Area 3 (NAVOCEANO, May 1968)
 - TI Area 5 (NAVOCEANO, Feb 1967b and Jun 1967)
 - TI Area 7/4 (NAVOCEANO, May 1969)
- Sound velocimeter data, furnished by Woods Hole Oceanographic Institution, from the following cruises:
 - R/V ATLANTIS Cruise No. 282, Jul 1962
 - R/V ATLANTIS II Cruise No. 11, Jun 1964




FIGURE 2

UNCLASSIFIED

- R/V ATLANTIS II Cruise No. 22, Jul-Aug 1966 (Beckerle, Oct 1968)
- R/V CHAIN Cruise No. 82, Aug-Sep 1968 (Katz, Jun 1969)
- R/V CHAIN Cruise No. 89, Mar 1969 (LaCasce, et al., Jun 1969)
- Sound velocimeter data taken by Lamont-Doherty Geological Observatory during Nov-Dec 1963, Apr-May 1964, and Jun-Jul 1965 (Piip, Jan 1966, Jun 1967, and Apr 1968, respectively)
- Sound velocimeter data taken by the USNS KANE during Jun-Jul 1970 as part of Phase I of the Northeast Atlantic Basin (NEAB) surveys (courtesy of C. Ostericher, Deep Ocean Surveys Division, NAVOCEANO)
- All available Nansen cast and salinity-temperature-depth (STD) recorder data taken as part of the Barbados Oceanographic and Meteorological Experiment (BOMEX) by the following ships:
 - USCGSS OCEANOGRAPHER, USCGC RAINIER, and USCGC (Environmental Science Services Administration, Dec 1969)
 - USNS GILLIS and R/V ADVANCE II (courtesy of P. Mazeika, NAVOCEANO BOMEX Coordinator)
- Additional Nansen cast and STD data taken in support of the International Ice Patrol by the following ships:
 - USCGC EVERGREEN, Apr-Jul 1966 (Wolford, Jun 1966)
 - USCGC EDISTO, Sep-Oct 1967 (Codispoti and Kravitz, May 1968)
 - USCGC WESTWIND, Sep-Oct 1969 (Bunce, Apr 1970)
- Additional Nansen cast data taken in support of the International Commission for the Northwest Atlantic Fisheries (ICNAF) in 1963 as follow:
 - at OCEAN WEATHER STATION POLAR FRONT I and II (same as OWS ALFA), Jun-Jul
 - by RS THALASSA, Apr
 - by CV AEGIR, May
 - by RS ACADEMICIAN KNIPOVITCH, Jun
 - by USCGC EVERGREEN, Jul
 - by R/V EXPLORER, Jul

(published by ICNAF, 1968)

UNCLASSIFIED

- Additional Canadian Nansen cast data taken by the following ships:
 - CSS HUDSON, Mar-Jul 1966 (Canadian Oceanographic Data Centre (CODC), 1969a)
 - CSS BAFFIN, Apr-May 1966 (CODC, 1969b)
 - CSS HUDSON, Jan-Mar 1967 (CODC, 1969c)
- Fleet Numerical Weather Central (Monterey) expendable bathythermograph (XBT) data processed as of 1 June 1970
- Additional XBT, sound velocimeter, and Nansen cast data taken as part of the first Northeast Atlantic (NEAT I) experiment during Sep-Oct 1969 by the USNS GIBBS, R/V CHAIN and ACS ST. MARGARETS (courtesy of Acoustics Division, NRL)
- Additional XBT data taken by the USNS GIBBS during May-Jun 1969 and Apr 1970 (courtesy of H. Fleming and G. Shaffer, Acoustics Division, NRL).

In addition, all available Nansen cast and sound velocimeter data taken by Project CAESAR in the North Atlantic Ocean between 1954 and 1967 (indexed by Bunce, Aug 1969) and by the NAVOCEANO Ocean Survey Program (OSP) are included in these figures. Nansen cast and STD data were converted into sound velocity profiles using the equation of Wilson (1960). Various XBT data were converted into sound velocity profiles using the equation of Wilson and historical salinity correction factors.

UPPER SOUND CHANNEL

(U) Figures I-6 through I-9, Figures II-6 through II-9, and Figures III-6 through III-9 show the areal extent and average axial depth of the upper sound channel for each of the four standard seasons. In constructing these figures, seasonal upper axial depths were compiled by one-degree square and averaged on a two-degree square (i.e., four one-degree squares) basis. Two-degree square averages then were contoured on an areal basis. Regions where an upper sound channel is present more than 80% of the time, 20-80% of the time, and less than 20% of the time are indicated on each figure. For purposes of the following discussion, the first case above will be referred to as a permanent upper sound channel, the second as transitory. In the third case, an upper sound channel effectively is absent.

(U) During winter, a permanent upper sound channel is present only east of about 20° W longitude between Cape Finisterre and the Canary Islands (see

UNCLASSIFIED

Figure II-6). A transitory upper sound channel surrounds this region north to about 49° N latitude and west to the Azores Islands. A transitory upper sound channel also is present during winter in a small area east of Grand Banks (see Figure I-6) and throughout the Sargasso Sea (see Figure III-6). In the Canary and Iberian Basins, upper axial depths range from approximately 300 meters near Cape Finisterre to greater than 900 meters north of the Canary Islands. South of the Canary Islands, the upper axial depths decrease. Axial depths in the Sargasso Sea and off Grand Banks range between 100 and 300 meters during winter, and are considerably shallower than those in the Canary Basin. This situation prevails during all seasons.

(U) During spring, there is a permanent upper sound channel east of about 20°W longitude between the Canary Islands and Iceland (see Figures I-7 and II-7) and in a small area east of Bermuda (see Figure III-7). However, during spring there are transitory upper sound channels north of about 25° N latitude except over the Mid-Atlantic Ridge and in the Labrador Basin. In the Canary and Iberian Basins, upper axial depths range from less than 300 meters off Cape Finisterre to greater than 900 meters in the vicinity of the Canary Islands. Throughout the remainder of the three areas, upper axial depths range between 100 and 200 meters.

(U) During summer, there is a permanent upper sound channel east of about 20° W longitude between the Canary Islands and Iceland, throughout the basin south of Iceland, and across the Reykjanes Ridge into the central Labrador Basin (see Figures I-8 and II-8). There is also a permanent upper sound channel in the Sargasso Sea (see Figure III-8). The summer extent of the transitory upper sound channel is similar to that during spring, with two exceptions: transitory upper sound channels are present farther to the west in the Labrador Basin during summer but are absent over larger sections of the Mid-Atlantic Ridge north of the Azores Islands. Upper axial depths in the Canary and Iberian Basins are approximately the same during spring and summer (300-900 meters). However, throughout Areas I and III summer axial depths range between 150 and 200 meters; while in Area II, upper axial depths are greater than 200 meters.

(U) During autumn, a permanent upper sound channel is present east of about 20° W longitude between the Canary Islands and the Faeroe Plateau (see Figures I-9 and II-9) and in a local area east and south of Bermuda (see Figure III-9). The extent of transitory upper sound channel is similar during spring and autumn. Upper axial depths in the Canary and Iberian Basins are similar to those during spring and summer. However, in the region north of Cape Finisterre and east of the Mid-Atlantic Ridge, autumn axial depths average about 200 meters (the deepest during the year). In the Sargasso Sea, autumn axial depths also are approximately 50 meters deeper than during summer. In the western

UNCLASSIFIED

half of Area I, autumn data are either insufficient to evaluate or insufficient to contour. However, axial depths probably are somewhat deeper during autumn than summer in this region.

(U) The causes of an upper sound channel east of the Mid-Atlantic Ridge have been described by Fenner and Bacon (Dec 1969). Briefly, south of about 43° N latitude (Cape Finisterre), an upper sound channel is formed by the interaction of relatively warm, saline Mediterranean Intermediate Water (MIW) with colder, more dilute North Atlantic Central Water (NACW) carried by an extension of the North Atlantic Current. In this case, upper axial depths coincide with the bottom of the NACW layer. North of about 50° N latitude, an upper sound channel is formed by warming of surface and near-surface layers characterized by permanent or transient positive velocity gradients during winter. In this case, upper axial depths correspond roughly with the maximum depth of summer warming. However, in cases where the maximum depth of summer warming exceeded the maximum depth of winter cooling, an upper sound channel was not found (e.g., south of the Canary Islands and over the Mid-Atlantic Ridge). Between 43° and 50° N latitude, an upper sound channel is formed by warming of surface layers combined with interaction of MIW and NACW. This latter region is characterized by a relatively strong oceanic front from the surface to depths in excess of 1200 meters caused by the interaction of NACW; a cold, dilute flow at depth (Arctic Intermediate Water - AIW); and MIW. Transitory upper sound channels generally were found in regions where various causative forces were attenuated (e.g., regions of MIW dilution, regions where winter cooling is attenuated by warm surface currents, etc.).

(U) West of the Mid-Atlantic Ridge (western half of Area I and Area III), upper sound channels apparently are caused by spring, summer, and autumn warming of the surface and near-surface layers. Such a hypothesis explains the lack of a widespread upper sound channel during winter throughout Area I and the greater extent of a transitory upper sound channel in the Labrador Basin during summer than during spring. Preferential summer warming in the regions influenced by the Irminger Current may explain the permanent upper sound channel west of the Reykjanes Ridge during summer. The localized occurrence of a transitory upper sound channel east of the Grand Banks during winter probably is a result of intensive mixing of waters carried by the North Atlantic and Labrador Currents. In the Sargasso Sea, an upper sound channel was associated with the "18° Water" of Worthington (1959). Somewhat cooler winter conditions near Bermuda and the relative absence of "18° Water" during winter throughout the Sargasso Sea may explain the absence of a permanent upper sound channel during winter in Area III.

UNCLASSIFIED

(U) Figures I-10, II-10, and III-10 show the annual average "strength" of the upper sound channel (if present) relative to the subsurface sound velocity maximum. This parameter was compiled by one-degree square and season, averaged by two-degree square on an annual basis, and generalized into seven categories. It is presented by two-degree squares on an annual basis because of the large temporal and spatial variability throughout the three areas. Generally, "A", "B", or "C" values (i.e., 6 to greater than 10 meters/second) indicate the effects of high concentrations of MIW and/or extreme winter cooling followed by intensive summer warming. "D" and "E" values (i.e., 2 to 6 meters/second) generally indicate the effects of intermediate concentrations of MIW and/or average winter cooling and summer warming conditions. "F" and "G" values (0.1 to 2 meters/second) generally are associated with transitory upper sound channels and/or subsurface sound velocity maxima and indicate attenuation of causative forces. The reader is cautioned that two-degree square values of this parameter often mask specific seasonal and areal tendencies.

SUBSURFACE SOUND VELOCITY MAXIMUM

(U) Figures I-11, II-11, and III-11 show the annual areal extent and average axial depth of the subsurface sound velocity maximum. This parameter was compiled by one-degree square and season, averaged by two-degree square on an annual basis, and then contoured on an areal basis. An annual presentation was chosen for this parameter since seasonal two-degree square averages vary by less than 100 meters from comparable annual averages throughout most of the three areas. However, in the vicinity of the Canary Islands, from 43° and 50° N latitude between about 15° and 30° W longitude, on either side of Iceland, and along 40° N latitude between 50° and 65° W longitude there are large seasonal variations in the depth of the subsurface sound velocity maximum. All of these regions roughly correspond to frontal zones either at the surface, at intermediate depths, or both. Regions where a subsurface sound velocity maximum is present greater than 80% of the time, 20-80% of the time, and less than 20% of the time are indicated on each figure. For purposes of the following discussion, the first case above will be referred to as a permanent subsurface sound velocity maximum and the second as a transitory subsurface sound velocity maximum. In the third case, a subsurface sound velocity maximum effectively is absent.

(U) A permanent subsurface sound velocity maximum is present throughout the year east of about 20° W longitude between the Canary Islands and Iceland, throughout the basin south of Iceland, and across the Reykjanes Ridge into the central Labrador Basin (see Figures I-11 and II-11). This areal extent is similar to that of the permanent upper sound channel during summer. There is also a

UNCLASSIFIED

permanent subsurface sound velocity maximum in separate regions south and east of the Grand Banks and in the central Sargasso Sea (see Figures I-11 and III-11). Transitory subsurface sound velocity maxima are found throughout all three areas north of about 25° N latitude except over the Mid-Atlantic Ridge, in a region directly off Grand Banks, and in the northern Labrador Basin. The annual average depth of the subsurface sound velocity maximum east of the Mid-Atlantic Ridge decreases to the north from 1300 meters near the Canary Islands to less than 500 meters south of Iceland. West of the Mid-Atlantic Ridge, depths range from less than 200 meters in the north to greater than 400 meters in the central Sargasso Sea.

(U) In the Northeast Atlantic (east of 30° W longitude), subsurface sound velocity maxima are caused solely by MIW south of about 43° N latitude, by winter cooling north of about 50° N latitude, and by a combination of both factors between about 43° and 50° N latitude and in the deep water channel between the Faeroe Plateau and Porcupine Bank (Fenner and Bucca, Dec 1969). In the first region, the depth of this parameter coincided well with the depth of the MIW high salinity core (salinity maximum). In the second region, the depth of the subsurface sound velocity maximum coincided roughly with the maximum depth of winter cooling. In the third region, MIW was found to accentuate the depth of subsurface sound velocity maxima originally formed by winter cooling.

(U) West of 30° W longitude (i.e., in the western half of Area I and in Area III), subsurface sound velocity maxima are considerably shallower than those in the Northeast Atlantic and roughly coincide either with layer depths or the bottom of the "18° Water" layer. Therefore, both the temporal and spatial variability of this parameter are greater west of the Mid-Atlantic Ridge. Despite intensive winter cooling throughout the Labrador Sea (to depths in excess of 500 meters), subsurface sound velocity maxima similar to those north of Porcupine Bank are not present in the western half of Area I. This is largely due to extremely low salinity water carried by the Labrador and West Greenland Currents (less than 34.9 ‰ according to Lee and Ellett, 1967). In this region, winter cooling causes positive velocity gradients throughout the water column (see Figure I-12), and summer warming causes a deep sound channel axis at a depth somewhat below the absolute salinity minimum (often at the surface). Therefore, a subsurface sound velocity maximum is not formed despite intensive winter cooling. The tongue with subsurface sound velocity maxima at greater than 400 meters that lies west of the Reykjanes Ridge at about 60° N latitude is apparently related to higher salinity and warmer waters that are transported by the Irminger Current. This tongue corresponds well with the shape of isohalines and isotherms shown by Wüst and Defant (1936) at 400 and 600 meters.

UNCLASSIFIED

(U) The Polar Front is shown clearly in the rapid transition of subsurface sound velocity maxima on either side of Iceland. Transitional regions also are found along the northern boundary of Area III (northern wall of Gulf Stream), in the region extending from about 43° to 50° N latitude between about 15° and 30° W longitude (frontal zone caused by interaction of NACW, AIW, and MIW), and in the vicinity of the Canary Islands (region of rapid dilution of MIW). In all these regions of rapid transition, the surface sound velocity maximum tends to merge with either the upper or deep sound channel to form a basically trilinear sound velocity profile.

DEEP SOUND CHANNEL

(U) Figures I-12, II-12, and III-12 show contours of the annual axial depth of the deep sound channel. This parameter was compiled and contoured in a manner similar to that used for the subsurface sound velocity maximum. An annual presentation was chosen for deep axial depth since in any given two-degree square seasonal variations were as great as annual variations. In addition, deep axial depths at any given latitude generally are not directly related to the annual heating and cooling cycle (i.e., deeper values in summer, shallower values in winter). A deep sound channel is found throughout the year in all three areas except in regions off Grand Banks and south and east of Greenland (see Figure I-12). In these two regions, intensive cooling results in positive velocity gradients during the colder months of January through April. During the summer, a deep sound channel is formed in these two regions by warming of the surface and near-surface layers. In a small region off Cape Farvel, positive velocity gradients are present throughout the year.

(U) Annual deep axial depths in the North Atlantic range from greater than 2000 meters off the Iberian coast (see Figure II-12) to less than 100 meters off the Grand Banks and Greenland (see Figure I-12). Two major features are apparent in the overall deep axial depth structure. The first consists of a tongue with anomalously deep values of this parameter that extends west southwest from the Iberian Coast across the North Atlantic to Bermuda. This tongue is bounded by the 1200-meter deep axial depth isoline and its shape corresponds well with the preferential flow of high salinity MIW shown by Worthington (1970). Deep axial depths shoal markedly to the north and south of this tongue. The second major feature consists of a tongue with anomalously shoal values of this parameter that extends along about 55° N latitude from about 40° to about 20° W longitude. This tongue is bounded by the 900-meter deep axial depth isoline and its shape corresponds well with the 35.0 ‰ and less isohalines shown at 400, 600, and 800 meters by Wüst and Defant (1936). The latter tongue corresponds to the flow of low salinity AIW. Deep axial depths tend to deepen to the north and south of 55° N latitude.

UNCLASSIFIED

(U) Zones of rapid change in deep axial depth are found to the east and west of Iceland (Polar Front), south and east of Grand Banks (at confluence of the Gulf Stream, North Atlantic Current system and the Labrador Current), and in the vicinity of the Canary Islands (region of rapid dilution of MIW). The extremely broad deep sound channel south of about 30° N latitude in both Areas II and III apparently is a result of diluted concentrations of high salinity MIW (Fenner and Bucca, Dec 1969). However, in the southwest corner of Area III, the width of the channel is constricted by intrusion of low salinity Antarctic Intermediate Water (AAIW) from the south. In the western half of Area I, the sound channel also is narrower because of low salinity AIW.

BATHYMETRY SHOALER THAN CRITICAL DEPTH

(U) Figures I-13, II-13, and III-13 show the amount of bathymetry extending above average critical depth during the composite six-month winter season (Nov-Apr). Figures I-14, II-14, and III-14 show the same parameter for the composite six-month summer season (May-Oct). These composite seasons were chosen because of the lack of data in various regions of the North Atlantic during one or more of the four standard seasons. In addition, six-month depth difference charts show relatively small changes in 500-fathom contours at three-month intervals. In preparation of these figures, critical depths were compiled by one-degree square for each of the two seasons, averaged by two-degree square, and then contoured on an areal basis. Bathymetric charts then were overlayed with the critical depth contours and areas where bathymetry extended above critical depth were outlined. Except in the Northeast Atlantic between 30° and 60° N latitude east of 30° W longitude, bathymetric charts compiled by NAVOCEANO for the National Intelligence Survey (NIS) prior to 1960 were used in constructing these figures. In the Northeast Atlantic, the charts of A. S. Laughton (National Institute of Oceanography, Wormley, England) were used. Critical depth contours are based on NODC serial sound velocity profiles processed as of January 1969, supplemented by sound velocimeter and Nansen cast data from the MGS and CAESAR programs and sound velocimeter data from Woods Hole Oceanographic Institution and Lamont-Doherty Geological observatory.

(U) During both seasons substantial areas of bathymetry shoaler than critical depth are associated with major physiographic features such as the Mid-Atlantic Ridge, the Reykjanes Ridge, the Faeroe Plateau, Porcupine Bank, various seamounts west of Gibraltar, the Canary Rise, the Bermuda Rise, the New England Seamounts, and the Grand Banks. The amount of bathymetry shoaler than critical depth is less during winter than during summer in all cases because of the shallower critical depths during this season. Of particular interest is the lesser amount of bathymetry shoaler than critical depth associated with the

UNCLASSIFIED

New England Seamounts, Bermuda Rise, and the intersection of the Reykjanes and Mid-Atlantic Ridges (Gibbs Fracture Zone) during winter (compare Figures I-13 and I-14 and Figures III-13 and III-14).

BATHYMETRIC INDEX CHARTS

(U) Figures I-15, II-15, and III-15 are indices of the best available bathymetric contour charts for the three LRAPP priority areas. Four types of charts were used in compiling these figures:

- NAVOCEANO North Atlantic Regional Charts compiled in 1969
- Charts of A. S. Loughton (National Institute of Oceanography, Wormley, England) compiled in 1966
- NAVOCEANO Bathymetric Contour charts (BC's) contoured in the indicated years
- Charts contoured prior to 1960 as part of the National Intelligence Survey (NIS) program.

Individual, large-scale charts of specific locations or bathymetric features have not been included on these indices.

SURFICIAL BOTTOM SEDIMENTS

Figures I-16, II-16, and III-16 show the location by one-degree square of bottom sediment samples held by NAVOCEANO. These figures reflect the same data distribution shown on Figure V-6 of H.O. Publication No. 700 (NAVOCEANO, 1965). Therefore, the bibliography contained in the above publication also applies to the three figures. There are large variations in the quantity and quality of bottom sediment data throughout the three areas. Only a few shallow nearshore regions have been sampled adequately.

Figures I-17, II-17, and III-17 show the classification of surficial bottom sediments on the basis of principle grain size constituents. These figures were taken directly from Figures V-7 and V-8 of H.O. Publication No. 700 (NAVOCEANO, 1965). Generally, coarser sediments are found over the Mid-Atlantic and Reykjanes Ridges and on the continental shelves. Other regions primarily contain muds, sands, and mud-sands.

BOTTOM CORE INVENTORIES

(U) Figures I-18, II-18, and III-18 show the location by one-degree square of unclassified bottom cores held by NAVOCEANO. A recently updated computer plot of these cores was provided by E. Wilcox, Oceanographic Analysis Division, NAVOCEANO. Cores indexed on these figures vary considerably in length and in type and quality of analysis.

CONTINUOUS SEISMIC PROFILE INVENTORIES

(U) Figures I-19, II-19, and III-19 show a partial inventory of continuous seismic profiles of the subbottom on a one-degree square grid. These figures are based on the following data sources.

- All data collected in the following NAVOCEANO MGS Atlantic Task Areas:
 - Alpine Area SF (NAVOCEANO, Feb 1967c)
 - Alpine Area ST (NAVOCEANO, Aug 1968)
 - Alpine Area I (NAVOCEANO, Sep 1966b)
 - Texas Instruments (TI) Area 2 (NAVOCEANO, Mar 1968)
 - TI Area 3 (NAVOCEANO, Oct 1968)
 - TI Area 5 (NAVOCEANO, Jan 1967)
 - TI Area 7/4 (NAVOCEANO, Apr 1969)
- All data collected on Cruise 9 of the USNS KANE (Lowrie and Escowitz, 1969)
- Data collected by the Woods Hole Oceanographic Institution on the following cruises of the R/V CHAIN:
 - No. 7, 1959 (Dunkle and Hays, May 1966)
 - No. 21, 1961 (Ibid.)
 - No. 34, 1962 (Ibid.)
 - No. 36, 1963 (Ibid.)
 - No. 39, 1963 (Ibid.)
 - No. 43, 1964 (Ibid.)
 - No. 44, 1964 (Ibid.)
 - No. 70 and No. 73, 1967 (Knott, et al., Jul 1968)
 - No. 82, 1968 (Dunkle, Jun 1969)

UNCLASSIFIED

- Data collected by the Lamont-Doherty Geological Observatory aboard the R/V VEMA and R/V CONRAD, as contained in the following references:

- Ewing, et al. (1960)
- Wilson (1963)
- Ewing, et al. (1964)
- Jones and Ewing (1969).

Major oceanographic, geological, and geophysical journals and serial publications from major national and international oceanographic institutions and organizations have been searched in preparing these figures. Tracks shown on these figures represent profiles taken with sparker, boomer, and air gun systems.

UNCLASSIFIED

LRAPP PRIORITY AREA ONE

UNCLASSIFIED

UNCLASSIFIED





FIGURE L-1: LOCATION OF LAMP ATLANTIC AREA I SURVEYS

FIGURE 1-2A: NUMBER OF WINTER
(Jan-Mar) OCEANOGRAPHIC
OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP
AXIAL DEPTH



CONFIDENTIAL





FIGURE I-2B: NUMBER OF WINTER (Jan-Mar) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH




240" W
65° N

35°

30°W
60°N

60°N
50°W

60°N
45°W



60

55

50
N

CONFIDENTIAL




FIGURE I-3B: NUMBER OF SPRING (Apr-Jun) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH

CONFIDENTIAL

CONFIDENTIAL




FIGURE I-3C

25

710:26 0193

CONFIDENTIAL

CONFIDENTIAL

FIGURE 1-4A: NUMBER OF SUMMER
(Jul-Sep) OCEANOGRAPHIC
OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP
AXIAL DEPTH




FIGURE 1-4A

CONFIDENTIAL

CONFIDENTIAL




FIGURE I-4B: NUMBER OF SUMMER (Jul-Sep) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP WATER DEPTH

CONFIDENTIAL

30°W

25°

20°

15°

55°
N
10°
W60°
N55°
N50°
N10°
W

FIGURE I-4C

28

CONFIDENTIAL

30°W
65°N

FIGURE 1-5A: NUMBER OF AUTUMN
(Oct-Dec) OCEANOGRAPHIC
OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP
AXIAL DEPTH




FIGURE I-5A

CONFIDENTIAL

710:26 0193




FIGURE I-5B: NUMBER OF AUTUMN (Oct-Dec) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH


30°W

25°

20°

15°

UN
10
W



Εύχρηστο
Νέο

55°
N

50'

10⁹
W

FIGURE I-5C

CONFIDENTIAL

UNCLASSIFIED




FIGURE I-6: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR WINTER (Jan-Mar)

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-7: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR SPRING (Apr-Jun)

UNCLASSIFIED

A map of the North Atlantic Ocean from 30°W to 10°W and 40°N to 65°N. The map displays depth contours (solid lines) and sound channel presence (dashed lines). A legend at the bottom explains the symbols: a solid line for 'Upper sound channel present > 80% of time', a dashed line for 'Average axial depth contours (in meters X 100)', a dash-dot line for 'Upper sound channel present 20-80% of time', and a dotted line for 'Upper sound channel present < 20% of time'. The map shows a large area of dotted pattern (sound channel present < 20% of time) in the central and eastern North Atlantic, with depth contours ranging from 100 to 3000 fathoms. A label '100 FATHOMS' is at the top. Latitude and longitude markers are provided along the edges.

34

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-9: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR AUTUMN (Oct-Dec)

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-10: ANNUAL AVERAGE "STRENGTH" OF UPPER SOUND CHANNEL (if present) RELATIVE TO SURFACE SOUND VELOCITY MAXIMUM

UNCLASSIFIED




FIGURE I-11: ANNUAL AREAL EXTENT AND AVERAGE AXIAL DEPTH OF SUBSURFACE SOUND VELOCITY MAXIMUM

UNCLASSIFIED

UNCLASSIFIED




FIGURE L-12: ANNUAL AVERAGE DEPTH OF DEEP SOUND CHANNEL AXIS
(in meters X 100)

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-13: BATHYMETRY SHOALER THAN AVERAGE CRITICAL DEPTH
FOR WINTER (Nov-Apr)
(Contour interval = 500 fathoms)

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-14: BATHYMETRY SHOALER THAN AVERAGE CRITICAL DEPTH
FOR SUMMER (May-Oct)
(Contour interval = 500 fathoms)

CONFIDENTIAL




FIGURE 1-15: INDEX OF BEST EXISTING BATHYMETRIC CHARTS

UNCLASSIFIED

FIGURE I-16A: NUMBER OF
BOTTOM SEDIMENT SAMPLES PER
ONE-DEGREE SQUARE

LEGEND:

$\frac{1}{3}$ — Number of surface
sediment samples from
cores
Number of surface sediment
samples from snappers,
grabs, etc.




FIGURE I-16A

UNCLASSIFIED




FIGURE I-16B: NUMBER OF BOTTOM SEDIMENT SAMPLES PER ONE-DEGREE SQUARE

LEGEND: see Figure I-16A


UNCLASSIFIED

25°

20°

15°

65°
N
10°
W



450°
N

55°
N

50
N

FIGURE I-16C

UNCLASSIFIED




FIGURE I-17: DISTRIBUTION OF SURFICIAL BOTTOM SEDIMENTS

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-18A: NUMBER OF
BOTTOM CORES PER ONE-DEGREE
SQUARE

FIGURE I-18A

UNCLASSIFIED

UNCLASSIFIED




FIGURE I-18B: NUMBER OF BOTTOM CORES PER ONE-DEGREE SQUARE


30°W

25°.

20°

15°

65°
N
1°
W



60°
N

55°
N

50

PAGE TWO I-1AC

48

19


UNCLASSIFIED

UNCLASSIFIED

FIGURE I-19A: PARTIAL INVENTORY
OF CONTINUOUS SEISMIC PROFILES

LEGEND:

- MCS data
- - - USNS KANE data
- WHOI data
- - - Lamont data



UNCLASSIFIED

UNCLASSIFIED





FIGURE I-19B: PARTIAL INVENTORY OF CONTINUOUS SEISMIC PROFILES

LEGEND: see Figure I-19A

UNCLASSIFIED



UNCLASSIFIED

UNCLASSIFIED

PRECEDING PAGE BLANK, NOT FILMED

LRAPP PRIORITY AREA TWO

UNCLASSIFIED

UNCLASSIFIED




FIGURE II-1: LOCATION OF LRAPP ATLANTIC AREA II SUBAREAS

CONFIDENTIAL





FIGURE II-2A

FIGURE II-2A: NUMBER OF WINTER (Jan-Mar) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE PER TIME DEPTH


CONFIDENTIAL

CONFIDENTIAL

FIGURE II-2B: NUMBER OF WINTER
(Jan-Mar) OCEANOGRAPHIC OBSERVATIONS
PER ONE-DEGREE SQUARE DEEPER THAN
DEEP AXIAL DEPTH



CONFIDENTIAL



CONFIDENTIAL

CONFIDENTIAL





FIGURE II-3A: NUMBER OF SPRING (Apr-Jun) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH


CONFIDENTIAL

CONFIDENTIAL

FIGURE II-3B: NUMBER OF SPRING
(Apr-Jun) OCEANOGRAPHIC OBSERVATIONS
PER ONE-DEGREE SQUARE DEEPER THAN
DEEP AXIAL DEPTH



CONFIDENTIAL



CONFIDENTIAL

CONFIDENTIAL





FIGURE II-1A

FIGURE II-1A: NUMBER OF SUMMER (Jul-Sep) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH

CONFIDENTIAL

FIGURE II-4B: NUMBER OF SUMMER
(Jul-Sep) OCEANOGRAPHIC OBSERVATIONS
PER ONE-DEGREE SQUARE DEEPER THAN
DEEP AXIAL DEPTH



CONFIDENTIAL




FIGURE II-4C: NUMBER OF SUMMER (Jul-Sep) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH

CONFIDENTIAL

CONFIDENTIAL





FIGURE II-5A

FIGURE II-5A: NUMBER OF AUTUMN (Oct-Dec) OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE SQUARE DEEPER THAN DEEP AXIAL DEPTH


CONFIDENTIAL

CONFIDENTIAL

FIGURE II-5B: NUMBER OF AUTUMN
(Oct-Dec) OCEANOGRAPHIC OBSERVATIONS
PER ONE-DEGREE SQUARE DEEPER THAN
DEEP AXIAL DEPTH



CONFIDENTIAL



UNCLASSIFIED




FIGURE II-6: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR WINTER (Jan-Mar)

UNCLASSIFIED

UNCLASSIFIED




FIGURE II-7: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR SPRING (Apr-Jun)

UNCLASSIFIED




FIGURE II-8: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR SUMMER (Jul-Sep)

UNCLASSIFIED




FIGURE II-9: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR AUTUMN (Oct-Dec)

UNCLASSIFIED

NOTE:

Letters represent difference (in meters/second) between velocity at upper sound channel axis and at subsurface sound velocity maximum, where:

- A = greater than 10
- B = 8 to 10
- C = 6 to 8
- D = 4 to 6
- E = 2 to 4
- F = 0.5 to 2
- G = 0.1 to 0.5




FIGURE II-10: ANNUAL AVERAGE "STRENGTH" OF UPPER SOUND CHANNEL (if present) RELATIVE TO SUBSURFACE SOUND VELOCITY MAXIMUM




FIGURE II-11: ANNUAL AREAL EXTENT AND AVERAGE AXIAL DEPTH OF SUBSURFACE SOUND VELOCITY MAXIMUM

UNCLASSIFIED




FIGURE II-12: ANNUAL AVERAGE DEPTH OF DEEP SOUND CHANNEL AXIS
(in meters X 100)

UNCLASSIFIED



LEGEND: see Figure I-13

FIGURE II-13: BATHYMETRY SHOALER THAN AVERAGE CRITICAL DEPTH
FOR WINTER (Nov-Apr)
(Contour interval = 500 fathoms)

UNCLASSIFIED



LEGEND: see Figure II-13

FIGURE II-14: BATHYMETRY OF DEEPER THAN AVERAGE CRITICAL DEPTH
FOR SUMMER (May-Oct)
(Contour interval = 500 fathoms)

Best Available Copy

UNCLASSIFIED

50° N

40°

30°

15° N

45° W

30°

15° W

REGION 5
1969

REGION 6
1969

100 FATHOMS

USOC CHART No. 2

USOC CHART No. 3

LEGEND:

0309 N
1969

0308 N
1969

0406 N
1951

0306 N
1952

0505 N
1966

0405 N
1966

0504 N
1953

0404 N
1953

0304 N
1969

0503 N
1953

0403 N
1952

103

104

103

104

103

104

28

29

42

43

59

60

79

80

81

Boundary of NAVOCEANO North Atlantic Regional Chart

Laughton Chart (National Institute of Oceanography)
Chart number (1966 compilation date)

NAVOCEANO BC Chart
Chart number
Compilation date

USOC Charts (pre-1960 compilation)

37

UNCLASSIFIED

UNCLASSIFIED




FIGURE II-16a


FIGURE II-16a: NUMBER OF BOTTOM SEDIMENT SAMPLES PER ONE-DEGREE SQUARE


UNCLASSIFIED

UNCLASSIFIED


FIGURE II-15B: NUMBER OF BOTTOM
SEDIMENT SAMPLES PER ONE-DEGREE
SQUARE

LEGEND: see Figure II-15A





UNCLASSIFIED



LEGEND: see Figure I-17

FIGURE II-17: DISTRIBUTION OF SURFICIAL BOTTOM SEDIMENTS

UNCLASSIFIED

UNCLASSIFIED





FIGURE II-19a

FIGURE II-19a: NUMBER OF BOTTOM CORES PER ONE-DEGREE SQUARE

UNCLASSIFIED

UNCLASSIFIED

FIGURE II-18B: NUMBER OF BOTTOM
COMES PER ONE-DEGREE SQUARE



UNCLASSIFIED




FIGURE II-130: NUMBER OF
BOTTOM CORES PER ONE-DEGREE
SQUARE

UNCLASSIFIED

UNCLASSIFIED





FIGURE II-19A: PARTIAL INVENTORY OF CONTINUOUS SEISMIC PROFILES

UNCLASSIFIED

FIGURE II-19B: PARTIAL INVENTORY
OF CONTINUED CLIMATIC PROFILES

LEGEND: see Figure II-19A



UNCLASSIFIED




FIGURE II-19C: PARTIAL INVENTORY
OF CONTINUOUS SEISMIC PROFILES

UNCLASSIFIED

PRECEDING PAGE, BLANK, NOT FILMED

UNCLASSIFIED

LRAPP PRIORITY AREA THREE

UNCLASSIFIED

UNCLASSIFIED

PRECEDING PAGE, BLANK, NOT FILMED




FIGURE III-1: LOCATION OF IRAPP ATLANTIC AREA III SUBAREAS

UNCLASSIFIED

CONFIDENTIAL

65° W

60°

55°

50° W

40° N

	2			1										
1	1	2			2									
3	1	2												
6	8	2	3	1										
7	4	2	1	2										1
6	2								1	1		1	1	
9	2	1				1	2	1		1	1			
70	1	2		1										
4	1	1					1		1					
5				2			1							
2						1	1	1						
5						1	1	1				1		
3	1						2	1						
2							1	1						
2	1						2				3			
	1					1	1				1			
		1				1	1					1		
2	1			1			1				1			
3	2	2	1	1			1							
3	2	5	2			1	1	1			2			

35°

30°

25°

20°

65° W

60°

55°

50° W

15° N

FIGURE III-2A: NUMBER OF
WINTER (Jan-Mar) OCEANOGRAPHIC
OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP
AXIAL DEPTH

CONFIDENTIAL




FIGURE III-2B: NUMBER OF WINTER (Jan-Mar)
OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN 1000 AXIAL DEPTH




FIGURE III-3A: NUMBER OF
SIGHTING (Apr-Jun) OCEANOGRAPHIC
OBSERVATION PER ONE-DEGREE
SQUARE DEEPER THAN THE
AXIAL DEPTH

CONFIDENTIAL




FIGURE III-3B: NUMBER OF SPRING (Apr-Jun)
OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE,
SQUARE DEEPER THAN TEEP AXIAL DEPTH

CONFIDENTIAL




FIGURE III-4A: NUMBER OF
SUMMER (Jul-Sep) OCEANOGRAPHIC
OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP
AXIAL DEPTH

CONFIDENTIAL





FIGURE III-4B: NUMBER OF SUMMER (Jul-Sep)
OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN DEEP AXIAL DEPTH



CONFIDENTIAL




FIGURE III-5B: NUMBER OF AUTUMN (Oct-Dec)
OCEANOGRAPHIC OBSERVATIONS PER ONE-DEGREE
SQUARE DEEPER THAN 1000 AXIAL DEPTH

CONFIDENTIAL

UNCLASSIFIED





FIGURE III-6: ARFAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR WINTER (Jan-Mar)

UNCLASSIFIED



LEGEND:

FIGURE III-7: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR SPRING (Apr-Jun)




FIGURE III-8: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR SUMMER (Jul-Sep)

UNCLASSIFIED





FIGURE III-9: AREAL EXTENT AND AVERAGE AXIAL DEPTH OF UPPER SOUND CHANNEL FOR AUTUMN (Oct-Dec)




FIGURE III-10: ANNUAL AVERAGE "STRENGTH" OF UPPER SOUND CHANNEL (if present) RELATIVE TO SUBSURFACE SOUND VELOCITY MAXIMUM



LEGEND:

FIGURE III-11: ANNUAL AREAL EXTENT AND AVERAGE AXIAL DEPTH OF SUBSURFACE SOUND VELOCITY MAXIMUM

UNCLASSIFIED




FIGURE III-12: ANNUAL AVERAGE DEPTH OF DEEP SOUND CHANNEL AXIS
(in meters X 100)

UNCLASSIFIED




FIGURE III-13: BATHYMETRY SHOALER THAN AVERAGE CRITICAL DEPTH
FOR WINTER (Nov-Apr)
(Contour interval = 500 fathoms)





FIGURE III-11: BATHYMETRY SHOALER THAN AVERAGE CRITICAL DEPTH
FOR SUMMER (May-Nov)
(Contour interval = 500 fathoms)

LEGEND:

 Boundary of NAVOCEANO North Atlantic Regional Chart



 NAVOCEANO BC Chart
 Chart number
 Compilation date


FIGURE III-15: INDEX OF BEST EXISTING BATHYMETRIC CHARTS

UNCLASSIFIED



UNCLASSIFIED

UNCLASSIFIED






FIGURE III-17: DISTRIBUTION OF SURFICIAL BOTTOM SEDIMENTS

UNCLASSIFIED




FIGURE III-18A: NUMBER OF
BOTTOM CORES PER ONE-DEGREE
SQUARE

UNCLASSIFIED





FIGURE III-18B: NUMBER OF BOTTOM CORES
PER ONE-DEGREE SQUARE

UNCLASSIFIED

UNCLASSIFIED



UNCLASSIFIED




FIGURE III-19B: PARTIAL INVENTORY OF
CONTINUOUS SEISMIC PROFILES

UNCLASSIFIED

REFERENCES (U)

- Beckerle, J. C., Oct 1968, A report on ATLANTIS II Cruise 22 - June, July, and August 1966 - Part II: sound velocity profiles: Woods Hole Oceanographic Inst. Reference no. 60-67 (unpublished manuscript).
- Bunce, J. A., Aug 1969, A 15-year summary of project CAESAR and related oceanographic surveys, 1954 to 1968 (U): U. S. Naval Oceanographic Office Informal Rept. no. 69-58. SECRET - NO FORN - LIMDIS
- , Apr 1970, Oceanographic cruise summary. Baffin Bay - Davis Strait - Labrador Sea, October 1969: U. S. Naval Oceanographic Office no. 70-16.
- Canadian Oceanographic Data Centre, 1969a, Labrador and Irminger Seas - March 12 to May 12, 1966: 1969 Data Record Series, Rept. no. 1, CODC Reference 10-66-001.
- , 1969b, Grand Banks to the Azores and Scotian Shelf - April 12 to May 5, 1966: 1969 Data Record Series, Rept. no. 2, CODC Reference 10-66-002.
- , 1969c, East Greenland, Denmark Strait, and Irminger Sea - January 16 to April 5, 1967: 1969 Data Record Series, Rept. no. 4, CODC Reference no. 10-67-001.
- Codispoti, L. A., and Kravitz, J. H., May 1968, Oceanographic cruise summary, Baffin Bay - Davis Strait - Labrador Sea. Summer 1967: U. S. Naval Oceanographic Office Informal Rept. no. 68-23.
- Dunkle, W. M., Jun 1969, R/V CHAIN cruise No. 82 - North Atlantic Ocean and Mediterranean Sea - Summary cruise report prepared by chief scientists, track charts, bathymetry, magnetics, and location of observations - May 10 to September 15, 1968: Woods Hole Oceanographic Inst. Reference no. 69-28 (unpublished manuscript).
- Dunkle, W. M., and Hayes, H. C., May 1968, Index charts to cruises conducted by members of the Geophysics Department - North Atlantic, South Atlantic, and Indian Oceans: Mediterranean and Red Seas - 1956-1965: Woods Hole Oceanographic Inst. Reference no. 66-19 (unpublished manuscript).

UNCLASSIFIED

REFERENCES (CONTINUED)

- Environmental Science Services Administration, Dec 1969, BOMEX preliminary data, salinity-temperature-depth observations: v. III, Parts 1 and 2 (unvalidated teletype messages).
- Ewing, J., Luskin, B., Roberts, A., and Hirshman, J., 1960, Subbottom reflection measurements on the continental shelf, Bermuda Banks, West Indies Arch, and West Atlantic Basins: Jour. Geophys. Research, v. 65, n. 9, p. 2849-2859.
- Ewing, M., Ewing, J. I., and Talwani, M., 1964, Sediment distribution in the oceans: Mid-Atlantic Ridge: Geol. Soc. America Bull., v. 75, p. 17-36.
- Fenner, D. F. and Bucca, P. J., Dec 1969, The upper and deep sound channel in the northeast Atlantic: U. S. Naval Oceanographic Office Informal Rept. no. 69-94.
- International Commission for the Northwest Atlantic Fisheries, 1968, Environmental surveys - NORWESTLANT 1-3, 1963: Part III. oceanographic data record, volumes I, II, and III: Special Pub. no. 7, prepared by the Canadian Oceanographic Data Centre.
- Jones, E. J. W. and Ewing, J. I., 1969, Age of the Bay of Biscay: evidence from seismic profiles and bottom samples: Science, v. 166, n. 3901, p. 102-105.
- Katz, E. J., Jun 1969, Variableness of the sound velocity profile about the Mid-Atlantic Ridge axis, Woods Hole Oceanographic Inst. Reference no. 69-50 (unpublished manuscript).
- Knott, S. T., Carter, J. C., and Brown, J. A., Jul 1968, R/V CHAIN cruise no. 70 (July 12, 1967 to August 2, 1967) and R/V CHAIN cruise no. 73 (September 6, 1967 to September 26, 1967) - summary cruise report prepared by chief scientists, track charts, bathymetry, magnetics, and location of observations - North Atlantic Ocean from Laurentian Channel to Cape Hatteras: Woods Hole Oceanographic Inst. Reference no. 68-50 (unpublished manuscript).
- LaCasce, E. O., Jr., Wooding, C. M., and Beckerle, J. C., Jun 1969, Sound velocity characteristics in the Sargasso Sea based on observations on cruise 89 of the R/V CHAIN: Woods Hole Oceanographic Inst. Reference no. 69-38 (unpublished manuscript).

UNCLASSIFIED

REFERENCES (CONTINUED)

Lee, A. and Ellett, D., 1967, On the water masses of the northwest Atlantic Ocean: Deep-Sea Research, v. 14, p. 183-190.

Lowrie, A. and Escowitz, E., editors, 1969, KANE 9: Global Ocean Floor Analysis and Research Data Series, v. 1, U. S. Naval Oceanographic Office Publication, U. S. Govt. Printing Office, Washington, D. C.

NAVOCEANO. This abbreviated reference, followed by the date, is used for the sake of brevity in the text to identify the following U. S. Naval Oceanographic Office publications:

U. S. Naval Oceanographic Office, 1965, Oceanographic atlas of the North Atlantic Ocean, Section V, Marine Geology: H. O. Pub no. 700.

———, Sep 1966b, Marine Geophysical Survey Program 65-67 western North Atlantic and eastern and central North Pacific oceans - area I, volume 8A, core, sound velocimeter, and bottom photographic stations - hydrographic stations including area II: U. S. Naval Oceanographic Office, SP-96-I-8A, prepared by Alpine Geophysical Associates, Inc. under contract N62306-1688.

———, Sep 1966b, Marine Geophysical Survey Program 65-67 western North Atlantic and eastern and central North Pacific oceans - area I, vol. 5, bathymetry and subbottom profiling U. S. Naval Oceanographic Office SP-96-I-5, prepared by Alpine Geophysical Associates, Inc. under contract N62306-1688.

———, Dec 1966, Marine Geophysical Survey Program 65-67 western North Atlantic and eastern central North Pacific oceans - area SF, volume 8A, core, sound velocimeter, and bottom photographic stations - sound velocity profiles: U. S. Naval Oceanographic Office, SP-96-SF-8A, prepared by Alpine Geophysical Associates, Inc. under contract N62306-1688.

———, Jan 1967, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 5 - volume 5, geology and geophysics: U. S. Naval Oceanographic Office, SP-95-5-5, prepared by Texas Instruments, Inc. under contract N62306-1687.

REFERENCES (CONTINUED)

- _____, Feb 1967a, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean, Norwegian Sea and Mediterranean Sea area 5 - volume 6, oceanographic stations and velocity profiles: U. S. Naval Oceanographic Office SP-95-7/4-6, prepared by Texas Instruments, Inc. under contract N62306-1687.
- _____, Feb 1967b, Marine Geophysical Survey Program 65-67 western North Atlantic and eastern and central North Pacific oceans - area 1, volume 8C, core, sound velocimeter, and bottom photographic stations - sound velocity profiles: U. S. Naval Oceanographic Office, SP-96-1-8C, prepared by Alpine Geophysical Associates, Inc. under contract N62306-1688.
- _____, Feb 1967c, Marine Geophysical Survey Program 65-67 western North Atlantic and eastern and central North Pacific oceans - area SF, vol. 5, bathymetry and subbottom profiling: U. S. Naval Oceanographic Office, SP-96-SF-5, prepared by Alpine Geophysical Associates, Inc. under contract N62306-1688.
- _____, Jun 1967, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 5 - volume 6S, oceanographic stations and velocity profiles (supplement): U. S. Naval Oceanographic Office, SP-95-5-6S, prepared by Texas Instruments, Inc. under contract N62306-1687.
- _____, Feb 1968, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 2 - volume 6, oceanographic stations and sound velocity profiles: U. S. Naval Oceanographic Office, SP-95-2-6, prepared by Texas Instruments, Inc. under contract N62306-1687.
- _____, Mar 1968, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 2 - volume 5, geology and geophysics: U. S. Naval Oceanographic Office, SP-95-2-5, prepared by Texas Instruments, Inc. under contract N62306-1688.
- _____, May 1968, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 3 - volume 6, oceanographic stations and

UNCLASSIFIED

REFERENCES (CONTINUED)

- sound velocity profiles: U. S. Naval Oceanographic Office, SP-95-3-6, prepared by Texas Instruments Inc. under contract N62306-1687.
- , Aug 1968, Marine Geophysical Survey Program 65-67 western North Atlantic Ocean and South China Sea area ST - volume 2, summary of geophysical information: U. S. Naval Oceanographic Office, SP-97-ST-2, prepared by Alpine Geophysical Associates, inc. under contract N62306-2020.
- , Oct 1968, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 3 - volume 5, geology and geophysics: U. S. Naval Oceanographic Office SP-95-3-5, prepared by Texas Instruments, Inc. under contract N62306-1687.
- , Mar 1969, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 7/4 - volume 6, oceanographic stations and sound velocity profiles: U. S. Naval Oceanographic Office, SP-95-7/4-6, prepared by Texas Instruments, Inc. under contract N62306-1687.
- , Apr 1969, United States Naval Oceanographic Office Marine Geophysical Survey Program 1965-1967 North Atlantic Ocean Norwegian Sea and Mediterranean Sea area 7/4 - volume 5, geology and geophysics: U. S. Naval Oceanographic Office, SP-95-7/4-5, prepared by Texas Instruments, Inc. under contract N62306-1687.
- Piip, A. T., Jan 1966, Precision sound velocity profiles in the ocean, volume I, the sound channel in the Bermuda-Barbados region (November-December 1963): Lamont Geological Observatory Tech. Rept. no. 3, prepared for NRL under contract Nonr 266(65).
- , Jun 1967, Precision sound velocity profiles in the ocean, volume II, the sound channel in the Bermuda-Barbados region (April-Jul 1964): Lamont Geological Observatory Tech. Rept. no. 4, prepared for NRL under contract Nonr 266(65).
- , Apr 1968, Precision sound velocity profiles in the ocean, volume IV, Canary Islands - Gibraltar - Bay of Biscay (June-July 1965): Lamont Geological Observatory Tech. Rept. no. 6, prepared for NRL under contract Nonr 266(65).

UNCLASSIFIED

REFERENCES (CONTINUED)

Wilson, W. D., 1960, Equation for the speed of sound in sea water: Jour. Acoustical Soc. America, v. 32, n. 10, p. 1357.

Wilson, J. T., 1963, The Mid-Labrador Sea Ridge: Nature, v. 200, no. 4911, p. 1085-1086.

Wolford, T. C., Jun 1966, Oceanography of the Grand Banks region and the Labrador Sea in 1966: U. S. Coast Guard Oceanographic Rept. no. 13 (CG 373-13).

Worthington, L. V., 1959, The 18° water in the Sargasso Sea: Deep-Sea Research, v. 5, p. 297-305.

———, 1970, The Norwegian Sea as a Mediterranean basin: Deep-Sea Research, v. 17, p. 77-84.

Wust, G. and Defant, A., 1936, Atlas zur schichtung und zirkulation des Atlantischen Ozeans, Walter de Gruyter and Co., Berlin.

UNCLASSIFIED

CONFIDENTIAL

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. NAVAL OCEANOGRAPHIC OFFICE ✓		2a. REPORT SECURITY CLASSIFICATION CONFIDENTIAL	
3. REPORT TITLE SOUND VELOCITY AND BOTTOM CHARACTERISTICS FOR LRAPP ATLANTIC AREAS I, II, AND III (U).		2b. GROUP 4	
4. DESCRIPTIVE NOTES (Type of report and effective dates) Informal Report.			
5. AUTHOR(S) (First name, middle initial, last name) Don F. Fenner, Paul J. Bucca, and Charles L. Davis			
6. REPORT DATE June 1971		7a. TOTAL NO. OF PAGES 122	7b. NO. OF REFS 46
8a. CONTRACT OR GRANT NO. b. PROJECT NO. 232 c. 714-EL-GJK d. NSO-IR-71-2		9a. ORIGINATOR'S REPORT NUMBER(S) IR NO. 71-2 ✓	
9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)			
10. DISTRIBUTION STATEMENT ~~This document may be further distributed by any holder only with specific prior approval of the Naval Oceanographic Office.~~			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U. S. Naval Oceanographic Office Washington, D. C. 20390	
13. ABSTRACT (U) Inventories and summaries of sound velocity, bathymetry, and bottom characteristics have been compiled for three areas in the North Atlantic Ocean between 15° and 65° N latitude and east of 50° to 65° W longitude. The following information is contained in this report: seasonal inventories of sound velocity profiles extending deeper than deep axial depth, charts of the seasonal extent and average axial depth of the upper sound channel, charts of the annual "strength" of the upper sound channel, charts of the annual extent and average depth of the subsurface sound velocity maximum, annual contour charts of deep axial depth, charts showing bathymetry shoaler than critical depth for summer and winter, an index of the best available bathymetric contour charts, inventories of surficial bottom sediment samples, charts of surficial bottom sediment analysis by grain size classes, an inventory of bottom cores, and a partial inventory of continuous seismic profiles. In addition, a brief analysis is given of water masses that effect sound velocity structures.			

DD FORM 1473

NOV 65

(PAGE 1)

S/N 0101-807-6801

(This page is unclassified)

UNCLASSIFIED

Security Classification

4B

Security Classification ~~TOP SECRET~~

CONFIDENTIAL

14

KEY WORDS

LINK A

LINK B

LINK C

SOLE

NY

	NAME	ROLE
1	Mr. J. Edgar Hoover	Director
2	Mr. Clegg	Chief Clerk
3	Mr. Glavin	Assistant Director
4	Mr. Ladd	Assistant Director
5	Mr. Nichols	Assistant Director
6	Mr. Rosen	Assistant Director
7	Mr. Tracy	Assistant Director
8	Mr. Egan	Assistant Director
9	Mr. Gurnea	Assistant Director
10	Mr. Harbo	Assistant Director
11	Mr. Hendon	Assistant Director
12	Mr. Pennington	Assistant Director
13	Mr. Quinn	Assistant Director
14	Mr. Nease	Assistant Director
15	Mr. Tamm	Assistant Director
16	Mr. Winterrowd	Assistant Director
17	Mr. Mohr	Assistant Director
18	Mr. Tele. Rm.	Telephone Room
19	Mr. Holmes	Mr. Tolson's Secretary
20	Miss Gandy	Miss Gandy

WT

[illegible]

NY T

Continuous seismic profile inventory

CONFIDENTIAL

DD FORM 1473 (BACK)
(PAGE 2) (This page is unclassified)

UNCLASSIFIED

Security Classification



DEPARTMENT OF THE NAVY

OFFICE OF NAVAL RESEARCH
875 NORTH RANDOLPH STREET
SUITE 1425
ARLINGTON VA 22203-1995

IN REPLY REFER TO:

5510/1
Ser 321OA/011/06
31 Jan 06

MEMORANDUM FOR DISTRIBUTION LIST

Subj: DECLASSIFICATION OF LONG RANGE ACOUSTIC PROPAGATION PROJECT (LRAPP) DOCUMENTS

Ref: (a) SECNAVINST 5510.36

Encl: (1) List of DECLASSIFIED LRAPP Documents

1. In accordance with reference (a), a declassification review has been conducted on a number of classified LRAPP documents.
2. The LRAPP documents listed in enclosure (1) have been downgraded to UNCLASSIFIED and have been approved for public release. These documents should be remarked as follows:

Classification changed to UNCLASSIFIED by authority of the Chief of Naval Operations (N772) letter N772A/6U875630, 20 January 2006.

DISTRIBUTION STATEMENT A: Approved for Public Release; Distribution is unlimited.

3. Questions may be directed to the undersigned on (703) 696-4619, DSN 426-4619.

A handwritten signature in black ink, appearing to read "B. F. Link", is positioned above the typed name.

BRIAN LINK
By direction

Subj: DECLASSIFICATION OF LONG RANGE ACOUSTIC PROPAGATION PROJECT
(LRAPP) DOCUMENTS

DISTRIBUTION LIST:

NAVOCEANO (Code N121LC – Jaime Ratliff)
NRL Washington (Code 5596.3 – Mary Templeman)
PEO LMW Det San Diego (PMS 181)
DTIC-OCQ (Larry Downing)
ARL, U of Texas
Blue Sea Corporation (Dr. Roy Gaul)
ONR 32B (CAPT Paul Stewart)
ONR 321OA (Dr. Ellen Livingston)
APL, U of Washington
APL, Johns Hopkins University
ARL, Penn State University
MPL of Scripps Institution of Oceanography
WHOI
NAVSEA
NAVAIR
NUWC
SAIC

Declassified LRAPP Documents

Report Number	Personal Author	Title	Publication Source (Originator)	Pub. Date	Current Availability	Class.
IR 71-2	Fenner, D. F., et al.	SOUND VELOCITY AND BOTTOM CHARACTERISTICS FOR LRAPP ATLANTIC AREAS I, II, AND III (U)	Naval Oceanographic Office	710601	ADC008372; ND	U
T-71-NJ-4508-C	Larsen, H. L., et al.	LRAPP DATA COLLECTION (U)	Tracor, Inc.	710831	AD0517012; ND	U
Unavailable	Anderson, C. G., et al.	ADAPTIVE BEAMFORMING ANALYSIS FOR DIRECTIONALITY USING DATA FROM A VERTICAL ARRAY IN THE MEDITERRANEAN	Naval Undersea Research and Development Center	710901	AD0517696	U
MC PLAN 06	Unavailable	IOMEDEX LRAPP OPERATION ORDER (U)	Maury Center for Ocean Science	710924	ND	U
NRLFR7322	Lawson, W. M.	POSITION-DETERMINING SYSTEM FOR SEA-SPIDER HYDROPHONE ARRAYS	Naval Research Laboratory	711230	ND	U
N00014-71-C-0088	Unavailable	CONTINUATION OF LRAPP FINAL REPORT (U)	Bell Laboratories	720201	AD0520426; NS; ND	U
Unavailable	Unavailable	PARKA II-A EXPERIMENT. VOLUME 2	Maury Center for Ocean Science	720201	AD0596342	U
Unavailable	Unavailable	NOISE POWER MEASUREMENTS FROM THE LAMBDA TOWED ARRAY EXPERIMENT	Texas Instruments, Inc.	720401	ND	U
Unavailable	Anderson, P. R., et al.	LRAPP VERTICAL ARRAY PRELIMINARY DESIGN STUDY, PHASE I	Westinghouse Research Laboratories	720414	AD0900477	U
Unavailable	Unavailable	SUPPLEMENTARY DATA. NOISE POWER MEASUREMENTS FROM THE LAMBDA TOWED ARRAY EXPERIMENT	Texas Instruments, Inc.	720419	ND	U
MC Report 6, Volume 2	Unavailable	PARKA OCEANOGRAPHIC DATA COMPENDIUM-PARKA II-A	Maury Center for Ocean Science	720501	ND	U
MC Report 6, Volume 3	Unavailable	PARKA OCEANOGRAPHIC DATA COMPENDIUM-PARKA II-B	Maury Center for Ocean Science	720501	ND	U
Unavailable	Unavailable	NORLANT 72 PHASE 2 OPERATION PLAN	Naval Underwater Systems Center	720628	AD0521225	U
Unavailable	Unavailable	NORLANT 72 PHASE 2 SCIENTIFIC PLAN	Naval Underwater Systems Center	720628	AD0521226	U
Unavailable	Unavailable	NORLANT 72 PHASE 3 OPERATION PLAN	Naval Underwater Systems Center	720628	AD0521227	U
Unavailable	Unavailable	NORLANT 72 PHASE 3 SCIENTIFIC PLAN	Naval Underwater Systems Center	720628	AD0521228	U
MC PLAN 08	Unavailable	OPERATION PLAN. LRAPP TASSRAP EXERCISE (TEX)	Maury Center for Ocean Science	720701	ND	U
Unavailable	Miller, R. R., III	CURRENT REGIME OF THE MALTESE OCEANIC FRONTAL ZONE	Naval Underwater Systems Center	720906	AD0749706	U