

COMPONENT PART NOTICE

THIS PAPER IS A COMPONENT PART OF THE FOLLOWING COMPILATION REPORT:

TITLE: Proceedings of the Medical Defense Bioscience Review (1993) Held  
in Baltimore, Maryland on 10-13 May 1993. Volume 3.

TO ORDER THE COMPLETE COMPILATION REPORT, USE AD-A275 669.

THE COMPONENT PART IS PROVIDED HERE TO ALLOW USERS ACCESS TO INDIVIDUALLY AUTHORED SECTIONS OF PROCEEDING, ANNALS, SYMPOSIA, ETC. HOWEVER, THE COMPONENT SHOULD BE CONSIDERED WITHIN THE CONTEXT OF THE OVERALL COMPILATION REPORT AND NOT AS A STAND-ALONE TECHNICAL REPORT.

THE FOLLOWING COMPONENT PART NUMBERS COMPRISE THE COMPILATION REPORT:

AD#: AD-P008 853 AD#: \_\_\_\_\_  
 AD#: thru AD#: \_\_\_\_\_  
 AD#: AD-P008 900 AD#: \_\_\_\_\_

**DTIC**  
**ELECTE**  
**S F D**  
 MAR 18 1994

This document has been approved for public release and sale; its distribution is unlimited.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail
A-1	

Ultrastructural Studies and Transendothelial  
Resistance of Phosgene-exposed  
Endothelial Cells

J.S. Madren-Whalley and R.J. Werrlein

United States Army Medical Research Institute  
of Chemical Defense, Aberdeen Proving Ground,  
Maryland 21010-5425

AD-P008 874  


ABSTRACT

Sheep pulmonary artery endothelial cells (PAEC) were grown on Millicell CM™ inserts pre-treated with type IV collagen. Using these inserts we successfully produced contiguous sheets of cells that mimic *in vivo* organization of endothelial tissues. Ultrastructural details, obtained by transmission electron microscopy showed that the plasmalemma of adjacent cells were partially joined by tight junctions and overlapping end processes. They were also joined by compressed basal lamina which lined the dorsoventral surfaces of apposing cells. In control populations, the basal lamina were uniform in dimension (0.1 to 0.35  $\mu\text{m}$ ) along their entire length and were filled with compressed extracellular matrix materials. Transendothelial resistance of these contiguous endothelial populations was low and averaged only 6.77  $\pm$  0.14 ohms per  $\text{cm}^2$ . Exposure to phosgene (137 ppm x 20 min) tore the basal lamina apart and produced large paracellular leakage paths. Doses as high as 561 ppm x 20 min. had no obvious effects on other ultrastructural details and did not decrease cell viability but did produce a 23% decrease in transendothelial resistance. We conclude that phosgene's effects on the basal lamina of PAEC may be linked by f-actin lesions to increased permeability and to cytoskeletal-membrane interactions that control normal barrier function.

94-08290



## INTRODUCTION

It has been known since the beginning of the century that the primary cause of death from phosgene inhalation is lung edema (Roos, 1914). The consequences of that fact were realized in the trenches of World War I where phosgene had devastating effects on military personnel and gave rise to extensive case studies on battlefield casualties and on the pathogenesis of phosgene toxicity. By 1978, phosgene was being commercially manufactured in million-ton volumes and concern about human exposure had been revitalized. However, despite occupational and military risks from exposure to phosgene, the mechanisms that disrupt barrier function and control permeability of pulmonary tissues remain poorly understood. Werrlein et al. (1991) have demonstrated that phosgene, in doses equivalent to 0.5, 1.0, 2.0 and 5.0 x LC<sub>50</sub> for sheep (3300 ppm · min), disrupts barrier function of sheep pulmonary artery endothelial cells (PAEC) grown in culture on cytodex-3 microcarrier beads. Dose-response effects from those studies indicated that the PAEC lesions produced by phosgene were peracellular. To confirm or refute that concept, we have investigated the effects of phosgene on transendothelial resistance and ultrastructure using the following *in vitro* model.

## MATERIALS AND METHODS

### *Sheep Pulmonary Artery Endothelial Cells (PAEC) Cultures*

PAEC from proliferating, low passage (4-7), stock cultures were seeded at 20,000 cells/cm<sup>2</sup> onto 12 mm Millicell CM™, PTFE inserts (Millipore Products, Bedford, MA). Inserts (Fig. 1A) were pre-coated with type IV collagen (Sigma Chemical Co., St. Louis, MO). Attached, PAEC cultures were subject to medium renewal 3 times per week (M-W-F) and were maintained in a Hotpack incubator (Hotpack Corp., Phila., PA) at 35 °C in an atmosphere of 5% CO<sub>2</sub> and air.




Fig. 1 Millicell CM™ insert (A) showing PTFE Biopore™ membrane, extracellular matrix of type IV collagen, layer of attached PAECs and surround of nutrient medium (apical and basal). Resistance measurement chamber (B) showing EVOM™ connectors, position of the Millicell CM™ insert relative to disc electrodes which face each other from opposite sides of the PAEC tissue sheet. Clearance between the top electrode and PTFE membrane was set at 1-2 mm.

### *Nutrient Medium*

PAEC were grown in HAM's F-12 medium containing 15% fetal bovine serum, 5% Omni<sup>TM</sup> serum (Advanced Biotechnologies Inc., Columbia, MD), L-glutamine (2mM), gentamicin sulfate (50 µg/ml) and endothelial cell growth supplement (50 µg/ml). All chemicals were obtained from Sigma Chemical Co., St. Louis, MO., unless otherwise stated.

### *Transendothelial Resistance Measurements*

Transendothelial tissue resistance was measured with an EVOM<sup>TM</sup> voltohm meter and an Endohm 12<sup>TM</sup> resistance measurement chamber (Fig. 1B), World Precision Instruments, Inc., New Haven, CT. Current passed between apposing faces of disc electrodes, placed above and below the attached sheet of endothelial tissues, was used to determine resistance to the current flow. The background resistance of a blank (collagen coated insert without cells) was subtracted from culture readings and the difference used to determine resistance of the cell sheet. Normalized values were expressed as ohms ( $\Omega$ ) · cm<sup>2</sup>.

### *Phosgene Exposure*

To expose Millicell-CM<sup>TM</sup> insert cultures, the overlaying, apical layer of medium was first reduced to a volume of 25 µl. That volume was sufficient to keep the PAEC wet and assure interaction with phosgene during the 20 min. dosing period. Immediately prior to exposure, replicate insert-cultures were placed in open, 60 mm Petri dishes containing 1.5 ml of freshly prepared medium. Replicates were then transferred to an environmentally controlled, 37 °C chamber and were exposed to a dose (137 ppm x 20 min) of phosgene in a humidified atmosphere of 5% CO<sub>2</sub> and air. The dose was verified by removing 10 µl gas samples from the exposure chamber at the beginning and end of each exposure and analyzing the sample by gas chromatography. Sham-treated control cultures were exposed to 5% CO<sub>2</sub> in air. At the end of an exposure period, replicate cultures were immediately prepared for transendothelial resistance and ultrastructural studies.

### *Ultrastructural Studies*

Sham-treated controls and phosgene-exposed populations were rinsed with 0.1 M sodium cacodylate buffer then fixed with 4% glutaraldehyde in 0.1 M sodium cacodylate buffer. The cultures were washed again in 0.1 M sodium cacodylate buffer and postfixated with 1% osmium tetroxide. After dehydration by passage through increasing alcohol gradients to propylene oxide, the cells were embedded in POLY/BED 812 (Polysciences Inc., Warrington, PA). Thin sections were cut on an LKB Ultramicrotome IV and stained with uranyl acetate and lead citrate.

## RESULTS

### *Phosgene's Effects on Ultrastructure*

Ultrastructural details show the major difference between sham-treated controls and phosgene-exposed populations was in tissue continuity. Untreated controls (Fig. 2A) had no obvious gaps between plasma membranes or apposing faces of adjacent cells. These populations were closely knit and cells that overlapped were held together by a thin scaffolding of compressed basal lamina. The basal lamina were uniform in dimension and defined an intercellular space of 0.1 to 0.35  $\mu\text{m}$  along their entire length. That space was filled with a ground substance of extracellular



Fig. 2 Ultrastructure of control (A) and phosgene exposed (B) PAECs. The plasmalemma of control cells were held together by a thin scaffolding of compressed basal lamina (BL). BL were typically 0.1 to 0.35  $\mu\text{m}$  wide along their entire length and were filled with a ground substance of extracellular matrix materials. When exposed to phosgene (137 ppm x 20 min), the basal lamina were rended apart and paracellular leakage paths (PLP) developed between the apposing faces of plasma membranes and cells that overlapped. Membranes joined by tight junctions (M) were not disrupted by phosgene.

matrix materials. In contrast, the basal lamina of sheep PAECs exposed to 2740 ppm min phosgene (Fig. 2B) were rended apart in a manner that separated the faces of apposed plasma membranes and produced large paracellular leakage paths. Cells from the exposed populations were slightly misshapen by the effects of phosgene but their organelles, e.g., nuclei, mitochondria, lysosomes and endoplasmic reticulum remained intact. Unions in

which plasma membranes were joined by tight junctions were resistant to separation by phosgene. Such unions frequently involved plasma membranes from adjacent cells that overlapped in thin folds at their dorsal and ventral edges.

#### *Transendothelial Resistance*

Populations of PAECs became confluent on Millicell CM™ inserts within 1 week of seeding. However, we found these cells to be poorly suited for resistance studies. Even when confluent for 3-4 weeks, the average transendothelial resistance was only  $6.77 \pm 0.14$  ohms  $\cdot$  cm<sup>2</sup>. All studies, therefore, required populations that were confluent for at least 4 weeks. High doses of phosgene did have a limited impact on transendothelial resistance but how these changes relate to barrier function is uncertain. Populations exposed to 11,220 ppm  $\cdot$  min (561 ppm  $\times$  20 min) showed a progressive 10% to 23% decrease in transendothelial resistance relative to sham-treated controls (Fig. 3) during the ensuing 24 hour period. Results from smaller doses (2740 ppm  $\cdot$  min) indicate that phosgene induced decreases in tissue resistance did persist for up to 72 hours (results not shown).




Fig. 3 Transendothelial resistance measurements of sheep PAECs in Millicell CM™ insert cultures. Sham-treated control populations (white bars) were exposed for 20 min to 5% CO<sub>2</sub> and air. Populations exposed to phosgene (561 ppm  $\times$  20 min) are shown as black bars. Average readings (ohms/cm<sup>2</sup>) and standard error bars compare replicate control (n = 9) and exposed (n = 17) populations at T = 0 hours (immediately after 20 min. exposure) and T = 24 hours post exposure. Results were subject to one way ANOVA \*\* = p < .01.

## DISCUSSION

Our ultrastructural studies indicate that doses of 137 ppm x 20 min disrupted basal lamina and produced paracellular leakage paths in Millicell CM™ cultures of sheep PAECs. These leakage paths are consistent with results from earlier studies (Werrlein et al., 1991) which showed that acute, 20 minute exposures to phosgene increased permeability of PAEC tissues immediately and in a dose-dependent fashion. Permeability was quantified in the 1991 study by transendothelial adsorption of erythrosin-B (a red viability stain) as it moved across the PAEC tissue barrier and adhered to the cell substrate, i.e., microcarrier beads. These direct and immediate effects of phosgene do not explain delayed onset of acute pulmonary edema, i.e., the "clinical latent phase." They may, however, help to explain why phosgene exposure in rats causes immediate increase in lung lavage fluid protein (Ghio et al., 1991). It is also interesting that F-actin depletion and paracellular leakage paths appear to develop in concert. That observation is in agreement with evidence which indicates that cellular interaction with basal lamina is F-actin dependent (Svorboda and Hay, 1987).

Average transendothelial resistance of sheep PAECs was very low ( $6.77 \pm 0.14 \text{ ohms} \cdot \text{cm}^2$ ). Phosgene, in high doses (11,220 ppm min) did produce a progressive decrease in tissue resistance. However, to a lesser extent, so did manipulation of the sham-control populations. Significant decreases in resistance were observed in exposed populations at T= 24 hours. While our resistance values are in agreement with the literature where pulmonary transendothelial resistance ranges from  $6.1 \pm 0.5 \text{ } \Omega \cdot \text{cm}^2$  to  $15 \text{ } \Omega \cdot \text{cm}^2$  (Territo et al., 1984; Shasby and Shasby, 1986; Naveb et al., 1986), the low resistance of PAECs does not lend itself to sensitive and routine testing for toxicity or decreased barrier function. The model does, however, have considerable potential for *in vitro* study of lung cells and tissues, especially the effects of toxic gases on ultrastructure and barrier function.

## CONCLUSIONS

Phosgene disrupted basal lamina of PAECs (in culture) and produced large paracellular leakage paths that indicate a mechanism for decreased barrier function.

Average transendothelial resistance of sheep PAECs is low ( $6.77 \pm 0.14 \text{ } \Omega \cdot \text{cm}^2$ ). High doses of phosgene (11,220 ppm min) caused a small but progressive and significant decrease in tissue resistance during a 24 hour post exposure period.

Phosgene's effects on the basal lamina and F-actin cytoskeleton of PAECs may work in concert to increase permeability of the lung's endothelial tissues.

## REFERENCES

Ghio, A.J., Kennedy, T.P., Hatch, G.E., and Tepper, J.S. (1991). Reduction of neutrophil influx diminishes lung injury and mortality following phosgene inhalation. *J. Appl. Physiol.* 71(2): 657-665.

Naveb, M., Hough, G.P., Berliner, J.A., Frank, J.A., Fogelman, A.M., Haberland, M.E., and Edwards, P.O.A. (1986). Rabbit beta-migrating very low density lipoprotein increases endothelial macromolecular transport without altering electrical resistance. *J. Clin. Invest.* 78: 388-397.

Roos, A. (1914). Phosgene poisoning. *Vjschr. gerichtl. Med.* 48: 67-101.

Shasby, M.D. and Shasby, S.S. (1986). Effects of calcium on transendothelial albumin transfer and electrical resistance. *J. Appl. Physiol.* 60(1): 71-79.

Svoboda, K.H. and Hay, E.D. (1987). Embryonic corneal epithelial interaction with exogenous laminin and basal lamina in F-actin dependent. *Develop. Biol.* 123: 455-469.

Territo, M., Berliner, J.A., and Fogelman, A.M. (1984). Effects of monocyte migration on low density lipoprotein transport across aortic endothelial cell monolayers. *J. Clin. Invest.* 74: 2279-2284.

Werrlein, R.J., Madren-Whalley, J., and Kirby, S.D. (1991). Phosgene and perfluoroisobutylene: Their effects on permeability of endothelial tissues in culture. in *Proc. of 1991 Med. Def. Biosci. Rev.*, U.S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, MD, pp 325-328. AD B158588