

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP012813

TITLE: Exciton-Polariton Band Structure in Quantum-Dot Lattices

DISTRIBUTION: Approved for public release, distribution unlimited

Availability: Hard copy only.

This paper is part of the following report:

TITLE: Nanostructures: Physics and Technology International Symposium
[6th] held in St. Petersburg, Russia on June 22-26, 1998 Proceedings

To order the complete compilation report, use: ADA406591

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP012712 thru ADP012852

UNCLASSIFIED

Exciton-polariton band structure in quantum-dot lattices

E. L. Ivchenko[†], *Y. Fu*[‡] and *M. Willander*[‡]

[†] Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

[‡] Physical Electronics and Photonics, Department of Microelectronics and Nanoscience, University of Gothenberg and Chalmers University of Technology, Fysikgränd 3, S-412 96 Göteborg, Sweden

Abstract. We develop a theory of exciton polaritons in three-dimensional quantum-dot lattices with the period comparable to the light wavelength. A system of the Maxwell equations and nonlocal material relation are used to derive the dispersion equation in a rather general and well-converging form. A possibility of analytical description of the dispersion is questioned and discussed. The photon band structure is calculated for a face-centered-cubic lattice with spherical dots of the radius exceeding the bulk-exciton Bohr radius. The dispersion along the $\Gamma - X$ and $\Gamma - L$ lines is characterized by a strong anticrossing between bare transverse photon and exciton branches and by remarkable overlapping band gaps. Approaching the U and W points the exciton-polariton branches converge and the gap becomes negligible.

Recently van Coevorden et al. [1] have calculated the optical band structure of a three-dimensional (3D) lattice of resonant two-level atoms. They have solved numerically the dispersion equation for light waves in a face-centered-cubic atomic lattice and demonstrated that, in the certain range of parameters, there exists an overlap of photonic band gaps in all directions in the frequency region near the two-level resonance. Here we consider the photonic (or more precisely, exciton-polaritonic) band structure of lattices formed by a 3D periodic array of quantum dots (QDs).

We start from the Maxwell equations

$$\begin{aligned}\Delta \mathbf{E} - \text{grad div } \mathbf{E} &= - \left(\frac{\omega}{c} \right)^2 \mathbf{D} , \\ \text{div } \mathbf{D} &= 0\end{aligned}\tag{1}$$

for the electric field $\mathbf{E}$ and the displacement vector $\mathbf{D}$. The nonlocal material equation relating $\mathbf{D}$ and $\mathbf{E}$ is taken in the form (see [2])

$$\mathbf{D}(\mathbf{r}) = \varepsilon_b \mathbf{E}(\mathbf{r}) + 4\pi \mathbf{P}_{exc}(\mathbf{r}) ,\tag{2}$$

$$4\pi \mathbf{P}_{exc}(\mathbf{r}) = T(\omega) \sum_{\mathbf{a}} \Phi_{\mathbf{a}}(\mathbf{r}) \int \Phi_{\mathbf{a}}(\mathbf{r}') \mathbf{E}(\mathbf{r}') d\mathbf{r}' .\tag{3}$$

Here $\mathbf{a}$ are the lattice translation vectors enumerating quantum dots, $\Phi_{\mathbf{a}}(\mathbf{r}) = \Phi_0(\mathbf{r} - \mathbf{a})$ is the envelope function $\Psi_{exc}(\mathbf{r}_e, \mathbf{r}_h; \mathbf{a})$ of an exciton excited in the $\mathbf{a}$ th QD at coinciding electron and hole coordinates: $\Phi_{\mathbf{a}}(\mathbf{r}) = \Psi_{exc}(\mathbf{r}, \mathbf{r}; \mathbf{a})$. The other notations are

$$T(\omega) = 2\pi \frac{\varepsilon_b \omega_{LT} \omega_0 a_B^3}{\omega_0^2 - \omega^2} ,\tag{4}$$

ω_{LT} and a_B are the exciton longitudinal-transverse splitting and Bohr radius in the corresponding bulk semiconductor, ω_0 is the QD-exciton resonance frequency, ε_b is the background dielectric constant which is assumed to coincide with the dielectric constant of the barrier material. In the following we neglect the overlap of exciton envelope functions $\Psi_{\mathbf{a}}$ and $\Psi_{\mathbf{a}'}$ with $\mathbf{a} \neq \mathbf{a}'$ so that excitons excited in different dots are assumed to be coupled only via electromagnetic field.

It follows from Eq. (2) that $\text{div } \mathbf{E} = -(4\pi/\varepsilon_b) \text{div } \mathbf{P}_{exc}$ which allows to rewrite the first Eq. (1) as

$$\Delta \mathbf{E} + k^2 \mathbf{E} = -4\pi k_0^2 (1 + k^{-2} \text{grad div}) \mathbf{P}_{exc}, \quad (5)$$

where $k_0 = \omega/c$, $k = k_0 n_b = \omega n_b/c$ and $n_b = \sqrt{\varepsilon_b}$.

We seek for Bloch-like solutions of Eq. (5) satisfying the translational symmetry $\mathbf{E}_{\mathbf{K}}(\mathbf{r} + \mathbf{a}) = \exp(i\mathbf{K}\mathbf{a}) \mathbf{E}_{\mathbf{K}}(\mathbf{r})$ where the wave vector $\mathbf{K}$ is defined within the first Brillouin zone. The exciton-polariton dispersion $\omega(\mathbf{K})$ can be shown to satisfy the equation

$$\text{Det}||\delta_{\alpha\beta} - R_{\alpha\beta}(\omega, \mathbf{K})|| = 0, \quad (6)$$

where $\alpha, \beta = x, y, z$, $\delta_{\alpha\beta}$ is the Kronecker symbol and, for QD lattices,

$$R_{\alpha\beta} = \frac{k_0^2 T(\omega)}{v_0} \sum_{\mathbf{b}} \frac{|I_{\mathbf{K}+\mathbf{b}}|^2 S_{\alpha\beta}(\mathbf{K} + \mathbf{b})}{|\mathbf{K} + \mathbf{b}|^2 - k^2}, \quad (7)$$

$$I_{\mathbf{Q}} = \int \Phi_0(\mathbf{r}) e^{i\mathbf{Q}\mathbf{r}} d\mathbf{r}, \quad S_{\alpha\beta} = \delta_{\alpha\beta} - \frac{Q_{\alpha} Q_{\beta}}{k^2}, \quad (8)$$

$\mathbf{b}$ are the reciprocal lattice vectors and v_0 is the volume of the lattice primitive cell.

Similarly to [1] we consider a face-centered-cubic lattice with the lattice constant a and the unit-cell volume $v_0 = a^3/4$. It is convenient to introduce a dimensionless parameter $P = (\pi\sqrt{3}c/a\omega_0 n_b)^3$ and the dimensionless frequency $\Omega = \omega/\omega_0$. The calculation is performed for spherical QDs with the radius R exceeding the Bohr radius a_B in which case we have

$$I_{\mathbf{Q}} = \pi \left(\frac{2R}{a_B} \right)^{3/2} \frac{\sin QR}{QR[\pi^2 - (QR)^2]}. \quad (9)$$

Then Eq. (7) can be transformed into

$$R_{\alpha\beta}(\Omega, \mathbf{K}) = N \frac{\Omega^2}{\Omega^2 - 1} \sigma_{\alpha\beta}(\Omega, \mathbf{K}), \quad (10)$$


$$\sigma_{\alpha\beta}(\Omega, \mathbf{K}) = \sum_{\mathbf{b}} \frac{f(|\mathbf{K} + \mathbf{b}|R) S_{\alpha\beta}(\mathbf{K} + \mathbf{b})}{\Omega^2 - \Omega^2(\mathbf{K} + \mathbf{b})}, \quad (11)$$

$$N = \frac{64}{\pi} \frac{\omega_{LT}}{\omega_0} \left(\frac{R}{a} \right)^3, \quad f(x) = \left[\frac{\pi^2 \sin x}{x(\pi^2 - x^2)} \right]^2, \quad (12)$$

$\Omega(\mathbf{Q}) = cQ/\omega_0 n_b$. Eq. (6) is equivalent to the three separate equations $R_j(\Omega, \mathbf{K}) = 1$ where R_j ($j = 1, 2, 3$) are eigenvalues of the matrix $R_{\alpha\beta}$. For high-symmetry points of the Brillouin zone, the symmetry imposes certain relations between the $R_{\alpha\beta}$ components

Table 1. Dispersion equations written in terms of $R_{\alpha\beta}$ for different $\mathbf{K}$ points in the Brillouin zone.

$\mathbf{K}$ ($2\pi/a$)	Nonzero components of $R_{\alpha\beta}$	Dispersion equations
Γ (0, 0, 0)	$R_{xx}=R_{yy}=R_{zz}$	$R_{xx}=1$
X (0, 0, 1)	$R_{xx}=R_{yy}, R_{zz}$	$R_{xx}=1, R_{zz}=1$
L (1/2, 1/2, 1/2)	$R_{\alpha\alpha}=R_{xx}, R_{\alpha\beta}=R_{xy}(\alpha \neq \beta)$	$R_{xx}-R_{xy}=1, R_{xx}+2R_{xy}=1$
W (1/2, 0, 1)	$R_{xx}, R_{yy}=R_{zz}$	$R_{xx}=1, R_{yy}=1$
K (3/4, 0, 3/4)	$R_{xx}=R_{zz}, R_{yy}, R_{xz}=R_{zx}$	$R_{xx}\pm R_{xz}=1, R_{yy}=1$
U (1/4, 1/4, 1)	$R_{xx}=R_{yy}, R_{zz}, R_{xy}=R_{yx}$	$R_{xx}\pm R_{xy}=1, R_{zz}=1$

**Fig 1.** Exciton-polariton dispersion near the exciton resonance frequency ω_0 in a face-centered-cubic lattice of spherical QDs characterized by the following set of parameters: $P = 1.1$, $R/a = 1/4$ and $\omega_{LT}/\omega_0 = 5 \times 10^{-4}$. The dashed lines show the photon dispersion in the empty lattice, i.e. for $\omega_{LT} = 0$, the dotted horizontal line indicates the value $\omega = \omega_0$.

and the eigenvalues R_j can be readily expressed via these components as illustrated in Table 1 for the points Γ, X, L, W, K and U .

Further simplifications follow taking into account a small value of the parameter N in Eq. (10) since, in semiconductors, the ratio ω_{LT}/ω_0 typically lies between 10^{-4} and 10^{-3} . Then in the frequency region given by the condition $|\Omega - 1| \ll P^{1/3} - 1$ one can readily use the approximate equation $\Omega - 1 \approx (N/2)\sigma_j(1, \mathbf{K})$ where σ_j are eigenvalues of the $\sigma_{\alpha\beta}$ matrix.

Fig. 1 shows the photonic band structure calculated for the dots of radius $R = a/4$ and for $P = 1.1$, $\omega_{LT}/\omega_0 = 5 \times 10^{-4}$. The dispersion on the Λ line is characterized by a giant anticrossing between the branches of bare transverse photon and exciton modes. At the X point, the gap is determined by the separation between the longitudinal and lower transverse branches, it is still remarkable and exceeds $0.5\omega_{LT}$. However near the points U and W the exciton-polariton branches converge and the gap almost disappears. Note that the anticrossing can be described with a high accuracy by retaining in the sum over $\mathbf{b}$ in Eq. (11) the two terms due to $\mathbf{b} = 0, -(4\pi/a)(0, 0, 1)$ for the Δ points and $\mathbf{b} = 0, -(2\pi/a)(1, 1, 1)$ for the Λ points.

The 3D QD arrays with periods comparable with the light wavelength ($P \approx 1$)

and with sizes exceeding the bulk-exciton Bohr radius could be grown artificially or by embedding semiconductor microcrystals into the pores of porous materials like the synthetic opal [3]. It should be mentioned that the developed theory takes into account a contribution of only one exciton resonance which is valid if the separation between the exciton size-quantization levels is much larger than the bulk value of the exciton longitudinal-transverse splitting, ω_{LT} . In the opposite limit of extremely large bulk-exciton translational effective mass one can use the local material relation $\mathbf{D}(\mathbf{r}) = \varepsilon(\mathbf{r}, \omega)\mathbf{E}(\mathbf{r})$ as it was done by Sigalas et al. [4] for phonon-polaritons in a two-dimensional lattice consisting of semiconductor cylinders.

E. L. I. acknowledges partial support from Russian Foundation of Basic Research.

References

- [1] D. V. van Coevorden, R. Sprik, A. Tip, and A. Lagendijk, *Phys. Rev. Lett.* **77** 2412 (1996).
- [2] E. L. Ivchenko and A. V. Kavokin, *Fiz. Tverd. Tela* **34** 1815 (1992) [*Sov. Phys. Solid State* **34** 968 (1992)].
- [3] Yu. A. Vlasov, V. N. Astratov, O. Z. Karimov, A. A. Kaplyanskii, V. N. Bogomolov and A. V. Prokofiev, *Phys. Rev. B* **55** 13357 (1997).
- [4] M. M. Sigalas, C. M. Soukoulis, C. T. Chan, and K. M. Ho, *Phys. Rev. B* **49** 11080 (1994).