

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP013955

TITLE: Wave Diffraction by Axially Symmetrical System of Finite Soft
Cylinders

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: 2002 International Conference on Mathematical Methods in
Electromagnetic Theory [MMET 02]. Volume 2

To order the complete compilation report, use: ADA413455

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP013889 thru ADP013989

UNCLASSIFIED

WAVE DIFFRACTION BY AXIALLY SYMMETRICAL SYSTEM OF FINITE SOFT CYLINDERS

Eylem Özkan¹, Fatih Dikmen¹, Yury A. Tuchkin^{1,2} Sergey I. Tarapov^{1,2}

1. Gebze Institute of Technology, PK. 141, 41400 Gebze - Kocaeli, Turkey.

2. Institute of Radiophysics and Electronics, NAS of the Ukraine, 12 Ac. Proscura St., Kharkov 61085, Ukraine.

ABSTRACT

A new strong mathematically rigorous and numerically efficient method for solving the boundary value problem of scalar wave diffraction by a system of infinitely thin circular cylindrical screens is proposed. The method is based on a combination of Orthogonal Polynomials Method [1-2] and Analytical Regularization Method as used in [3,4,5]. The solution is generalization of the investigation done for one cylinder [6] and the method has been demonstrated on flat soft circular ring [6,7,8]. As a result of the suggested regularization procedure, the initial boundary value problem was equivalently reduced to the infinite system of the linear algebraic equations of the second kind, i.e. to an equation of the type $(I + H)x = b$, $x, b \in l_2$ - in the space l_2 of square summable sequences. This equation can be solved numerically by means of truncation method with, in principle, any required accuracy. Pilot experiments show good perspective of such cylindrical reflector for development of individual antenna tag for rescue radar or broadcast systems in mm waveband.

Let surface S have the following property,

$$S = \bigcup_{j=1}^N S_j, S_j \cap S_{j+1} = \emptyset \quad (1)$$

S is a system of finite circular cylinders located on z -axis defining, (Figure 1)

$$S_j = \{(z, \rho, \varphi) : z \in [\zeta_j - L_j, \zeta_j + L_j], \rho = a_j, \varphi \in [-\pi, \pi]\}, j=1, 2, \dots, N. \quad (2)$$




Figure 1

The following integral equation of the first kind is equivalent to the diffraction problem posed.

$$\int_S J_D(p) G(q, p) ds_p = -u^i(q); q \in S \quad (3)$$

where, $u^i(q)$ is known incident wave, $J_D(p)$ is unknown function i.e. current density like, $J_D(p) = [d(p)]^{1/2} H(p)$, $p \in S$, where $H(p)$ is a smooth function on surface S , $d(p)$ is the distance to the nearest edge of a ring, $G(q, p)$ is the Green's function of free space.

The axially symmetry of the system of obstacles leads to an infinite system of one dimensional, non-interacting integral equations of the first kind below,

$$2\pi \sum_{j=1}^N a_j \int_{\zeta_j-L_j}^{\zeta_j+L_j} Z_m^j(z_p) G_m^{j,l}(|z_p - z_q|) dz_p = g_m^l(z_q), \quad z_q \in [L_l, L_l], \quad l=1,2,\dots,N, \quad (4)$$

in terms of Fourier coefficients.

Proper parametrization of the variable in the equation is required to use the Fourier-Chebyshev expansions defined on the interval $[-1,1]$. The parametrization to reduce the points on each S_j surface to such an interval is the following:

$$z_p^j = \eta^j(v) = \zeta_j + L_j v, \quad v \in [-1,1]; \quad (5)$$

$$z_q^l = \eta^l(u) = \zeta_l + L_l u, \quad u \in [-1,1]. \quad (6)$$

Then we have (4) reduced to:

$$\int_{-1}^1 \left\{ -\frac{1}{\pi} \ln|u-v| + K_m^{l,l}(u,v) \right\} \tilde{Z}_m^l(v) dv + \sum_{j=1, j \neq l}^N \int_{-1}^1 \tilde{G}_m^{j,l}(u,v) \tilde{Z}_m^j(v) dv = \tilde{g}_m^l(u);$$

$$u \in [-1,1], \quad l=1,2,\dots,N \quad (7)$$

Here $K_m^{l,l}$ are sufficiently and $\tilde{G}_m^{j,l}$ are infinitely smooth functions and it is possible to express those functions and the unknown function $\tilde{Z}_m^j(v)$ using Fourier-Chebyshev series.

Therefore the final algebraic system of the first kind that (5) will be reduced to is

$$\gamma_n^{-2} z_n^l + \sum_{s=0}^{\infty} \left[k_{ns}^{l,l} z_s^l + \sum_{\substack{j=1 \\ j \neq l}}^N k_{ns}^{j,l} z_s^j \right] = b_n^l, \quad n=0,1,2,\dots \quad l=1,2,3,\dots,N \quad (8)$$

and will be subject to analytical regularization in the same manner done for single obstacles (as done in [6-7-8]), easily by introducing new variables $\hat{z}_n^l = z_n^l / \gamma_n$ and multiplying each term in (8) by γ_q . Here, $\gamma_0 = (\ln 2)^{-1/2}$; $\gamma_n = |n|^{1/2}$, $n \neq 0$, for every $m=0,\pm 1,\pm 2,\pm 3,\dots$ z_n^l and z_n^j are the Fourier - Chebyshev coefficients of the unknown function, b_n^l is Fourier - Chebyshev coefficients of the excitation term, $k_{ns}^{j,l}$ and $k_{ns}^{l,l}$ are Fourier - Chebyshev coefficients of the smooth kernels in (7).

Numerical results of the system $ka_1=2$, $ka_2=6$, $ka_3=10$, $kL_1=kL_2=kL_3=4$ in case of a normally incident plane wave, are following. In figure 4 the approximate locations of the surfaces are indicated.




Figure 2: Current Density




Figure 3: Far Field




Figure 4: Near Field

- [1] G.Ya.Popov. On one approximate method for solving the integral equation of wave diffraction by finite width strip - Zhurnal tekhnicheskoy fiziki, 1965, v. 35, n. 3, p.p. 381-389 (in Russian).
- [2] G.Ya.Popov. On Orthogonal polynomials method in contact problem of the elasticity theory - Prikladnaya matematika i. mekhanika, 1969, V. 33, N3 (in Russian)
- [3] Yu.A.Tuchkin. Wave scattering by unclosed cylindrical screen of arbitrary profile with Dirichlet boundary condition. - Soviet Physics Doclady, 1985, v. 30, p.p. 1027-1030
- [4] Yu.A.Tuchkin. Wave scattering by unclosed cylindrical screen of arbitrary profile with Neumann boundary condition. - Soviet Physics Doclady, 1987, v. 32, p.p. 213-216
- [5] Yu.A.Tuchkin. Regularization of boundary value problem of wave diffraction by toroidal screen of arbitrary profile - in the book: Electrodynamics of open structure of millimetre and sub-millimetre wave range. - Publishing house of IRE Acad. Sci. The Ukr. SSSR.Kharkov, 1990 (in Russian)
- [6] Fatih Dikmen, Eylem Özkan, Yury A. Tuchkin. Scalar Wave Diffraction from infinitely thin perfectly soft finite-length circular cylinder. Day On Diffraction 2001, International Seminar, St. Petersburg, May 29-31, 2001.
- [7] F. Dikmen, E. Karacuha, Yu.A. Tuchkin. Scalar Wave Diffraction by a Perfectly Soft Infinitely Thin Circular Ring. Turkish Journal of Electrical Engineering and Computer Sciences "Elektrik", Vol:9 No:21, 2001.
- [8] Yu.A. Tuchkin, E. Karacuha, F. Dikmen. Scalar Wave Diffraction from Infinitely Thin Circular Ring, Proc. of International Symposium on Mathematical Methods in Electromagnetic Theory - MMET'98, Kharkov, Ukraine.
- [9] E. Özkan, Fatih Dikmen, Eylem Özkan, Yury A. Tuchkin. Scalar Wave Diffraction by Perfectly Soft Thin Circular Cylinder of Finite Length ; Analytical Regularization. Turkish Journal of Electrical Engineering and Computer Sciences "Elektrik"(accepted, will be published).